


This article was downloaded by: [National Taiwan University]

On: 11 September 2008

Access details: Access Details: [subscription number 788856278]

Publisher Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Applied Financial Economics

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title-content=t713684415>

On the use and improvement of Hull and White's control variate technique

San-Lin Chung ^a; Mark B. Shackleton ^b

^a Department of Finance, National Taiwan University, Taipei, Taiwan, ROC ^b Department of Accounting and Finance, Lancaster University, UK

Online Publication Date: 01 November 2005

To cite this Article Chung, San-Lin and Shackleton, Mark B.(2005)'On the use and improvement of Hull and White's control variate technique',Applied Financial Economics,15:16,1171 — 1179

To link to this Article: DOI: 10.1080/09603100500359195

URL: <http://dx.doi.org/10.1080/09603100500359195>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

On the use and improvement of Hull and White's control variate technique

San-Lin Chung^{a,*} and Mark B. Shackleton^b

^a*Department of Finance, National Taiwan University, Taipei, Taiwan, ROC*

^b*Department of Accounting and Finance, Lancaster University, UK*

A study of the use and improvement of Hull and White's (1988) control variate technique in pricing options is provided. It contributes to the literature in two ways. First it is shown that it is not optimal to use the entire error of a control variate against its known price (usually a closed-form solution) to correct and improve the unknown error of the unknown price of a complex option and a better error correction fraction is derived. Secondly, while Hull and White only advocated the use of the simplest European option control variate, it is shown how to choose better controls to reduce pricing errors more effectively and the role of so called static hedges as the best theoretical control variates is discussed.

I. Introduction

Many problems in finance must ultimately be solved through the use of a numerical technique as the only practical and universally robust solution method. By numerical technique, we mean that the state space in question must be partitioned and discretized in some manner (either using a tree or a grid) before the value of the asset is evaluated at each and every of the many resultant points. Moreover these points (tree nodes or grid points) have to be chosen before the prices at those points are known.

In the field of options and particularly for American options¹ (where no closed form solution is generally available), such a numerical brute force method is necessary because it is the only way to guarantee that increasing computational effort results in increased numerical accuracy. Other areas where numerical techniques are necessary include all

dynamic programming techniques where the value of a strategy must be evaluated simultaneously with its optimal action, for instance in optimal portfolio control under transaction costs and similar problems.

It is important to be able to determine the price of a financial claim to arbitrary accuracy, first because a financial intermediary may be concerned about overpaying or undercharging. Secondly and more importantly, because the financial intermediary is often left with the task of replicating the claim over its life, he must be able to evaluate the comparative statics (the hedge parameters or so called Greeks of the option) repeatedly and quickly if he is to respond to revised hedging needs. For exotic options (especially path dependent options), this task may be sensitive to the method chosen and remaining errors will be a function of the computational time spent (see Boyle and Lau, 1994; Figlewski and Gao, 1999).

*Corresponding author. E-mail: chungsl@management.ntu.edu.tw

¹ Asian options and other exotics also fall into this category.




Fig. 1. The American put option price with regard to the number of time steps (binomial method): (—) no CV, (—) c^* , (---) HW($c=1$), (—) true value

The parameters are $S=40$, $K=45$, $T=7$ months, $r=0.07$, $q=0.02$, and $s=0.2$. The coefficient c^* is estimated using binomial trees with 20 to 50 steps. The accurate American option price is estimated using the BBSR method with 10 000 steps.

Although several approximations² are available for American options (also for Asians), many of which have the right asymptotic properties and converge to the true price in the limit of one variable, these do not allow the price of a general option to be determined to arbitrary accuracy. Even if a series solution is actually available for pricing an option to arbitrary accuracy (e.g. Geske and Johnson, 1984), it may still be more efficient³ to employ a numerical method compared to the task of evaluating a complex series. This is because the required multivariate cumulative normal function (and other high dimensional numerical integrations on which many compound options rely) although expressed as an integral and computer coded very efficiently, is not strictly a closed form solution in that it requires a numerical method itself for computation. Arbitrary accuracy is only achieved through increasing computational power and so other methods are typically preferred to the multivariate cumulative normal function.

Furthermore, the numerical methods themselves are subject to various problems. The rate of convergence of the estimated price to the true price, may not be fast and the convergence may not be monotone in that the sign and magnitude of the error may oscillate as a finer time grid or tree is used (see Fig. 1). This is because there are two sources of discretization error.

First, discretization at the terminal boundary will yield imperfect prices. Interpolation techniques and trapezoidal integration rules can minimize these errors for smooth payoff functions but the first derivative at the final exercise point of an option may not be smooth so the grid or lattice density here is critical.⁴

Secondly, the chosen numerical array must be used to evaluate the point of optimal action simultaneously with the desired prices. This will involve further forms of discretization error, one in determining the location of the optimal exercise points itself and one when subsequently valuing the option earlier in time, which is dependent on the value at, and position of, the optimal exercise boundary (Wilmott *et al.* (1995) detail a complimentary condition that can reduce this boundary error). This will cause problems particularly if the American exercise boundary is far from linear or if it is discontinuous at maturity.

This is to say that the critical points in the option pricing problem (that coincide with an optimal exercise boundary) are as important as the option prices away from the boundary. Indeed the boundary conditions have equal standing with the asset partial differential equation itself in pricing the claim because it is the boundary conditions alone that distinguish one type of option claim from another. It is the (prior) positioning of the nodes in state space relative to these critical thresholds in the option pricing problem that cause the second type of pricing error. Furthermore it is the non-linear way that the state space grid or tree interacts with a boundary of general shape that causes these errors to depend chaotically⁵ on the grid size (see Fig. 1). A small error in the

² Approximate solutions for American options include the work of Geske and Johnson (1984), MacMillan (1986), Barone-Adesi and Whaley (1987), Omberg (1987), and Ju (1998) etc. Although quick to evaluate, none of these methods will allow accuracy to be increased with more computational effort.

³ Here, efficient means that the method takes less computational time for the same accuracy or gives more accurate estimates for the same computational time.

⁴ Figlewski and Gao (1999) propose a method termed Adaptive Mesh Model (AMM) to reduce the 'non-linearity error' at the terminal boundary. In their method, one or more small sections of fine high-resolution lattice are added onto a tree with coarser time and price steps. This will incorporate more accurate fine-mesh values for critical nodes into the coarse tree and thus reduce the non-linearity error.

⁵ A chaotic variable is one that could be evaluated exactly with sufficient computing power; without such resource it may appear to be 'random' (i.e. there may be no apparent short cut to its determination). Since the emphasis here is to use as little computational power as possible, these chaotic errors will be treated as if they were truly random (i.e. without deterministic form).

placement of the grid next to a boundary can lead to a large pricing error. By way of example, Boyle and Lau (1994) show that convergence is very slow when using binomial methods to price Barrier options because of the severe non-linearity present as the mesh or lattice points oscillate around the boundary with decreasing mesh size.

One way to reduce these types of errors and hence improve the computational time required is the so called 'control variate' method proposed by Hull and White (1988, hereafter HW). They suggested that a similar claim for which a closed form solution is available be priced on the same grid and made subject to the same discretization errors. Comparison of the resultant price with the analytic price is instructive for two reasons, first the magnitude of the errors against the known price may be related to the errors on the unknown price, secondly the sign of the two may be related. In their paper, the use of a Black-Scholes control spared the use of a finer grid and thus more computation.

In addition to HW, this paper contributes to the literature in two directions. First HW used 100% of the error in this control variate against its known price to correct and improve the unknown error of the unknown price. It will be shown that the full use of the control variable may not be optimal. In some cases using the *entire error* to correct the unknown error will increase rather than decrease the latter.

Secondly it is interesting to note that HW only advocated the use of a simple control variate, the corresponding Black and Scholes European option price. This will help deal with the first type of discretization error at maturity but will not necessarily help with the second source of error in the location of and value near the boundary. Other instruments are suggested that may well help in the second area.

The rest of this paper is organized as follows. Section II discusses the methodology and the sources of pricing errors when using discrete lattice methods. Section III presents the numerical results and suggests how to choose good controls to effectively correct errors. Section IV concludes.

II. Methodology

Numerical methods based on a finite number of calculations will always contain error, in particular for binomial methods only for an arbitrarily large

number of tree nodes will the binomial price converge to the true continuous price. That is for a given set of option parameters (e.g. stock, strike, risk free, volatility, maturity) $\theta = (S_0, X, r, \sigma, T)$ and number of time steps n implemented in the tree, the estimated option price $\hat{P}(\theta, n)$ will only converge to the true price $P(\theta) = \lim_{n \rightarrow \infty} \hat{P}(\theta, n)$ as $n \rightarrow \infty$. As mentioned previously, even then there is a problem since convergence to the limit may be oscillatory.

An option error $\varepsilon(n)$ can be defined for any given numerical method or choice of n , the magnitude of which should decrease as $n \rightarrow \infty$

$$\varepsilon(n) = \hat{P}(\theta, n) - P(\theta) \quad \lim_{n \rightarrow \infty} \varepsilon(n) = 0$$

The control variate (CV) technique is applicable when there are two similar⁶ options, labelled C and T . The Target option T is the (difficult) option which needs to be priced while C is a Control option, called a Control Variate (or CV), which is similar to option T but has an analytical (or closed-form) solution P_C . To apply the control variate technique, the values of T and C have to be evaluated using the same numerical procedure (i.e. with the same value of n if trees are used). The numerical price estimates of T and C are denoted as \hat{P}_T and \hat{P}_C , respectively and they depend on n . The control variate estimate for the value of T , \tilde{P}_T , is given by

$$\tilde{P}_T = \hat{P}_T + c(P_C - \hat{P}_C) \tag{1}$$

$$\varepsilon_C = \hat{P}_C - P_C \tag{2}$$

$$\varepsilon_T \approx c\varepsilon_C \tag{3}$$

$$\tilde{P}_T = \hat{P}_T - c\varepsilon_C \tag{4}$$

This is to say that the error of the control option ε_C can be used to improve the error of the target option ε_T because it may be closely linked to it.

Hull and White recommended the use of a corresponding Black-Scholes European option price as a control variable P_C for an American option and they then used the pricing error of the discretized price estimate on the European to improve the American, i.e. the *entire error* $c=1$ of the control variate to correct the unknown error of the price estimate of the target option.

Without any prior knowledge of the dependency between the target errors and the control errors, the $c=1$ assumption is probably best but examination of actual error dependency may yield a different error sensitivity and potentially a better use of the control. The idea is to always chose the value of that leads to

⁶ By similar options it is meant that both option prices should follow the same partial differential equation (e.g. European versus American options), both option payoffs should be highly correlated (geometric average versus arithmetic average Asian options) or that their boundary conditions should be congruent (continuously versus discretely monitored barrier options).

the minimum target error, at least over a limited range of estimations n .

This then leads to the question of how to estimate the dependency between the control and target errors. This could be examined across different methods, but within a particular numerical method such as binomial trees the errors are investigated as a function of different tree step numbers n indexed by i . This is always done for the same parameter set θ and generate option price estimates $\hat{P}(\theta, i)$ for many i . For a given set of target option inputs, it is not desirable to include control errors from other parameter input combinations θ' (since they may not respond/correlate in the same manner) so instead other numerical estimate errors are employed across different i but for the same θ . The idea is to use a relatively small number of time steps i between two reference levels l, m to determine the target/control error dependency quickly and then use this to apply to a much larger tree of size n that is only *evaluated once* ($l < m \ll n$).

Option prices for the control and target will be evaluated for different numbers of time steps between l, m inclusive (i.e. a total of $m - l + 1$) in order to determine the correlation coefficient and optimal control ratio. This optimal control ratio will then be applied to errors from the control and target calculated from a much larger tree with $n \gg m, l$ in order to obtain a more accurate estimate of the limiting value $n \rightarrow \infty$.

If it is desired to minimize the error variance of \tilde{P}_T then the optimal correction to use is

$$c^*(l, m) = \text{Cov}(T, C) / \text{Var}(C) = \rho(T, C) \sigma_T / \sigma_C \quad (5)$$

$$= \frac{\sum_{i=l}^m \varepsilon_C(i) \varepsilon_T(i)}{\sum_{i=l}^m \varepsilon_C^2(i)} \quad (6)$$

where $\rho(T, C)$ is the correlation coefficient between \hat{P}_T and \hat{P}_C , σ_T^2 (σ_C^2) is the variance of \hat{P}_T (\hat{P}_C), and c^* is the optimal coefficient in which case the variance of the estimate will be minimized at

$$\text{Var}(\hat{P}_T + c^*(P_C - \hat{P}_C)) = (1 - \rho(l, m)^2) \sigma_T^2 \quad (7)$$

a quantity that could be zero if the correlation were perfect.

Most importantly, it will not generally be the case that the HW choice of $c = 1$ will yield the best results, especially not if $\rho(T, C) \ll 1$. In fact if this is the case, the resulting error may well be greater if $c = 1$ is used,




Fig. 2. American against European option errors for tree steps between 20 and 100

and clearly a variate with $\rho(T, C) = 0$ cannot help improve pricing. This is the case in Fig. 1 where the American pricing error is plotted as a function of the number of grid points used. As can be seen the errors using $c = 1$ can actually be greater than using $c = 0$ (i.e. no control is used).⁷ Figure 2 shows a scatter graph of the American option pricing errors as a function of the European errors, i.e. a visualisation of the correlation and slope coefficients used in this paper, the dependency between the two is clear from this figure.

There are in fact two attributes of the control variate error that matter for improving target prices, one is the *magnitude* of its pricing errors and the second is the *correlation* of its errors with those on the unknown price of the target option.

This paper is concerned with the choice of control variate, the correlation of target errors and finally the magnitude of the errors that result from an optimal use of the CV technique. A range of control variates are examined, the correlation of their errors and their ability to enhance pricing.

It is proposed to evaluate the performance of CV techniques under $c = 1$ and $c^*(l, m)$ methods in order to evaluate the performance of the Black–Scholes (BS) control variate. It is also proposed to evaluate the performance of other control variates that may reduce discretization error near an exercise boundary. For example, the analytical approximations of Barone-Adesi and Whaley (1987), MacMillan (1986) (BAWM) and Omberg (1987) (Om) both allow determination of an approximate (but not optimal) exercise boundary that can be used to correct the discretization error near the true exercise boundary. For each and every array location, it is possible to say

⁷ Note that c^* can in fact be greater than one if the errors in T exceed the errors in C and the correlation is high, i.e. the variance of the target option price must be greater than that of the control price, a situation which may be possible for barrier options controlled with standard European options.

if it lies in or outside the boundary as approximated by each of the formulae. Thus the evaluation of these methods under grid pricing and closed form may yield extra information about the nature of the discretization error near a boundary that is a reasonable approximation for the true boundary. The performance of these new forms compared to existing control variates will elucidate something of the nature of the discretization errors that are incurred at different points of the grid.

For each CV (European, Omberg options etc.) the optimal $c^*(l, m)$ is established as follows. The prices of the control option and the target option are calculated using one lattice method, e.g. simple binomial method, for a reasonable number of time steps, say $l=20$ to $m=50$ steps. Then two series of prices are obtained which allow us to estimate their variance-covariance and hence $c^*(l, m)$. A much larger tree ($n=10\,000$) is then used and the prices it generates for both target and control along with the optimal $c^*(l, m)$ use of these controls.

III. Numerical Results

The most efficient use of a control variate is not necessarily $c=1$, for a given step size estimation and use of c^* will generally improve price estimates. However it takes time to estimate c^* , therefore it is important that c^* can be estimated using only a few samples (Monte Carlo simulation) or time steps (binomial method) we estimate c^* as follows. The prices of the target option and the control option are calculated using binomial trees using different numbers of time steps, for example, $l=20$ to $m=100$, this generates two series of prices which allow $c^*(l, m)$ to be estimated. In Table 1, the target options are American put options and the control options are their counterpart European options. Table 1 shows that the estimates of c^* are quite similar for different ranges of time steps. In other words, c^* can be estimated using only few (e.g. 20–40) time steps but apply it for trees with more time steps. Therefore it is beneficial to use c^* rather than the

Table 1. Estimated optimal control $c^*(l, m)$ values using different sets of time steps

K	T (months)	σ	$c^*(l, m)$ and $\rho(l, m)$			
			(20, 40)	(20, 100)	(150, 200)	(250, 300)
35	1	0.2	0.96 (0.999)	0.99 (0.999)	0.96 (0.998)	0.95 (0.999)
35	1	0.3	0.95 (0.999)	0.98 (0.999)	0.96 (0.999)	0.95 (0.999)
35	1	0.4	0.95 (0.999)	0.98 (0.999)	0.96 (0.999)	0.95 (0.999)
35	4	0.2	0.90 (0.990)	0.92 (0.989)	0.90 (0.993)	0.87 (0.994)
35	4	0.3	0.88 (0.989)	0.92 (0.990)	0.88 (0.995)	0.90 (0.996)
35	4	0.4	0.88 (0.993)	0.92 (0.992)	0.92 (0.997)	0.90 (0.997)
35	7	0.2	0.83 (0.984)	0.86 (0.983)	0.82 (0.991)	0.81 (0.994)
35	7	0.3	0.82 (0.985)	0.87 (0.985)	0.87 (0.993)	0.84 (0.994)
35	7	0.4	0.87 (0.990)	0.90 (0.990)	0.86 (0.995)	0.88 (0.997)
40	1	0.2	0.81 (0.999)	0.84 (0.995)	0.84 (1.000)	0.84 (1.000)
40	1	0.3	0.86 (0.999)	0.89 (0.997)	0.88 (1.000)	0.88 (1.000)
40	1	0.4	0.88 (1.000)	0.91 (0.998)	0.91 (1.000)	0.91 (1.000)
40	4	0.2	0.70 (0.998)	0.73 (0.991)	0.73 (1.000)	0.73 (1.000)
40	4	0.3	0.77 (0.998)	0.80 (0.993)	0.80 (1.000)	0.80 (1.000)
40	4	0.4	0.81 (0.999)	0.84 (0.995)	0.84 (1.000)	0.84 (1.000)
40	7	0.2	0.64 (0.998)	0.66 (0.991)	0.66 (1.000)	0.66 (1.000)
40	7	0.3	0.72 (0.999)	0.75 (0.992)	0.75 (1.000)	0.75 (1.000)
40	7	0.4	0.77 (0.998)	0.79 (0.994)	0.79 (1.000)	0.79 (1.000)
45	1	0.2	0.00 (0.000)	0.00 (0.000)	0.00 (0.000)	0.00 (0.000)
45	1	0.3	0.51 (0.967)	0.51 (0.946)	0.57 (0.994)	0.57 (0.996)
45	1	0.4	0.79 (0.984)	0.81 (0.970)	0.79 (0.996)	0.77 (0.997)
45	4	0.2	0.36 (0.931)	0.37 (0.933)	0.35 (0.977)	0.33 (0.974)
45	4	0.3	0.63 (0.972)	0.68 (0.953)	0.67 (0.987)	0.67 (0.993)
45	4	0.4	0.77 (0.973)	0.82 (0.965)	0.78 (0.993)	0.75 (0.991)
45	7	0.2	0.40 (0.899)	0.41 (0.918)	0.41 (0.976)	0.38 (0.983)
45	7	0.3	0.64 (0.955)	0.68 (0.948)	0.65 (0.987)	0.64 (0.981)
45	7	0.4	0.75 (0.987)	0.79 (0.976)	0.74 (0.995)	0.68 (0.989)

Notes: The options are puts with other parameters: $S=40$, $r=0.0488$, $q=0.0$ (dividend yield). The bracket (e.g. 20, 40) means that c^* and ρ are estimated using binomial trees with step numbers from 20 to 40. The correlation coefficients (ρ) are given in parentheses.

Table 2. The root-mean-squared (RMS) relative errors using the following four methods to price European options

Binomial	BBS	BBSR	AMM
4.59E-05 = 0.459bp	5.41E-05 = 0.541bp	2.46E-06 = 0.0246bp	2.98E-04 = 2.98bp

Note: The options are European puts. The root-mean-squared relative errors are defined as follows

$$\text{RMS} = \left(\frac{1}{m} \sum_{j=1}^m e_j^2 \right)^{1/2}$$

where $e_j = (P_j^* - P)/P_j$ is the relative error, P_j is the accurate option price (Black–Scholes), P_j^* is the estimated option price. The number of time steps in each method is $n = 10\,000$. There are 243 sets of parameters are used: $\sigma = 40$, $K = 35, 40, 45$, $s = 0.2, 0.3, 0.4$, $T = 1, 4, 7$ months, $r = 3, 5, 7\%$, and $q = 2, 5, 8\%$.

prior of $c = 1$ because it takes little computational time to estimate it.

To compute the benchmark values of the American options used in validating the approximation methods, the accuracy of four lattice approaches is first investigated; the binomial method of Cox *et al.* (1979), the binomial Black and Scholes (BBS) method and the binomial Black and Scholes method with Richardson extrapolation (BBSR) of Broadie and Detemple (1996),⁸ and the Adaptive Mesh Model (AMM) of Figlewski and Gao (1999) (see footnote 4). The procedure in Broadie and Detemple (1996, 1997) is followed to run a comparison. First a large set of options ($3^5 = 243$ options; 3 values of each of 5 variables) is chosen with parameters of practical interest (values from other work). Then for each method the test set of options is priced and percentage error measures computed. The root-mean-squared (RMS) relative percentage error is used as the error measure and is defined by:

$$\text{RMS} = \left(\frac{1}{T} \sum_{j=1}^T e_j^2 \right)^{1/2} \quad (8)$$

where $T = 243$ is the number of options under consideration, $e_j = (\widehat{P}_j(\theta_j) - P_j(\theta_j))/P_j(\theta_j)$ is the relative error⁹ of the j th option, $P_j(\theta_j)$ is the true option price (Black–Scholes), $\widehat{P}_j(\theta_j)$ is the estimated option price.

The results (Table 2) indicate that the BBSR method produces the most accurate values of these four methods. Therefore, the BBSR with $n = 10\,000$

time steps is used to produce benchmark values for American options in the following numerical evaluations.

Optimal $c^*(l, m)$ is defined as the coefficient of the target and control option estimated from using the binomial prices with few time steps, l to m ($l < m$). Treating this numerical experiment as a statistical experiment when evaluating pricing errors for a number of tree steps between l, m , terminology is borrowed and $c^*(l, m)$ labelled as an *in sample coefficient*. This coefficient is used as a proxy for the unknown *out of sample coefficient* $c^*(l, n)$ and the errors used in making this assumption are evaluated.

Table 3 shows that the CV technique can significantly reduce the RMS relative errors of price estimates for the binomial method. The CV technique using c^* improves the RMS relative error by about 20 to 25 times for the binomial method. Combining with the results (their Figs 2 to 8) of Broadie and Detemple (1996), it can be found that the binomial method coupled with the CV technique using c^* seems more efficient than other proven (numerically) efficient methods such as the BBSR method. Moreover, Table 3 suggests that the CV technique using c^* works well for both in-sample and out-of-sample tests.¹⁰

Another objective of the CV technique is actually to seek a better instrument (with closed or analytic form pricing solution) whose numerical errors have as high a correlation as possible with the target option prices errors and then to use these as corrections.

⁸The BBS method is a modification of the binomial method where the Black–Scholes formula replaces the usual ‘continuation value’ at the last time step just before option maturity. The BBSR method adds Richardson extrapolation to the BBS method. In particular, the BBSR method with n steps computes the corresponding BBS price for $n/2$ steps $P(\theta, n/2)$ and n steps $P(\theta, n)$ and then sets the BBSR approximation to the asymptotic price equal to $P(\theta, \infty) = 2P(\theta, n) - P(\theta, n/2)$.

⁹Note that these *percentage* errors are calculated using the same number of steps ($n = 10\,000$) but across *different* parameter sets j i.e. $\% e(j) = (\widehat{P}(\theta(j), i) - P(\theta(j), i))/P(\theta(j), i)$ as oppose to the previous absolute errors which were calculated for the same parameter set θ but for different number of tree steps i i.e. $\epsilon(i) = \widehat{P}(\theta, i) - P(\theta, i)$ for $i = l$ to m .

¹⁰Because c^* is estimated using binomial trees with 20 to 100 time steps, 30, 50, and 80 time steps are called in-sample tests, and 200, 400, and 600 time steps out-of-sample tests.

Table 3. The root-mean-squared (RMS) relative errors of price estimates

	In-sample test			Out-of-sample test		
	30	50	80	200	400	600
no CV	1.74E-02 174bp	1.46E-02 146bp	6.38E-03 63.8bp	2.38E-03 23.8bp	1.88E-03 18.8bp	8.11E-04 8.11bp
$c = 1$	1.14E-03 11.4bp (93.4%)	6.86E-04 6.86bp (95.3%)	5.26E-04 5.26bp (91.8%)	1.83E-04 1.83bp (92.3%)	9.98E-05 0.998bp (94.7%)	6.45E-05 0.645bp (92.0%)
c^*	7.43E-04 7.43bp (95.7%)	4.36E-04 4.36bp (97.0%)	3.80E-04 3.80bp (94.0%)	1.19E-04 1.19bp (95.0%)	7.82E-05 0.782bp (95.8%)	4.77E-05 0.477bp (94.1%)

Note: The options are American puts. The root-mean-squared relative errors are defined as follows:

$$RMS = \left(\frac{1}{m} \sum_{j=1}^m e_j^2 \right)^{1/2}$$

where $e_j = (P_j^* - P_j)/P_j$ is the relative error, P_j is the accurate American option price (estimated by the BBSR method with 10 000 steps), P_j^* is the estimated option price. There are 243 sets of parameters are used: $S=40$, $K=35, 40, 45$, $\sigma=0.2, 0.3, 0.4$, $T=1, 4, 7$ months, $r=3, 5, 7\%$, and $q=2, 5, 8\%$. The coefficient c^* is estimated using trees with 20 to 100 steps. The numbers in parentheses are the error reduction percentages using the control variate technique with c^* and $c = 1$.

To date the CV technique has used Black–Scholes (since its errors are positively correlated because a deep out of the money American, is priced as a European anyway) but other closed form formulae or indeed approximating formulae could be used instead. For instance Omberg (1987) has a closed form expression for American option prices assuming an exponential boundary (since this boundary is not actually optimal the formula will undervalue options), this boundary can be numerically applied and a binomial price for Omberg compared to the theoretical Omberg price. *It may not matter that the Omberg price is not actually an optimal price of an option,¹¹ what matters is that its errors may correlate very well with the true American price errors.¹²* This is so because its arbitrary boundary does not get too far from the true one and it will suffer errors in the same regions as the true American (near the boundary).

To verify the above argument, the performance of European options were compared to those of Omberg options. Table 4 shows that the average correlation coefficients between American options and

Omberg options are high (0.974). However, it is a little surprising that the performance of Omberg options¹³ as the controls is only slightly better than European options. A possible explanation is that the oscillating binomial prices may be not able to distinguish a good control from a bad control.¹⁴ Overall, the results still suggest that an option with wisely¹⁵ chosen exercise boundary is a good control for American options but that the final maturity boundary errors due to grid point positioning still matter.

Note that the control can be an option or a portfolio of options and theoretically the best control option(s) are those that have the highest correlation coefficient with the price of the target option. Recently the *static hedge* literature¹⁶ has been used to replicate an exotic option using a *portfolio* of standard (closed form) options. Since an effective static hedge portfolio can approximately replicate the option position, it will also provide an effective control to reduce the variance of the numerical estimates of option prices.¹⁷

¹¹ Although it corresponds to the price of a claim with an arbitrary, but optimized, boundary shape.

¹² The option bias between the actual and binomial Omberg prices should be small.

¹³ In terms of percentage the absolute values of correlation coefficients of Omberg options are still generally (65.1%) higher than those of European options.

¹⁴ One evaluation criterion to determine a good numerical method may be its ability to distinguish good and bad controls. From this point of view, the binomial method may fail in some cases.

¹⁵ Meaning that the arbitrary boundary is very close to the true exercise boundary, for example, Ju (1998) proposes a multipiece exponential exercise boundary which can approximate the true boundary much better than Omberg (1987).

¹⁶ The basic principle behind static hedge method is as follows. If the prices of the target option and the static hedge portfolio follow the same partial differential equation and they are worth the same on a certain boundary, they are also worth the same at all interior points of the boundary (see Derman *et al.*, 1995 or Carr *et al.*, 1998 for details).

¹⁷ However it may be very difficult and complicated to formulate a perfect static hedge portfolio in practice thus reducing the benefits of the CV method.

Table 4. The absolute value of correlation coefficients between the American option prices and the control option prices

	European option	Omberg option
mean	0.97246	0.97405
max	0.99967	0.99973
min	0.29175	0.32823

Notes: The options are American puts. There are 180 sets of parameters are used: $S=40$, $K=30$ to 50 in increments of 5, $\sigma=0.2, 0.4, 0.6$, $T=6, 12, 18, 24$ months, $r=3, 5, 7\%$, and $q=0$. The correlation coefficient is estimated using binomial trees with 60 to 100 steps. The first control is the counterpart European option with maturity date T . The second control is the counterpart Omberg option which has an exponential exercise boundary.

Table 5. The root-mean-squared (RMS) relative errors for the barrier option

	No control	Portfolio 1	Portfolio 2	Portfolio 3
RMS	6.66E-03	5.79E-03	4.34E-03	2.42E-03
	66.6bp	57.9bp	43.4bp	24.2bp

Note: The barrier option is an up-and-out call with $S=100$, $K=100$, $H=120$, $T=1$ year, $r=0.1$ (annually compounded), $q=0.05$ (annually compounded), and $\sigma=0.25$. Portfolio 1 is the counterpart European call option. Portfolio 2 consists of long one European call option with $K=100$, short one European call option with $K=H$, and short one cash-or-nothing (digital) call option with $K=H$ and cash $Q=H-100$. Portfolio 3 is a static hedge portfolio from Derman *et al.* (1995) (their Portfolio 4). The root-mean-squared relative errors are defined as follows:

$$\text{RMS} = \left(\frac{1}{m} \sum_{j=1}^m e_j^2 \right)^{1/2}$$

where $e_j = (P_j^* - P_j)/P_j$ is the relative error, P_j is the accurate barrier option price (closed-form solution), P_j^* is the estimated option price using n_i steps per half-year, where n_i equals to 250 to 300 steps.

As an example, the performance of three controls which partially match the boundary conditions of a barrier option were investigated. The barrier option considered in this paper is an up-and-out call option with $S=100$, $K=100$, $H=120$ (barrier), $T=1$ year, $r=0.1$ (annually compounded risk free rate), $q=0.05$ (annually compounded dividend yield), and $\sigma=0.25$ (volatility). The first control (1) is the counterpart European option which exactly matches

the boundary if the barrier is never crossed before maturity. The second control (2) matches the boundary of the barrier option at maturity. The third control (3) replicates the value of the barrier option; (a) at expiration below the barrier, and (b) exactly on the 120 barrier at one year and 6 months prior to expiration (half way). It is obvious that portfolio (3) should be a better control than portfolio (1) because it can match more nodes on the boundary of the barrier option. Table 5 indicates that the third control is the best control and can reduce the RMS relative errors by a factor of about three.

Following Derman *et al.* (1995), other controls can easily be found which match as many nodes on the boundary as desired.¹⁸ In other words, theoretically the static hedge portfolio can be used to find a perfect control which can correct all the errors although it will involve an increasing number of terms.

IV. Conclusion

This paper has considered the use and improvement of Hull and White's (1988) control variate technique in option pricing. First the sources of pricing errors when using discrete lattice method were discussed and good controls that effectively correct target errors recommended. Secondly, examples have been given that showed where Hull and White's control variate technique is not optimal and potentially it may increase rather than decrease the pricing errors.

To effectively reduce pricing errors, the control variate technique has to be customized to both the type of numerical methods and the options in question. The benefits of this effort are reduced errors and a deeper understanding of their origin and control.

It would be interesting to study the possibility of combining recent advanced numerical methods with the control variate technique to price complex options efficiently. For example, Barraquand and Martineau (1995) and Longstaff and Schwartz (2001) have proposed a method for pricing multi-dimensional American options by Monte Carlo simulation. Given the flexibility for pricing exotic options under complex asset price processes, combining their methods with control variate technique may provide a powerful tool for both academics and practitioners.

¹⁸ Derman *et al.* (1995) have shown that a replicating portfolio using 24 options to match the zero boundary value on the barrier at half-month intervals can hedge the (continuous) barrier option very well.

References

- Barone-Adesi, G. and Whaley, R. E. (1987) Efficient analytic approximation of American option values, *Journal of Finance*, **42**, 301–20.
- Barraquand J. and Martineau, D. (1995) Numerical valuation of high dimensional multivariate American securities, *Journal of Financial and Quantitative Analysis*, **30**, 381–405.
- Boyle, P. and Lau, S. H. (1994) Bumping up against the barrier with the binomial model, *Journal of Derivatives*, **1**, 6–14.
- Broadie, M. and Detemple, J. (1996) American option valuation: new bounds, approximations, and a comparison of existing methods, *Review of Financial Studies*, **9**, 1211–50.
- Broadie, M. and Detemple, J. (1997) The valuation of American options on multiple assets, *Mathematical Finance*, **7**(3), 241–86.
- Carr, P., Ellis, K. and Gupta, V. (1998) Static hedging of exotic options, *Journal of Finance*, **53**, 1165–90.
- Cox, J., Ross, S. and Rubinstein, M. (1979) Option pricing: a simplified approach, *Journal of Financial Economics*, **7**, 229–63.
- Derman, E., Ergener, D. and Kani, I. (1995) Static option replication, *Journal of Derivatives*, **2**, 78–95.
- Figlewski, S. and Gao, B. (1999) The adaptive mesh model: a new approach to efficient option pricing, *Journal of Financial Economics*, **53**, 313–51.
- Geske, R. and Johnson, H. E. (1984) The American put option valued analytically, *Journal of Finance*, **39**, 1511–24.
- Hull, J. and White, A. (1988) The use of the control variate technique in option pricing, *Journal of Financial and Quantitative Analysis*, **23**, 237–51.
- Ju, N. (1998) Pricing an American option by approximating its early exercise boundary as a multipiece exponential function, *Review of Financial Studies*, **11**, 627–46.
- Longstaff, F. A. and Schwartz, E. S. (2001) Valuing American options by simulations: a simple least-squares approach, *The Review of Financial Studies*, **14**, 113–47.
- MacMillan, L. W. (1986) Analytic approximation for the American put option, *Advances in Futures and Options Research*, **1**, 119–39.
- Omberg, E. (1987) The valuation of American puts with exponential exercise policies, *Advances in Futures and Options Research*, **2**, 117–42.
- Wilmott, P., Howison, S. and Dewynne, J. (1995) *The Mathematics of Financial Derivatives: A Student Introduction*, Cambridge University Press, Cambridge.