

Synthesis and reactivity of ditungsten helical complex $W_2(CO)_6(\mu-Ph_2PC\equiv CPPH_2)_3$

Wen-Yann Yeh^{a,*}, Shie-Ming Peng^b, Gene-Hsiang Lee^b

^a Department of Chemistry, National Sun Yat-Sen University, Kaohsiung 804, Taiwan

^b Department of Chemistry, National Taiwan University, Taipei 106, Taiwan

Received 26 December 2002; received in revised form 14 January 2003; accepted 30 January 2003

Abstract

Reaction of $W(CO)_3(Me_3tach)$ ($Me_3tach=1,3,5$ -trimethyl-1,3,5-triazacyclohexane) with $Ph_2PC\equiv CPPH_2$ at room temperature affords a triply-bridged complex $W_2(CO)_6(\mu-Ph_2PC\equiv CPPH_2)_3$ (**1**) and a vinylidene complex $W_2(CO)_6(\mu-Ph_2PC\equiv CPPH_2)[\mu-C_4H(PPh_2)_3]$ (**2**). Compound **2** can be obtained by treating **1** with Me_3tach in dichloromethane. The crystal structures of **1** and **2** are determined by an X-ray diffraction study. The structure of **1** depicts a helical M_2L_3 framework with an idealized D_3 symmetry. The vinylidene group of **2** is not linear, with the $W=C=C$ bond angle of $158.6(4)^\circ$.

© 2003 Elsevier Science B.V. All rights reserved.

Keywords: Me_3tach ; Bis(diphenylphosphino)acetylene; 1,3,5-Trimethyl-1,3,5-triazacyclohexane

1. Introduction


The ‘coordination clusters’ assembling from mono-metallic complexes and multidentate bridging ligands currently attract much interest [1,2]. Frequently, the final shape of the self-assembled clusters can be predicted, which is not only defined through the metal coordination geometry but also through the orientation of the interaction sites in a given ligand [3–7]. Bis(diphenylphosphino)acetylene (dppa) is a potentially trifunctional ligand. The two phosphorus centers of dppa are normally coordinated to metals in advance of the acetylene group due to stronger net donor capability of the phosphine ligand compared with alkyne [8]. However, the rigidity of the linear $-C\equiv C-$ unit between two phosphorus centers forces the dppa ligand to form bonds between different metal atoms. Thus, a number of bimetallic complexes bridged by one [9,10], two [11] and three [12] dppa moieties are known. In this paper, we report the synthesis of a tris(μ -dppa) ditungsten helical complex and its reactivity.

2. Results and discussion

The reaction of $W(CO)_3(Me_3tach)$ and diphenylacetylene has been shown to give the tris(alkyne) complex $W(CO)(\eta^2-PhC\equiv CPh)_3$ [13]. However, treatment of $W(CO)_3(Me_3tach)$ with $Ph_2PC\equiv CPPH_2$ in dichloromethane solvent at room temperature for 2 days produces $W_2(CO)_6(\mu-Ph_2PC\equiv CPPH_2)_3$ (**1**) in 74% yield and $W_2(CO)_6(\mu-Ph_2PC\equiv CPPH_2)[\mu-C_4H(PPh_2)_3]$ (**2**) in 3% yield (Scheme 1). The reaction can be completed within 5 h in refluxing tetrahydrofuran (THF) but affording little **2**. The triply-bridged **1** forms a pale-yellow, air-stable crystalline solid which is only sparingly soluble in THF, CH_2Cl_2 and acetone. It is thermally stable in refluxing toluene ($110^\circ C$) and *n*-butyl ether ($142^\circ C$) without decomposition. The IR spectrum of **1** in the carbonyl region presents two strong absorptions at 1950 and 1866 cm^{-1} , consistent with a *facial* configuration for each $W(CO)_3$ unit [14]. The 1H -NMR spectrum of **1** displays only one type of phenyl groups, and the ^{31}P -NMR spectrum shows one sharp singlet at -1.75 ppm with appropriate ^{183}W satellites for the equivalent phosphorus atoms. These spectral data are in agreement with a D_3 symmetry for **1** in solution.

* Corresponding author. Fax: +88-6-7-52539908.

E-mail address: wenyann@mail.nsysu.edu.tw (W.-Y. Yeh).



Scheme 1.

Compound **1** can be treated like three-bladed propellers by looking at the molecule down the W(1)···W(2) axis, where a counterclockwise (Λ) and a clockwise (Δ) arrangement can result [15]. The crystalline material is achiral with each unit cell containing two molecules of each enantiomer. The ORTEP diagram for the Λ -enantiomer is shown in Fig. 1. The molecule consists of two facially substituted W(CO)₃ moieties bridged by three dppa ligands. Each tungsten atom is bonded to three terminal carbonyl groups with individual W–C–O

angles in the range 172(1)–177(1)°. The coordination about each tungsten atom is a distorted octahedral such that the C–W–C angles range from 84.3(5)° to 90.0(5)° and the P–W–P angles are in the range 91.2(1)–97.7(1)°. The W–P–C_(acetylene) angles show a significant deviation from the ideal tetrahedral with angles ranging from 118.2(4)° to 120.6(4)°, consistent with steric strains arising from *syn*-coordination of the dppa ligands. The P–C≡C–P backbones are slightly bowed as indicated by the P–C≡C angles (average 177(1)°). There is a helical




Fig. 1. Molecular structure of **1**. The phenyl groups have been artificially omitted, except the ipso carbon atoms, for clarity. Selected bond distances (Å): W(1)–C(1), 2.02(1); W(1)–C(2), 2.01(2); W(1)–C(3), 2.02(2); W(1)–P(1), 2.495(3); W(1)–P(3), 2.509(3); W(1)–P(5), 2.501(3); W(2)–C(4), 2.01(1); W(2)–C(5), 2.02(2); W(2)–C(6), 1.98(2); W(2)–P(2), 2.514(3); W(2)–P(4), 2.504(3); W(2)–P(6), 2.501(3); P(1)–C(7), 1.75(1); P(2)–C(8), 1.76(1); P(3)–C(9), 1.79(1); P(4)–C(10), 1.78(1); P(5)–C(11), 1.76(1); P(6)–C(12), 1.77(1); C(7)–C(8), 1.23(2); C(9)–C(10), 1.19(2); C(11)–C(12), 1.21(2). Selected bond angles (°): C(1)–W(1)–C(2), 86.6(5); C(1)–W(1)–C(3), 84.3(5); C(2)–W(1)–C(3), 88.8(5); P(1)–W(1)–P(3), 93.9(1); P(1)–W(1)–P(5), 95.7(1); P(3)–W(1)–P(5), 91.2(1); C(4)–W(2)–C(5), 90.0(5); C(4)–W(2)–C(6), 84.6(5); C(5)–W(2)–C(6), 86.0(5); P(2)–W(2)–P(4), 93.0(1); P(2)–W(2)–P(6), 92.8(1); P(4)–W(2)–P(6), 94.3(1); W(1)–P(1)–C(7), 118.9(4); W(1)–P(3)–C(9), 118.7(4); W(1)–P(5)–C(11), 118.8(4); W(2)–P(2)–C(8), 118.2(4); W(2)–P(4)–C(10), 120.6(4); W(2)–P(6)–C(12), 118.3(4); P(1)–C(7)–C(8), 178(1); P(2)–C(8)–C(7), 176(1); P(3)–C(9)–C(10), 176(1); P(4)–C(10)–C(9), 176(1); P(5)–C(11)–C(12), 178(1); P(6)–C(12)–C(11), 178(1).

twist of ca. 21° about the W(1)··W(2) axis, taking as the average of the torsion angles: P(1)–W(1)–W(2)–P(2) 19.3° , P(3)–W(1)–W(2)–P(4) 19.3° and P(5)–W(1)–W(2)–P(6) 24.7° . The related helical tris(dppa) complexes are Ni₂(CO)₂(dppa)₃ [16], [Ag₂(dppa)₃]²⁺ [17], Mo₂(CO)₆(dppa)₃ [12] and (BuTe)₂Cu₂(dppa)₃ [18]. There are substantially intramolecular aromatic contacts between the 12 phenyl groups, as shown in Fig. 2. In combination of these contacts should restrict free rotation of the phenyl groups to account for poor solubility of **1** in organic solvents. There is small void at the center, such that the W(1)··W(2) distance is 7.39 Å and the distances between C≡C centroids are 3.54 Å in average.

Compound **2** forms a purple-red, air-stable crystalline solid. It is soluble in common organic solvents. The IR spectrum in the carbonyl region displays four stretching absorptions, suggesting different environments for the two W(CO)₃ groups. The ¹H-NMR spectrum shows a very complex multiplet in the range 7.85–6.43 ppm for the phenyl protons and a multiplet of ¹H at 5.84 ppm for an alkene proton. The most striking feature is that the ³¹P-NMR spectrum presents five multiplets of about equal intensities at 57.8, 49.4, 31.8, –5.1 and –6.3 ppm with each accompanied by appropriate ¹⁸³W satellites. Because of the absence of diagnostic spectral features to reveal the structure of **2**, a single-crystal X-ray diffraction study was performed.

The molecular structure of **2**, shown in Fig. 3, consists of two *fac*-W(CO)₃ groups bridged by a dppa ligand and a tris(diphenylphosphino)vinylvinylidene species. The coordination about the W(2) atom resembles the parent compound **1**, while the W(1) atom is connected to three terminal carbonyls, two phosphines and a vinylidene ligand. The atoms C(7), C(8), C(9), P(2) and P(3) are

about on the same plane, while the C(9)–C(10) and C(7)–P(1) vectors are bent away from this plane by ca. 10° . The C(7)–C(8) and C(9)–C(10) lengths of 1.345(6) and 1.329(6) Å are characteristic of a carbon–carbon double bond, which are shorter than the C(8)–C(9) single-bond distance of 1.475(6) Å and longer than the C(11)–C(12) triple-bond distance of 1.214(6) Å. The C(10) atom appears to form a double-bond to the W(1) atom with the interatomic distance of 1.954(4) Å, which is comparable with those measured for the tungsten–carbene lengths in W(C₂Ph₂)(C₈Ph₈) (2.02 Å) [19] and W(CO)(C₂Ph₂)(C₈Ph₈) (2.01 Å) [20]. Noticeably, the vinylidene W(1)=C(10)=C(9) angle of $158.6(4)^\circ$ is significantly deviated from linearity by 21.4° , presumably due to steric strains inside the WPC₄ ring.

Compound **1** is stable in dichloromethane, but it decomposes slowly when Me₃tach is added into the solution to produce **2** as one of the products. Apparently the base ‘Me₃tach’, either being added or released from W(CO)₃(Me₃tach), might induce **1** to undergo sequential transformations, including C–C coupling, PPh₂ elimination and hydrogen abstraction, to give **2**, though the exact pathways are still unclear. There are three uncoordinated acetylene units in **1**. However, attempts to react **1** with Co₂(CO)₈ or Ni(COD)₂ gave no reactions. Hong et al. [21] have shown that treating Co₂(CO)₈ with dppa yields a tetracobalt cluster bonded to both the phosphine and the acetylene moieties. It is probably the intramolecular phenyl stacking (Fig. 2) that prevents metal atom(s) from binding the acetylene groups. We are currently investigating if a small metal cation can be trapped inside the cavity of **1** through π -interactions.

3. Experimental

3.1. General methods

All manipulations were carried out under an atmosphere of purified dinitrogen with standard Schlenk techniques. W(CO)₃(Me₃tach) (Me₃tach=1,3,5-trimethyl-1,3,5-triazacyclohexane) was prepared from W(CO)₆ as described in the literature [13]. Ph₂PC≡CPh₂ was purchased from Aldrich and used as received. Solvents were dried over appropriate reagents under dinitrogen and distilled immediately before use. Preparative thin-layer chromatographic (TLC) plates were prepared from silica gel (Kieselgel, DGF₂₅₄). Infrared spectra were recorded with a 0.1 mm path CaF₂ solution cell on a Hitachi I-2001 IR spectrometer. ¹H- and ³¹P-NMR spectra were obtained on a Varian VXR-300 spectrometer. Fast-atom-bombardment (FAB) mass spectra were recorded by using a VG Blotch-5022 mass spectrometer. Elemental analyses




Fig. 2. Space filling representation of the phenyl groups in **1** to show the intramolecular aromatic contacts.




Fig. 3. Molecular structure of **2**. The phenyl groups have been artificially omitted, except the ipso carbon atoms, for clarity. Selected bond distances (Å): W(1)–C(1), 2.040(5); W(1)–C(2), 2.088(5); W(1)–C(3), 1.990(5); W(1)–C(10), 1.954(5); W(1)–P(1), 2.495(1); W(1)–P(4), 2.535(1); W(2)–C(4), 1.985(5); W(2)–C(5), 1.983(5); W(2)–C(6), 1.982(6); W(2)–P(2), 2.509(1); W(2)–P(3), 2.500(1); W(2)–P(5), 2.525(1); P(1)–C(7), 1.809(5); P(2)–C(8), 1.862(5); C(7)–C(8), 1.345(6); P(3)–C(9), 1.826(5); C(8)–C(9), 1.475(6); C(9)–C(10), 1.329(6); P(4)–C(11), 1.778(5); P(5)–C(12), 1.770(5); C(11)–C(12), 1.214(6). Selected bond angles (°): P(1)–W(1)–C(10), 74.0(1); W(1)–C(10)–C(9), 158.6(4); C(10)–C(9)–C(8), 115.3(4); C(10)–C(9)–P(3), 124.3(3); C(8)–C(9)–P(3), 119.8(3); C(9)–C(8)–C(7), 124.8(4); C(9)–C(8)–P(2), 115.8(3); C(7)–C(8)–P(2), 119.5(3); C(8)–C(7)–P(1), 123.2(4); C(7)–P(1)–W(1), 113.3(2); C(11)–C(12)–P(5), 173.8(4); C(12)–C(11)–P(4), 173.8(4).

were performed at the National Chen-Kung University, Tainan, Taiwan.

3.2. Reaction of $W(CO)_3(Me_3tach)$ with $Ph_2PC\equiv CPh_2$

$W(CO)_3(Me_3tach)$ (500 mg, 1.26 mmol) and $Ph_2PC\equiv CPh_2$ (800 mg, 2.01 mmol) were placed in an oven-dried 100 ml Schlenk flask, equipped with a magnetic stir bar and a rubber serum stopper. Freshly distilled dichloromethane (100 ml) was introduced into the flask via a syringe through the serum stopper. The mixture was stirred at room temperature for 2 days, resulting in an orange solution. The mixture was filtered into a Schlenk tube, and the filtrate was carefully layered with *n*-hexane (100 ml) and left to settle at ambient temperature. Bright yellow crystals of $W_2(CO)_6(\mu-Ph_2PC\equiv CPh_2)_3$ (**1**; 805 mg, 0.47 mmol, 74% based on W atoms) were afforded. The mother liquid was evaporated to dryness on a rotary evaporator, and the residue was subjected to TLC with *n*-hexane/dichloromethane (1:1, v v⁻¹) as eluant. Crystallization of the material forming the orange band from benzene/*n*-hexane produced purple-red, plate crystals of $W_2(CO)_6(\mu-Ph_2PC\equiv CPh_2)[\mu-C_4H(PPh_2)_3]$ (**2**; 35 mg, 3.4%).

1: IR (CH₂Cl₂, ν_{CO}): 1950vs, 1866vs cm⁻¹. ¹H-NMR (CD₂Cl₂, 25 °C): 7.40 (t, 12H), 7.25 (t, 24H), 7.04 (t, *J* = 8 Hz, 24H) ppm. ³¹P{¹H}-NMR (CD₂Cl₂, 25 °C): –1.75 (s, with ¹⁸³W satellites, *J*_{W-P} = 235 Hz) ppm.

Anal. found: C, 58.22; H, 3.40. C₈₄H₆₀O₆P₆W₂ Anal. Calc.: C, 58.69; H, 3.52%.

2: IR (CH₂Cl₂, ν_{CO}): 1966s, 1944vs, 1906w, 1850 cm⁻¹. ¹H-NMR (CD₂Cl₂, 25 °C): 7.85–6.43 (m, 50H, Ph), 5.84 (m, 1H, CH=) ppm. ³¹P{¹H}-NMR (CD₂Cl₂, 25 °C): 57.8 (m, 1P, with ¹⁸³W satellites, *J*_{W-P} = 226 Hz), 49.4 (m, 1P, with ¹⁸³W satellites, *J*_{W-P} = 228 Hz), 31.8 (m, 1P, with ¹⁸³W satellites, *J*_{W-P} = 204 Hz), –5.1 (m, 1P, with ¹⁸³W satellites, *J*_{W-P} = 215 Hz), –6.3 (m, 1P, with ¹⁸³W satellites, *J*_{W-P} = 242 Hz) ppm. MS (FAB) *m/z* 1534 [M⁺, ¹⁸⁴W].

3.3. Transformation of **1** into **2**

A solution of **1** (20 mg) in dichloromethane (20 ml) was added into a Schlenk flask under dinitrogen, and the mixture was stirred at room temperature for 24 h, showing no reactions as evidenced by IR and TLC. *Me*₃tach (10 μl) was then introduced, and the mixture was stirred for another 24 h, resulting an orange solution. Compound **2** was obtained after separation, accompanied with several uncharacterized products.

3.4. Structure determination for **1** and **2**

Crystals of **1** and **2** suitable for X-ray diffraction studies were grown from dichloromethane/*n*-hexane and benzene/*n*-hexane at ambient temperature, respectively. They were each mounted in a thin-walled glass capillary and aligned on the Nonius Kappa CCD diffractometer

Table 1
Crystal data and refinement details for **1** and **2**

	1	2
Formula	C ₈₄ H ₆₀ O ₆ P ₆ W ₂	C ₇₂ H ₅₁ O ₆ P ₅ W ₂
<i>T</i> (K)	150(1)	150
Crystal system	triclinic	orthorhombic
Crystal solvent		C ₆ H ₆
Space group	<i>P</i> $\bar{1}$	<i>P</i> bcn
Unit cell dimensions		
<i>a</i> (Å)	13.5488(3)	38.3452(2)
<i>b</i> (Å)	13.6484(3)	17.3030(1)
<i>c</i> (Å)	23.1937(7)	20.2987(1)
α (°)	73.814(1)	90
β (°)	88.412(1)	90
γ (°)	60.599(1)	90
<i>V</i> (Å ³)	3557.8(2)	13467.9(1)
<i>Z</i>	2	8
<i>D</i> _{calc} (g cm ⁻³)	1.604	1.592
<i>F</i> (0 0 0)	1700	6376
Radiation λ (Å)	0.71073	0.71073
μ (mm ⁻¹)	3.421	3.586
θ range (°)	1.74–25.00	1.06–27.50
<i>R</i> ₁	0.0712	0.0391
w <i>R</i> ₂	0.1716	0.0826
GOF on <i>F</i> ²	1.269	1.102

with graphite-monochromated Mo–K α radiation ($\lambda = 0.71073$ Å). The data were collected at 150 K. All data were corrected for the effects of absorption. The structures were solved by the direct method and refined by full-matrix least-squares on *F*². The program used was the SHELXTL package [22]. All non-hydrogen atoms were refined with anisotropic displacement parameters. Hydrogen atoms were included but not refined. A summary of relevant crystallographic data is provided in Table 1.

4. Supplementary material

Crystallographic data for the structural analysis of **1** and **2** have been deposited with the Cambridge Crystallographic Data Centre, CCDC No. 199849 and 199850, respectively. Copy of this information may be obtained free of charge from: The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (Fax: +44-1223-

336033 or email: deposit@ccdc.cam.ac.uk or www: http://www.ccdc.cam.ac.uk).

Acknowledgements

We are grateful to the National Science Council of Taiwan for support of this work.

References

- [1] N. Takeda, K. Umemoto, K. Yamaguchi, M. Fujita, *Nature* 398 (1999) 794.
- [2] B. Olenyuk, J.A. Whiteford, A. Fechtenkotter, P.J. Stang, *Nature* 398 (1999) 769.
- [3] M. Fujita, *Chem. Soc. Rev.* (1998) 417.
- [4] S. Leininger, B. Olenyuk, P.J. Stang, *Chem. Rev.* 100 (2000) 853.
- [5] M. Fujita, K. Umemoto, M. Yoshizawa, N. Fujita, T. Kusukawa, K. Biradha, *Chem. Commun.* (2001) 509.
- [6] B.J. Holliday, C.A. Mirkin, *Angew. Chem. Int. Ed.* 40 (2001) 2022.
- [7] S.R. Seidei, P.J. Stang, *Acc. Chem. Res.* 35 (2002) 972.
- [8] R.H. Crabtree, *The Organometallic Chemistry of the Transition Metals*, 3rd ed., Wiley, New York, 2001.
- [9] B.F.G. Johnson, J. Lewis, A.D. Massey, P.R. Raithby, W.T. Wong, *J. Organomet. Chem.* 397 (1990) C28.
- [10] E. Louattani, J. Suades, *J. Organomet. Chem.* 604 (2000) 234.
- [11] L.R. Falvello, J. Forníés, J. Gómez, E. Lalinde, A. Martín, F. Martínez, M.T. Moreno, *J. Chem. Soc. Dalton Trans.* (2001) 2132.
- [12] G. Hogarth, T. Norman, *Polyhedron* 15 (1996) 2859.
- [13] M.V. Baker, M.R. North, *J. Organomet. Chem.* 565 (1998) 225.
- [14] S.C.N. Hsu, W.-Y. Yeh, *J. Chem. Soc. Dalton Trans.* (1998) 125.
- [15] G.L. Miessler, D.A. Tarr, *Inorganic Chemistry*, 2nd ed., Prentice-Hall, Englewood Cliffs, NJ, 2000.
- [16] A.J. Carty, A. Efraty, T.W. Ng, *Can. J. Chem.* 47 (1969) 1429.
- [17] S.L. James, E. Lozano, M. Nieuwenhuyzen, *Chem. Commun.* (2000) 617.
- [18] M. Semmelmann, D. Fenske, J.F. Corrigan, *J. Chem. Soc. Dalton Trans.* (1998) 2541.
- [19] W.-Y. Yeh, C.-L. Ho, M.Y. Chiang, I.-T. Chen, *Organometallics* 16 (1997) 2698.
- [20] W.-Y. Yeh, S.-M. Peng, G.-H. Lee, *J. Chem. Soc. Chem. Commun.* (1993) 1056.
- [21] F.-E. Hong, H. Chang, Y.-C. Chang, B.-T. Ko, *Inorg. Chem. Commun.* 4 (2001) 723.
- [22] G.M. Sheldrick, SHELXTL, University of Göttingen, Germany, 1985.