

變異係數模式之多項式平滑迴歸法(2/2)

執行期間：90 年 8 月 1 日至 91 年 7 月 31 日

計畫參與人員：

☐赴國外出差或研習心得報告一份

☐赴大陸地區出差或研習心得報告一份

☒出席國際學術會議心得報告及發表之論文各一份

☐國際合作研究計畫國外研究報告書一份

中 華 民 國 91 年 10 月 28 日

行政院國家科學委員會專題研究計畫成果報告

變異係數模式之多項式平滑迴歸法(2/2)

(Simultaneous Smoothing Splines for Varying Coefficient Models)

計畫編號：NSC 90-2118-M-002-013

執行期限：90 年 8 月 1 日至 91 年 7 月 31 日

主持人：江金倉 執行機構及單位名稱：台灣大學數學系

共同主持人： 執行機構及單位名稱

計畫參與人員： 執行機構及單位名稱

一、中文摘要

提出多項式平滑回歸估計方法，當中之反應變數為時間函數，而解釋變數為一常值，在此估計方法中，吾等考慮將其解釋變數之變異消除，此目的乃在於避免估計式受隨機平滑參數之影響及理論推導之複雜性。對此研究主題，吾已推導出估計式之大樣本理論並借助實例及模擬發現好的有限樣本之性質。

關鍵詞：

Abstract

Keywords:

二、緣由與目的

三、結果與討論

四、計畫成果自評

五、參考文獻

(以上討論於所附之研究成果)

Simultaneous Smoothing Splines for Varying Coefficient Models

Chin-Tsang Chiang
Department of Mathematics
National Taiwan University
Taipei 106, Taiwan, R.O.C.

Abstract

In this paper, simultaneous smoothing spline estimation methods are proposed to estimate the coefficient curves on the varying coefficient model with repeatedly measured (longitudinal) dependent variable and time invariant covariates. The estimators are obtained from the penalized least squares with adjustment for the variations of covariates in the penalized terms. We do this mainly to avoid the penalized terms being influenced by the scales of the covariates and the random smoothing parameters appearing in the estimators, which complicates the derivation of the properties of the estimators. We also develop the asymptotic properties of the simultaneous smoothing splines. When the smoothing parameters within each estimator are set to be equal, the simultaneous smoothing spline estimators are shown to have smaller variances than the componentwise smoothing spline estimators. Through a Monte Carlo simulation and two empirical examples, the simultaneous smoothing splines are also found to be more accurate in the variances.

1 Introduction

In biomedical and epidemiology studies, longitudinal data with time invariant covariates and repeatedly measured (longitudinal) dependent variable over time are frequently encountered. Generally speaking, this type of data is collected from n randomly selected

Key words and phrases: componentwise smoothing splines, longitudinal data, mean squared error, penalized least squares, simultaneous smoothing splines, smoothing parameters, varying coefficient model.

subjects. For the i th subject, let m_i , t_{ij} , Y_{ij} , and $\mathbf{X}_i^T = (X_{i0}, \dots, X_{ik})$ with $X_{i0} = 1$, respectively denote the number of the repeated measurements, the time of the j th measurement, the observed outcome at time t_{ij} , and the observed time invariant covariate vector. Here, the total number of observations is denoted by $N = \sum_{i=1}^n m_i$.

To model the relationship between the dependent variable $Y(t)$ and the time dependent or time invariant covariates $\mathbf{X}^T(t) = (X_0(t), \dots, X_k(t))$ with $X_0(t) = 1$, Hoover, Rice, Wu and Yang (1998) considered a more flexible varying coefficient model of Hastie and Tibshirani (1993)

$$Y(t) = \mathbf{X}^T(t)\boldsymbol{\beta}(t) + \varepsilon(t), \quad (1)$$

where $\boldsymbol{\beta}(t) = (\beta_0(t), \dots, \beta_k(t))^T$ are smooth functions of t , and $\varepsilon(t)$ is a mean zero stochastic process and is independent of $\mathbf{X}(t)$. They also proposed a class of smoothing methods to estimate the coefficient curves. Based on model (1), Hoover, *et al.* (1998), and Wu, Chiang and Hoover (1998) have developed inferences for the kernel estimators. Under some specific designs, Fan and Zhang (2000) provided a simply implemented two-step smoothing alternative. When the covariates are time dependent, Wu, Yu and Chiang (2000) also proposed a two-step smoothing method to avoid large biases appearing on the estimates.

In this paper, we focus on the covariates which are invariant with respect to the time points. Under this data setting, model (1) will be reduced to

$$Y(t) = \mathbf{X}^T\boldsymbol{\beta}(t) + \varepsilon(t), \quad (2)$$

where $\mathbf{X}^T = (X_0, \dots, X_k)$ with $X_0 = 1$. Substituting $Y(t)$, \mathbf{X} and t with observations Y_{ij} , \mathbf{X}_i and t_{ij} , model (2) becomes

$$Y_{ij} = \mathbf{X}_i^T\boldsymbol{\beta}(t_{ij}) + \varepsilon(t_{ij}), \quad i = 1, \dots, n; j = 1, \dots, m_i. \quad (3)$$

Based on (3), Wu and Chiang (2000), and Chiang, Rice and Wu (2001) modified the methods of Hoover, *et al.* (1998) into componentwise estimation methods to significantly simplify the computations. They also derived the asymptotic properties of the estimators through the explicit expressions of their asymptotic risk representations. Meanwhile, through a Monte Carlo simulation, the sample variances of their estimators are found to be smaller than those of Fan and Zhang (2000). However, in succeeding sections, the componentwise smoothing spline estimators are shown not as accurate as it is expected in the variances under both the finite sample and the infinite sample. This is mainly caused by the unexpected non-negative terms, which are functions of the moments of the covariates \mathbf{X} and the parameter curves $\beta(t)$, in the variances of the estimators.

Instead of using the componentwise estimation methods, we propose the simultaneous smoothing spline estimation methods based on the penalized least squares with adjustment for the variations of the covariates in the penalized terms, which avoid the penalized terms being influenced by the scales of the covariates. There are two features of our simultaneous smoothing spline estimation methods: First, our estimators are unlike the smoothing spline estimators of Hoover, *et al.* (1998), which are smoothen by the random smoothing parameters and are more complicated in terms of deriving the properties of the estimators. Second, when the smoothers within each estimator are set to be equal, it is shown that the mean squared errors of our simultaneous smoothing spline estimators are smaller than the corresponding componentwise smoothing splines.

The contents of this paper are organized as follows. In Section 2, we propose the simultaneous smoothing spline estimation methods, and summarize the componentwise smoothing spline estimation methods. The asymptotic mean squared errors of the pro-

posed estimators with or without equal smoothing parameters for each estimator are developed in Section 3. For the sake of comparison, the asymptotic mean squared errors of the componentwise smoothing splines are also stated in this section. In Section 4, a Monte Carlo simulation is implemented to examine the finite sample properties of the simultaneous smoothing spline estimators. Applications of our estimation methods are also demonstrated in Section 5 through two empirical examples — a CD4 depletion study and an opioid detoxification study. Finally, the proofs of the main results are shown in the Appendix.

2 Estimation

Assume that the support of the design time points $\{t_{ij}\}$ is contained in a compact set $[a, b]$, and $\beta_l(t), l = 0, \dots, k$, are twice differentiable. Also, let $\mathcal{H}_{[a,b]}$ be the set of compact supported functions such that

$$\mathcal{H}_{[a,b]} = \left\{ g(t) : g(t) \text{ and } g^{(1)}(t) \text{ are absolutely continuous, and } g^{(2)}(t) \in L^2[a, b] \right\}.$$

The simultaneous smoothing splines $\hat{\beta}_{(s)}(t; \boldsymbol{\lambda}) = (\hat{\beta}_{0(s)}(t; \boldsymbol{\lambda}), \dots, \hat{\beta}_{k(s)}(t; \boldsymbol{\lambda}))^T$ of $\boldsymbol{\beta}(t)$ proposed here are obtained by minimizing the penalized least squares with adjustment for the variations of the covariates in the penalized terms

$$J_{1s}(\boldsymbol{\beta}; \boldsymbol{\lambda}) = \sum_{i=1}^n \sum_{j=1}^{m_i} w_i \left[Y_{ij} - \left(\sum_{l=0}^k X_{il} \beta_l(t_{ij}) \right) \right]^2 + \sum_{l=0}^k \lambda_l s_l^2 \int_a^b \left(\beta_l^{(2)}(t) \right)^2 dt, \quad (4)$$

where $\boldsymbol{\lambda} = (\lambda_0, \dots, \lambda_k)$ are non-negative smoothing parameters, $\mathbf{w} = (w_1, \dots, w_n)$ are non-negative constant weights with $\sum_{i=1}^n m_i w_i = 1$, $s_l = \sum_{i=1}^n (w_i m_i) X_{il}^2$, and $\beta_l^{(p)}(t)$ denotes the p th derivative of $\beta_l(t)$ with respect to t . In practice, w_i 's are usually assigned to be $1/N$ and $1/(nm_i)$ which provide equal weight to each single observation and each

single subject, respectively. However, as mentioned in Remark 8 of Chiang, *et al.* (2001), there may not have the explicit risk representations for the estimators with $w_i = 1/(nm_i)$ or more general weights. When the numbers of the repeated measurements are bounded, it was suggested by Lin and Carroll (2000) that $w_i = 1/N$ leads to asymptotically optimal kernel smoothers for the generalized estimating equations. For the sake of comparison with the existing estimators in the literature, the weights are assigned to be $1/N$ in the succeeding discussions. Setting the Gateaux derivatives of $J_{1s}(\beta; \lambda)$ to zero, $\hat{\beta}_{(s)}(t; \lambda)$ uniquely minimize $J_{1s}(\beta; \lambda)$ if they satisfy the following normal equations

$$\sum_{i=1}^n \sum_{j=1}^{m_i} \left[\frac{X_{il}}{Ns_l^2} \left(Y_{ij} - \sum_{l_1=0}^k \hat{\beta}_{l_1(s)}(t_{ij}; \lambda) X_{il_1} \right) g_l(t_{ij}) \right] = \lambda_l \int_a^b \hat{\beta}_{l(s)}^{(2)}(t; \lambda) g_l^{(2)}(t) dt, \quad (5)$$

for $l = 0, \dots, k$, and all g_l 's in a dense subset of $\mathcal{H}_{[a,b]}$. A similar argument as in Wahba (1975) shows that there is a symmetric function $S_{\lambda_l, X_l}(t, s)$ in $\mathcal{H}_{[a,b]}$ so that $\hat{\beta}_{l(s)}(t; \lambda)$ is given by

$$\hat{\beta}_{l(s)}(t; \lambda) = \sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{Ns_l^2} (Y_{ij} - \sum_{l_1 \neq l} \hat{\beta}_{l_1(s)}(t_{ij}; \lambda) X_{il_1}) S_{\lambda_l, X_l}(t, t_{ij}). \quad (6)$$

By substituting (6) into (5) and rearranging the terms, we have the characterization of the spline function $S_{\lambda_l, X_l}(t, t_{ij})$,

$$\int_a^b S_{\lambda_l, X_l}(t, t_{ij}) g_l(t) dF_{N, X_l}(t) + \lambda_l \int_a^b S_{\lambda_l, X_l}^{(2)}(t, t_{ij}) g_l^{(2)}(t) dt = g_l(t_{ij}), \quad (7)$$

where $F_{N, X_l}(t) = \sum_{i=1}^n \frac{m_i X_{il}^2}{Ns_l^2} 1_{[t_{ij} \leq t]}$.

In (6), we can see that the estimator $\hat{\beta}_{l(s)}(t; \lambda)$ is influenced not only by the smoothing parameter λ_l but also by the other smoothers. To make $\hat{\beta}_{l(s)}(t; \lambda)$ being smoothen only by λ_l , the smoothing parameters λ are set equal to $\lambda_l \mathbf{1}$. Then, the by-product estimators, denoted by $\hat{\beta}_{l(s)}(t; \lambda_l)$, can be expressed as

$$\hat{\beta}_{l(s)}(t; \lambda_l) = \sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{Ns_l^2} (Y_{ij} - \sum_{l_1 \neq l} \hat{\beta}_{l_1(s)}(t_{ij}; \lambda_l) X_{il_1}) S_{\lambda_l, X_l}(t, t_{ij}). \quad (8)$$

For the smoothing spline estimation methods of Hoover, *et al.* (1998), their estimators are obtained by minimizing

$$J_{2s}(\boldsymbol{\beta}; \boldsymbol{\lambda}) = \frac{1}{N} \sum_{i=1}^n \sum_{j=1}^{m_i} \left(Y_{ij} - \sum_{l=0}^k X_{il} \beta_l(t_{ij}) \right)^2 + \sum_{l=0}^k \lambda_l \int_a^b \left(\beta_l^{(2)}(t) \right)^2 dt. \quad (9)$$

The same reasoning as in the derivation of $\hat{\beta}_{l(s)}(t; \boldsymbol{\lambda})$ shows that the corresponding minimizers, say, $\hat{\beta}_{l(s)}^*(t; \boldsymbol{\lambda}^*)$ of $J_{2s}(\boldsymbol{\beta}; \boldsymbol{\lambda})$ are given by

$$\hat{\beta}_{l(s)}^*(t; \boldsymbol{\lambda}^*) = \sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{N s_l^2} \left(Y_{ij} - \sum_{l_1 \neq l} \hat{\beta}_{l_1(s)}^*(t_{ij}; \boldsymbol{\lambda}^*) X_{il_1} \right) S_{\lambda_l^*, X_l}(t, t_{ij}), \quad (10)$$

where $\boldsymbol{\lambda}^* = (\lambda_0^*, \dots, \lambda_k^*)$ with $\lambda_l^* = \lambda_l / s_l^2$. As we can see, our procedures work like the usual smoothing methods of Hoover, *et al.* (1998). However, the random smoothing parameters $\boldsymbol{\lambda}^*$ will cause complexity in deriving the properties of $\hat{\beta}_{l(s)}^*(t; \boldsymbol{\lambda}^*)$. By contrast, the simultaneous smoothing spline estimators $\hat{\beta}_{l(s)}(t; \boldsymbol{\lambda})$ can avoid this problem, and are more common in many practical applications. In the following sections, we will focus only on this discussion of $\hat{\beta}_{l(s)}(t; \boldsymbol{\lambda})$.

To avoid intensive computation in the estimation of the coefficient curves $\boldsymbol{\beta}(t)$, Chiang, *et al.* (2001) proposed the componentwise smoothing spline estimation methods for the varying coefficient model (2). Even though these methods are fast in computation, the variances of the estimators, as shown later, are not accurate enough. Especially, computing considerations are no longer a major consideration in modern computer equipment. For the sake of comparison in succeeding sections, we summarize here their estimation methods.

Assume that the inverse of $E[\mathbf{X}\mathbf{X}^T]$, denoted by $E_{\mathbf{X}\mathbf{X}^T}^{-1}$, exists. By multiplying the both sides of (2) by \mathbf{X} and taking expectations, $\boldsymbol{\beta}(t)$ can be expressed as

$$\boldsymbol{\beta}(t) = E[\mathbf{Z}(t)], \quad (11)$$

where $\mathbf{Z}(t) = (Z_0(t), \dots, Z_k(t))^T$ with $Z_l(t) = \sum_{r=0}^k e_{lr} X_r Y(t)$ and e_{lr} the (l, r) th element of $E_{\mathbf{X}\mathbf{X}^T}^{-1}$. Since $E[\mathbf{X}\mathbf{X}^T]$ is unknown and is invariant with respect to time t , it is naturally estimated by the sample mean

$$E[\widehat{\mathbf{X}\mathbf{X}^T}] = n^{-1} \sum_{i=1}^n (\mathbf{X}_i \mathbf{X}_i^T). \quad (12)$$

Assume further that the inverse of $E[\widehat{\mathbf{X}\mathbf{X}^T}]$, denoted by $\widehat{E}_{\mathbf{X}\mathbf{X}}^{-1}$, exists. Substituting $E_{\mathbf{X}\mathbf{X}^T}^{-1}$ with $\widehat{E}_{\mathbf{X}\mathbf{X}}^{-1}$, the componentwise estimator, say, $\widehat{\beta}_{l(c)}(t; \lambda_l)$ of $\beta_l(t)$ is obtained by minimizing the following penalized least squares

$$J_c(\beta_l; \lambda_l) = \frac{1}{N} \sum_{i=1}^n \sum_{j=1}^{m_i} (\widehat{Z}_{ijl} - \beta_l(t_{ij}))^2 + \lambda_l \int_a^b (\beta_l^{(2)}(t))^2 dt, \quad (13)$$

where $\widehat{Z}_{ijl} = \sum_{r=0}^k \widehat{e}_{lr} X_{il} Y_{ij}$ is the estimated observed value of Z_{ijl} with \widehat{e}_{lr} being the (l, r) th element of $\widehat{E}_{\mathbf{X}\mathbf{X}}^{-1}$. The minimizer $\widehat{\beta}_{l(c)}(t; \lambda_l)$ of $J_c(\beta_l; \lambda_l)$ can be expressed as

$$\widehat{\beta}_{l(c)}(t; \lambda_l) = \frac{1}{N} \sum_{i=1}^n \sum_{j=1}^{m_i} \widehat{Z}_{ijl} S_{\lambda_l}(t, t_{ij}), \quad (14)$$

where $S_{\lambda_l}(t, s) = S_{\lambda_l, X_0}(t, s)$ is in $\mathcal{H}_{[a, b]}$.

As derived in that paper, the asymptotic variance of the estimator $\widehat{\beta}_{l(c)}(t; \lambda_l)$ contains the unexpected non-negative terms, which are functions of the moments of the covariates and the parameter curves. In Section 3, we will show that both $\widehat{\beta}_{l(s)}(t; \lambda_l)$ and $\widehat{\beta}_{l(c)}(t; \lambda_l)$ have the same asymptotic bias, but the asymptotic variance of $\widehat{\beta}_{l(s)}(t; \lambda_l)$ is smaller. Through a Monte Carlo simulation in Section 4 and two empirical examples in Section 5, it appears that $\widehat{\beta}_{l(s)}(t; \lambda_l)$ is more accurate in variance than $\widehat{\beta}_{l(c)}(t; \lambda_l)$. Except the evidence from the finite and the infinite sample properties of the estimators, the unexpected terms in the variances of $\widehat{\beta}_{l(c)}(t; \lambda_l)$ can also be explained by the following reasoning:

From (2) and the definition of $\mathbf{Z}(t)$, $\mathbf{Z}(t)$ can be reexpressed as

$$\begin{aligned}
\mathbf{Z}(t) &= E_{\mathbf{X}\mathbf{X}^T}^{-1} \mathbf{X}Y(t) \\
&= E_{\mathbf{X}\mathbf{X}^T}^{-1} (\mathbf{X}\mathbf{X}^T) \boldsymbol{\beta}(t) + E_{\mathbf{X}\mathbf{X}^T}^{-1} \mathbf{X}\varepsilon(t) \\
&= \boldsymbol{\beta}(t) + \varepsilon^*(t),
\end{aligned} \tag{15}$$

where

$$\varepsilon^*(t) = E_{\mathbf{X}\mathbf{X}^T}^{-1} (\mathbf{X}\mathbf{X}^T - E[\mathbf{X}\mathbf{X}^T]) \boldsymbol{\beta}(t) + E_{\mathbf{X}\mathbf{X}^T}^{-1} \mathbf{X}\varepsilon(t).$$

One can observe that the new error process $\varepsilon^*(t)$ consists of two components: the variabilities of the covariates and the original stochastic error process $\varepsilon(t)$. Thus, the variances of $\hat{\beta}_{l(c)}(t; \lambda_l)$'s are enlarged by the extra non-negative terms.

3 Asymptotic Properties

The asymptotic mean squared errors of the simultaneous smoothing spline estimators $\hat{\beta}_{l(s)}(t; \boldsymbol{\lambda})$, $l = 0, \dots, k$, will be derived in this section. Without loss of generality, we focus on the interval $[0, 1]$. Extension to the general interval $[a, b]$ can be carried out by the affine transformation $u = (t - a)/(b - a)$ for $t \in [a, b]$. For the succeeding discussions, we make the following assumptions:

(A1) The time design points $\{t_{ij}\}$ are nonrandom and satisfy

$$D_N = \sup_{t \in [0, 1]} |F_N(t) - F(t)| \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

for some distribution function F with strictly positive density f on $[0, 1]$, where $F_N(t) = N^{-1} \sum_{i=1}^n \sum_{j=1}^{m_i} 1_{[t_{ij} \leq t]}$, and f is three times differentiable and uniformly continuous on $[0, 1]$ with $f^{(\nu)}(0) = f^{(\nu)}(1) = 0$ for $\nu = 1, 2$.

(A2) The coefficient curve $\beta_l(t)$ is four times differentiable and satisfies the boundary conditions $\beta_l^{(\nu)}(0) = \beta_l^{(\nu)}(1) = 0$ for $\nu = 2, 3$. The fourth derivative $\beta_l^{(4)}(t)$ is Lipschitz continuous in the sense that $|\beta_l^{(4)}(s_1) - \beta_l^{(4)}(s_2)| \leq c_{1l}|s_1 - s_2|^{c_{2l}}$ for all $s_1, s_2 \in [0, 1]$ and some positive constants c_{1l} and c_{2l} .

(A3) The fourth moment $E[X_l^4]$ exists.

(A4) Define $D_{N,l} = \sup_{t \in [0,1]} |F_{N,X_l}(t) - F(t)|$. $\lambda_l = O(\lambda_0) \rightarrow 0$, $N\lambda_l^{1/4} \rightarrow \infty$ and $\lambda_l^{-5/4} D_{N,l} \xrightarrow{a.s.} 0$ as $n \rightarrow \infty$.

(A5) Define $\sigma^2(t) = E[\varepsilon^2(t)]$ and $\rho(t) = \lim_{t' \rightarrow t} E[\varepsilon(t)\varepsilon(t')]$. Both $\sigma^2(t)$ and $\rho(t)$ are continuous at t .

Under the validity of assumption A3, it is straightforward to show by the law of large numbers that $\sup_{t \in [0,1]} |F_{N,X_l}(t) - F_N(t)|$ converges to zero almost surely. With assumption A1 and the property $D_{N,l} \leq D_N + \sup_{t \in [0,1]} |F_{N,X_l}(t) - F_N(t)|$, we can show that $D_{N,l}$ in assumption A4 converges to zero almost surely. In general, $\sigma^2(t)$ need not be equal to $\rho(t)$. As discussed in Zeger and Diggle (1994), the strict inequality appears when $\varepsilon(t)$ consists of a stationary process of t and an independent measurement error. Since the spline function $S_{\lambda_l, X_l}(t, s)$ in (7) does not have an explicit expression, it may be approximated by the Green function $G_{\lambda_l}(t, s)$ of the 4th order differential equation

$$\lambda_l g_l^{(4)}(t) + f(t)g_l(t) = f(t)\beta_l(t), \quad t \in [0, 1], \quad (16)$$

with $g_l^{(\nu)}(0) = g_l^{(\nu)}(1)$ for $\nu = 2, 3$. By substituting the smoothing spline function $S_{\lambda_l, X_l}(t, s)$ with the Green function $G_{\lambda_l}(t, s)$ and using the exponential bound of $|S_{\lambda_l, X_l}(t, s) - G_{\lambda_l}(t, s)|$, the asymptotic properties of $\hat{\beta}_{l(s)}(t; \lambda)$ can be conveniently derived. It was also shown by Abramovich and Grinshtein (1999) and Chiang, *et al.* (2001) that the Green function $G_{\lambda_l}(t, s)$ can be approximated by

$$\begin{aligned} H_{\lambda_l}(t, s) &= \frac{(\lambda_l/\gamma^4)^{-1/4}}{2} \Gamma^{(1)}(s) (f(s))^{-1} \sin \left(\frac{\pi}{4} + \frac{(\lambda_l/\gamma^4)^{-1/4}}{\sqrt{2}} |\Gamma(t) - \Gamma(s)| \right) \\ &\quad \times \exp \left(-\frac{(\lambda_l/\gamma^4)^{-1/4}}{\sqrt{2}} |\Gamma(t) - \Gamma(s)| \right), \end{aligned} \quad (17)$$

where $\gamma = \int_0^1 (f(s))^{1/4} ds$ and $\Gamma(t) = \gamma^{-1} \int_0^t (f(s))^{1/4} ds$. Some important properties of the functions $S_{\lambda_l}(t, s)$, $G_{\lambda_l}(t, s)$ and $H_{\lambda_l}(t, s)$, which will be used in the main results, are stated in the following Lemma.

Lemma 1. Under assumptions A1 and A4, there are positive constants α_1 , α_2 , κ_1 and κ_2 so that

$$|G_{\lambda_l}(t, s) - H_{\lambda_l}(t, s)| \leq \kappa_1 \exp \left(-\alpha_1 \lambda_l^{-1/4} |t - s| \right), \quad (18)$$

$$\left| \frac{\partial^\nu}{\partial t^\nu} G_{\lambda_l}(t, s) \right| \leq \kappa_1 \lambda_l^{-(\nu+1)/4} \exp \left(-\alpha_2 \lambda_l^{-1/4} |t - s| \right), \quad 0 \leq \nu \leq 3, \quad (19)$$

$$|S_{\lambda_l, X_l}(t, s) - G_{\lambda_l}(t, s)| \leq \kappa_2 \lambda_l^{-1/2} D_{N,l} \exp \left(-\alpha_1 \lambda_l^{-1/4} |t - s| \right) \text{ a.s.}, \quad (20)$$

and

$$\left| \frac{\partial^\nu}{\partial t^\nu} S_{\lambda_l, X_l}(t, s) \right| \leq \kappa_2 \lambda_l^{-(\nu+1)/4} D_{N,l} \exp \left(-\alpha_2 \lambda_l^{-1/4} |t - s| \right) \text{ a.s.} \quad (21)$$

hold uniformly for $t, s \in [0, 1]$ and $0 \leq \nu \leq 3$.

Proof. The proof can be derived along the same lines as the proof in Lemma 3.1 of Chiang, *et al.* (2001). \square

Let $\mathbf{B}(\hat{\beta}_{(s)}(t; \lambda))$ and $\mathbf{V}(\hat{\beta}_{(s)}(t; \lambda))$ be the bias and the variance of $\hat{\beta}_{(s)}(t; \lambda)$. By the decomposition principle of the mean squared error, we can separately evaluate the bias and the variance of $\hat{\beta}_{(s)}(t; \lambda)$.

Theorem 1. Suppose that assumptions (A1)-(A5) are satisfied and $t \in (0, 1)$. Then, for sufficiently large n , the bias and the variance of $\hat{\beta}_{(s)}(t; \lambda)$ are given by

$$\mathbf{B}(\hat{\beta}_{(s)}(t; \lambda)) = -(f(t))^{-1} \left[(E_{\mathbf{X}\mathbf{X}^T}^{-1} \Lambda E[\mathbf{X}\mathbf{X}^T]) \beta^{(4)}(t) \right]^T (1 + o(1)), \quad (22)$$

and

$$\begin{aligned} \mathbf{V}(\hat{\beta}_{(s)}(t; \lambda)) &= \frac{1}{2\sqrt{2}N} (f(t))^{-3/4} \sigma^2(t) (E_{\mathbf{X}\mathbf{X}^T}^{-1} \Lambda_{\mathbf{X}\mathbf{X}^T} E_{\mathbf{X}\mathbf{X}^T}^{-1}) (1 + o(1)) \\ &+ \left(\sum_{i=1}^n \left(\frac{m_i}{N} \right)^2 \rho(t, t) \right) E_{\mathbf{X}\mathbf{X}^T}^{-1} (1 + o(1)), \end{aligned} \quad (23)$$

where $\Lambda = \text{diag}(\lambda_0, \dots, \lambda_k)$, $\beta^{(4)}(t) = (\beta_0^{(4)}(t), \dots, \beta_k^{(4)}(t))^T$, and $\Lambda_{\mathbf{X}\mathbf{X}^T} = [(\lambda_{l_1 l_2} E[X_{l_1} X_{l_2}])]$ with $\lambda_{l_1 l_2} = (\lambda_{l_1}^{1/4} + \lambda_{l_2}^{1/4})^{-1}$.

Proof. See Appendix. \square

Following the arguments in Hoover, *et al.* (1998), the variance of $\hat{\beta}_{l(s)}(t; \lambda)$ asymptotically converges to zero if and only if $\max_{1 \leq i \leq n} (m_i/N) \rightarrow 0$. When the smoothing parameters λ in $\hat{\beta}_{l(s)}(t; \lambda)$ are set to be equal to $\lambda_l \mathbf{1}$, the asymptotic properties of $\hat{\beta}_{l(s)}(t; \lambda_l)$ in (8) can be derived straightforward from Theorem 1.

Lemma 2. Suppose that assumptions (A1)-(A5) are satisfied and $t \in (0, 1)$. Then, for sufficiently large n , the bias and the variance of $\hat{\beta}_{l(s)}(t; \lambda_l)$, $l = 0, \dots, k$ are given by

$$B(\hat{\beta}_{l(s)}(t; \lambda_l)) = -(f(t))^{-1} \beta_l^{(4)}(t) \lambda_l (1 + o(1)), \quad (24)$$

and

$$\begin{aligned}
V(\hat{\beta}_{l(s)}(t; \lambda_l)) &= \frac{1}{4\sqrt{2}}(f(t))^{-3/4}(N\lambda_l^{1/4})^{-1}e_{ll}\sigma^2(t)(1 + o(1)) \\
&+ \left(\sum_{i=1}^n \frac{m_i}{N}\right)^2 e_{ll}\rho(t, t)(1 + o(1)). \tag{25}
\end{aligned}$$

To compare the asymptotic properties between $\hat{\beta}_{l(s)}(t; \lambda_l)$ and $\hat{\beta}_{l(c)}(t; \lambda_l)$, we summarize the asymptotic properties of $\hat{\beta}_{l(c)}(t; \lambda_l)$ in Theorem 2.

Theorem 2. Suppose that assumptions (A1)-(A5) are satisfied and $t \in (0, 1)$. When n is sufficiently large, conditional on \mathbf{X} , the bias and the variance of $\hat{\beta}_{l(c)}(t; \lambda_l)$, $l = 0, \dots, k$, are given by

$$B(\hat{\beta}_{l(c)}(t; \lambda_l)) = -(f(t))^{-1}\beta_l^{(4)}(t)\lambda_l(1 + o_p(1)) + O_p(n^{-1/2}), \tag{26}$$

and

$$\begin{aligned}
V(\hat{\beta}_{l(c)}(t; \lambda_l)) &= \frac{1}{4\sqrt{2}}(f(t))^{-3/4}(N\lambda_l^{1/4})^{-1}(M_l(t) + e_{ll}\sigma^2(t))(1 + o_p(1)) \\
&+ \left(\sum_{i=1}^n \left(\frac{m_i}{N}\right)^2\right)(M_l(t) + e_{ll}\rho(t, t))(1 + o_p(1)) + O_p(n^{-1}), \tag{27}
\end{aligned}$$

where $M_l(t) = \sum_{l_1=1}^k \sum_{l_2=1}^k (\beta_{l_1}(t)\beta_{l_2}(t)E[X_{l_1}X_{l_2}(\sum_{l_3=0}^k e_{ll_3}X_{l_3})^2]) - (\beta_l(t))^2$.

Proof. See Chiang, *et al.* (2001).

It follows from Lemma 2 and Theorem 2 that both the dominating terms in the biases of $\hat{\beta}_{l(s)}(t; \lambda_l)$ and $\hat{\beta}_{l(c)}(t; \lambda_l)$ are same. However, the dominating term in the variance of $\hat{\beta}_{l(s)}(t; \lambda_l)$ is smaller than that of $\hat{\beta}_{l(c)}(t; \lambda_l)$ since $M_l(t)$ is nonnegative.

4 Monte Carlo Simulation

Consider the varying coefficient model (2) with coefficient curves

$$\begin{aligned}\beta_0(t) &= 3.5 + 6.5 \sin\left(\frac{t\pi}{60}\right), \\ \beta_1(t) &= -0.2 - 1.6 \cos\left(\frac{(t-30)\pi}{60}\right), \\ \beta_2(t) &= 0.25 - 0.0074 \left(\frac{30-t}{10}\right)^3\end{aligned}$$

and covariate vector $\mathbf{X} = (X_0, X_1, X_2)^T$, where X_1 and X_2 are independent Bernoulli and Gaussian random variables with joint density

$$f(x_1, x_2) = \frac{1}{8(2\pi)^{1/2}} \exp\left(-\frac{x_2^2}{32}\right) 1_{\{0,1\}}(x_1) 1_{(-\infty, \infty)}(x_2).$$

In this simulation, 400 subjects are scheduled to appear at equally spaced time points $0, 1, \dots, 30$ with a 60% probability of missing for each of the 31 “appointments”. The covariates of each subject are independently generated from the above distribution. Under the given time points $\{t_{ij}\}$, the errors $\varepsilon(t_{ij})$, which are independent of the covariates \mathbf{X}_i , are generated from the mean zero Gaussian process with covariance function

$$\text{cov}(\varepsilon(t_{i_1 j_1}), \varepsilon(t_{i_2 j_2})) = \begin{cases} 0.0625 \exp(-|t_{i_1 j_1} - t_{i_2 j_2}|), & \text{if } i_1 = i_2, \\ 0, & \text{if } i_1 \neq i_2. \end{cases}$$

Finally, the dependent variables Y_{ij} are automatically obtained by substituting \mathbf{X}_i , t_{ij} and $\varepsilon(t_{ij})$ into (3).

Based on the above design, the longitudinal data are repeatedly generated 500 times. In each set of simulated data, $\hat{\beta}_{l(s)}(t; \lambda_l)$ and $\hat{\beta}_{l(c)}(t; \lambda_l)$ are computed by (8) and (14) with appropriate smoothing parameters. As mentioned in Chiang, *et al.* (2001), the

“leave one subject out” cross-validation procedure of Rice and Silverman (1991) may sometimes select inadequate smoothing parameters. It is usually preferable to have a set of smoothing parameters which has the corresponding cross validation score close to the minimum and gives better estimators. For the purpose of comparison, the smoothing parameters $(\lambda_0, \lambda_1, \lambda_2) = (1, 1, 1)$ from their simulation are used to both estimators. Table 1 through Table 3 show the true curves, the 500 averages of the estimated curves and the standard errors of the 500 simulation estimates at nine selected time points. As shown in these tables, the variances of the componentwise smoothing spline estimators are enlarged by the values of $M_l(t)$ s, and thus are larger than those of the simultaneous smoothing splines. The results are consistent with the asymptotic properties discussed in Section 3.

5 Application

In this section, the proposed simultaneous smoothing spline estimation methods and the componentwise ones are applied to two empirical examples. These longitudinal data sets arise from a CD4 depletion study and an opioid detoxification study.

5.1 A CD4 Depletion Study

The first data set is from the Multicenter AIDS Cohort Study (MACS), which includes 283 homosexual men who were infected by HIV-1 virus. Measurements taken include CD4 percentage, the cigarette smoking status, pre-HIV infection CD4 percentage, and age at HIV infection. Individuals were repeatedly measured at scheduled semi-annual visits between 1984 and 1991. During the study period, many individuals missed some

of their scheduled visits. Thus, the numbers of repeated measurements may differ among subjects. Details of the design and the methods of this study are described in Kaslow, *et al.* (1987).

In this study, the objective is to evaluate the effects of cigarette smoking, pre-HIV infection CD4 percentage, and age at HIV infection on the mean post-HIV CD4 percentage at any given time since the infection among seroconverters. Based on model (2), the simultaneous smoothing spline estimation methods and the componentwise smoothing spline estimation methods are used to estimate the effects of the concerned covariates. Estimators $\hat{\beta}_{l(s)}(t; \lambda_l)$ and $\hat{\beta}_{l(c)}(t; \lambda_l)$ are separately computed from (8) and (14) with the smoothing parameters $(\lambda_0, \lambda_1, \lambda_2, \lambda_3) = (0.1, 0.01, 1, 0.1)$, which have the corresponding cross-validation score close to the minimum.

Figures (1a)-(1d) show the estimated curves and their 95% pointwise bootstrap confidence intervals. From these graphs, we can see that $\hat{\beta}_{l(c)}(t; \lambda_l)$ and $\hat{\beta}_{l(s)}(t; \lambda_l)$ have similar physical explanations. However, the confidence bands of $\hat{\beta}_{l(c)}(t; \lambda_l)$ are wider than $\hat{\beta}_{l(s)}(t; \lambda_l)$. As mentioned in Section 3 and Section 4, the simultaneous smoothing spline estimation methods are more reliable. From Figure (1a), the mean CD4 percentage for the non-smoking group with average pre-infection CD4 percentage and average age at HIV infection appears depleting rather quickly at the beginning of HIV infection, but the rate of depletion seems to be slowing down for the later period of the study after infection. No significant effects are detected in Figures (1b) and (1d) for cigarette smoking and age at HIV infection. However, it appears in Figure (1c) that the pre-HIV infection CD4 percentage associated positively with higher CD4 percentage after the infection.

5.2 An Opioid Detoxification Study

The second data set is from the National Institute on Drug Abuse (NIDA) opioid detoxification study, which includes 60 opioid dependent (DSM-IV) heroin users seeking detoxification treatment. In the study design, 32 patients are randomly assigned to the naltrexone-buprenorphine group and 28 to the placebo-buprenorphine group. Measurements were taken at 9 scheduled times per day by a trained nurse for a total 72 (8×9) measurements. During an 8 day inpatient clinical trial, each patient was subjected to the observer-rated opioid withdrawal scale (OOW) measurement, a scale to rate opioid withdrawal symptoms. Since some patients randomly missed some scheduled measurements or quit the treatment altogether, the number of measurements may be different for each patient. Details of this design and its medical implications can be found in Umbricht-Schneiter, *et al.* (1999).

The objective here is to detect the effects of treatment status and the centered baseline OOW score on the OOW scores over the trials. Similar to the process of analysis in Section 5.1, two estimation methods are used to detect the effects of the interesting covariates. Here, estimators $\hat{\beta}_{l(c)}(t; \lambda_l)$ and $\hat{\beta}_{l(s)}(t; \lambda_l)$ are computed with the smoothing parameters $(\lambda_0, \lambda_1, \lambda_2) = (0.001, 0.001, 0.1)$.

Figures (2a)-(2c) show the estimated curves and their 95% pointwise bootstrap confidence intervals. From these graphs, two estimation methods provide similar explanations. Also, the confidence bands for some of the componentwise smoothing spline estimators are close to the corresponding simultaneous ones. This can be explained by the small effect of $M_l(t)$ to the variance of $\hat{\beta}_{l(c)}(t; \lambda_l)$ in (27) for some l . It is shown that the placebo mean stays very close to a constant throughout the trials, while the

naltrexone treatment is generally associated with lower OOW scores roughly after the later half of the trial. The peak at the beginning of the trial for the naltrexone treatment is mainly caused by the patient's initial negative reaction to the treatment. As expected, the baseline OOW score has a positive association with the OOW scores.

APPENDIX A.

Before conducting the proof of Theorem 1, a technical lemma is stated first.

Lemma A. Suppose assumptions A1 and A4 are satisfied, and let function $M(t)$ be continuous for all $t \in [0, 1]$. Then, for sufficiently small λ_t ,

$$\int_0^1 G_{\lambda_{t_1}}(t, s) G_{\lambda_{t_2}}(t, s) M(s) f(s) ds = \frac{1}{2\sqrt{2}} (f(t))^{-3/4} M(t) \lambda_{t_1 t_2} (1 + o(1)), \quad (\text{A.1})$$

and

$$\int_0^1 G_{\lambda_t}(t, s) M(s) f(s) ds = M(t) (1 + o(1)). \quad (\text{A.2})$$

for all $t \in [\tau, 1 - \tau]$ with some $\tau > 0$.

Proof: It is easy to see that the quantity $\int_0^1 G_{\lambda_{t_1}}(t, s) G_{\lambda_{t_2}}(t, s) M(s) f(s) ds$ can be expressed as

$$\begin{aligned} & \int_0^1 G_{\lambda_{t_1}}(t, s) G_{\lambda_{t_2}}(t, s) M(s) f(s) ds \\ &= \int_0^1 (G_{\lambda_{t_1}}(t, s) - H_{\lambda_{t_1}}(t, s)) G_{\lambda_{t_2}}(t, s) M(s) f(s) ds + \int_0^1 H_{\lambda_{t_1}}(t, s) (G_{\lambda_{t_2}}(t, s) \\ & \quad - H_{\lambda_{t_2}}(t, s)) M(s) f(s) ds + \int_0^1 H_{\lambda_{t_1}}(t, s) H_{\lambda_{t_2}}(t, s) M(s) f(s) ds. \end{aligned} \quad (\text{A.3})$$

From (17), Lemma 1, and the properties of double exponential distributions, there exists a positive constant c_1 so that, as $\lambda_{l_1} \rightarrow 0$, $\lambda_{l_2} \rightarrow 0$, and $\lambda_{l_1} = O(\lambda_{l_2})$,

$$\begin{aligned}
& \left| \int_0^1 (G_{\lambda_{l_1}}(t, s) - H_{\lambda_{l_1}}(t, s)) G_{\lambda_{l_2}}(t, s) M(s) f(s) ds \right| \\
& \leq \int_0^1 |G_{\lambda_{l_1}}(t, s) - H_{\lambda_{l_1}}(t, s)| |G_{\lambda_{l_2}}(t, s)| |M(s)| f(s) ds \\
& \leq \int_0^1 (\kappa_1)^2 \lambda_{l_2}^{-1/4} \exp\left(-(\alpha_1 \lambda_{l_1}^{-1/4} + \alpha_2 \lambda_{l_2}^{-1/4})|t - s|\right) |M(s)| f(s) ds \\
& = c_1 |M(t)| f(t) (1 + o(1)).
\end{aligned} \tag{A.4}$$

Similarly,

$$\begin{aligned}
& \left| \int_0^1 H_{\lambda_{l_1}}(t, s) (G_{\lambda_{l_2}}(t, s) - H_{\lambda_{l_2}}(t, s)) M(s) f(s) ds \right| \\
& \leq c_2 |M(t)| f(t) (1 + o(1)),
\end{aligned} \tag{A.5}$$

for a positive constant c_2 . Let $u = \Gamma(t)$ and $v = \Gamma(s)$. Again, using the properties of double exponential distributions, it can be shown that

$$\begin{aligned}
& \int_0^1 H_{\lambda_{l_1}}(t, s) H_{\lambda_{l_2}}(t, s) M(s) f(s) ds \\
& = \frac{(\gamma^2 (\lambda_{l_1} \lambda_{l_2})^{-1/4})}{4} \int_0^1 \sin\left(\frac{\pi}{4} + \gamma \frac{\lambda_{l_1} |u - v|}{\sqrt{2}}\right) \sin\left(\frac{\pi}{4} + \gamma \frac{\lambda_{l_2} |u - v|}{\sqrt{2}}\right) \\
& \quad \cdot \exp\left(-\gamma \frac{\lambda_{l_1}^* |u - v|}{\sqrt{2}}\right) M(\Gamma^{-1}(v)) \gamma^{-1} (f(\Gamma^{-1}(v)))^{-3/4} dv \\
& = \frac{1}{2\sqrt{2}} (f(t))^{-3/4} M(t) \lambda_{l_1} \lambda_{l_2} (1 + o(1)),
\end{aligned} \tag{A.6}$$

where $\lambda_{l_1 l_2}^* = \lambda_{l_1}^{-1/4} + \lambda_{l_2}^{-1/4}$. By substituting (A.4), (A.5) and (A.6) into (A.3), (A.1) is then obtained. Similar arguments can be used to show that (A.2) also holds.

APPENDIX B.

Proof of (22)

From (3) and (6), we can derive the equations

$$\begin{aligned} \hat{\beta}_{l(s)}(t; \lambda) - \sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{N s_l^2} \left(\beta_l(t_{ij}) X_{il} - \sum_{l_1 \neq l} B_{l_1(s)}(t_{ij}; \lambda) X_{il_1} \right) S_{\lambda_l, X_l}(t, t_{ij}) \\ = \sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{N s_l^2} \varepsilon(t_{ij}) S_{\lambda_l, X_l}(t, t_{ij}), \quad l = 0, \dots, k, \end{aligned} \quad (\text{B.1})$$

where $B_{l_1(s)}(t_{ij}; \lambda) = (\hat{\beta}_{l_1(s)}(t_{ij}; \lambda) - \beta_{l_1}(t_{ij}))$. By the law of large numbers and assumptions A1, A3, it can be shown that

$$\sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il} X_{il_1}}{N s_l^2} 1_{[t_{ij} \leq t]} \xrightarrow{a.s.} \left(\frac{E[X_l X_{l_1}]}{E[X_l^2]} \right) F(t). \quad (\text{B.2})$$

Using the properties (20)-(21) in Lemma 1, (B.2), assumption A4, and taking expectation of the left hand side of (B.1),

$$\begin{aligned} E \left[\hat{\beta}_{l(s)}(t; \lambda) - \sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{N s_l^2} \left(\beta_l(t_{ij}) X_{il} - \sum_{l_1 \neq l} B_{l_1(s)}(t_{ij}; \lambda) X_{il_1} \right) S_{\lambda_l, X_l}(t, t_{ij}) \right] \\ = B_I + B_{II}, \end{aligned} \quad (\text{B.3})$$

where

$$B_I = E \left[\hat{\beta}_{l(s)}(t; \lambda) \right] - \int_0^1 \beta_l(s) G_{\lambda_l}(t, s) f(s) ds (1 + o(1)),$$

and

$$B_{II} = \sum_{l_1 \neq l} \frac{E[X_l X_{l_1}]}{E[X_l^2]} \int_0^1 \left(E \left[\widehat{\beta}_{l_1(s)}(s; \boldsymbol{\lambda}) \right] - \beta_{l_1}(s) \right) G_{\lambda_l}(t, s) f(s) ds (1 + o(1)).$$

Let $g_l(t) = \int_0^1 G_{\lambda_l}(t, s) \beta_l(s) f(s) ds$. It follows by the definition of the Green function in (16) and Lemma 6.1 of Nychka (1995) that

$$\begin{aligned} g_l(t) - \beta_l(t) &= \frac{-\lambda_l}{f(t)} g_l^{(4)}(t) (1 + o(1)) \\ &= \frac{-\lambda_l}{f(t)} \beta_l^{(4)}(t) (1 + o(1)). \end{aligned} \quad (\text{B.4})$$

Thus, from (B.4) and Lemma A, we can get

$$\begin{aligned} B_I &= E \left[\widehat{\beta}_{l(s)}(t; \boldsymbol{\lambda}) \right] - \left(\beta_l(t) - \frac{\lambda_l}{f(t)} \beta_l^{(4)}(t) + o(\lambda_l) \right) (1 + o(1)) \\ &= \left(E \left[\widehat{\beta}_{l(s)}(t; \boldsymbol{\lambda}) \right] - \beta_l(t) + \frac{\lambda_l}{f(t)} \beta_l^{(4)}(t) \right) (1 + o(1)) \end{aligned} \quad (\text{B.5})$$

and

$$\begin{aligned} B_{II} &= \sum_{l_1 \neq l} \frac{E[X_l X_{l_1}]}{E[X_l^2]} \left(E \left[\widehat{\beta}_{l_1(s)}(t; \boldsymbol{\lambda}) \right] - \left(\beta_{l_1}(t) - \frac{\lambda_{l_1}}{f(t)} \beta_{l_1}^{(4)}(t) + o(\lambda_{l_1}) \right) \right) (1 + o(1)) \\ &= \sum_{l_1 \neq l} \frac{E[X_l X_{l_1}]}{E[X_l^2]} \left(E \left[\widehat{\beta}_{l_1(s)}(t; \boldsymbol{\lambda}) \right] - \beta_{l_1}(t) + \frac{\lambda_{l_1}}{f(t)} \beta_{l_1}^{(4)}(t) \right) (1 + o(1)) \\ &= \sum_{l_1 \neq l} \frac{E[X_l X_{l_1}]}{E[X_l^2]} \left(B \left(\widehat{\beta}_{l_1(s)}(t; \boldsymbol{\lambda}) \right) + \frac{\lambda_{l_1}}{f(t)} \beta_{l_1}^{(4)}(t) \right) (1 + o(1)). \end{aligned} \quad (\text{B.6})$$

By substituting (B.5) and (B.6) into (B.3),

$$\begin{aligned} &E \left[\widehat{\beta}_{l(s)}(t; \boldsymbol{\lambda}) - \sum_{\{i,j\}} \frac{X_{il}}{N s_l^2} \left(\beta_l(t_{ij}) X_{il} - \sum_{l_1 \neq l} B_{l_1(s)}(t_{ij}; \boldsymbol{\lambda}) X_{il_1} \right) S_{\lambda_l, X_l}(t, t_{ij}) \right] \\ &= \sum_{l_1=0}^k \frac{E[X_l X_{l_1}]}{E[X_l^2]} \left(B \left(\widehat{\beta}_{l_1(s)}(t; \boldsymbol{\lambda}) \right) + \frac{\lambda_{l_1}}{f(t)} \beta_{l_1}^{(4)}(t) \right) (1 + o(1)). \end{aligned} \quad (\text{B.7})$$

Since $\varepsilon_i(t)$ is a mean zero stochastic process and is independent of the covariates \mathbf{X} , it implies that

$$E \left[\sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{Ns_l^2} \varepsilon(t_{ij}) S_{\lambda_l, X_l}(t, t_{ij}) \right] = 0. \quad (\text{B.8})$$

Finally, from (B.7) and (B.8), the bias of $\hat{\beta}_{(s)}(t; \lambda)$ in (22) is obtained.

Proof of (23)

Following (B.1), we can get the following equations

$$\begin{aligned} & \prod_{\{l=l_1, l_2\}} \left(\hat{\beta}_{l(s)}(t; \lambda) - \sum_{\{i,j\}} \frac{X_{il}}{Ns_l^2} \left(\beta_l(t_{ij}) X_{il} - \sum_{l_3 \neq l} B_{l_3(s)}(t_{ij}; \lambda) X_{il_3} \right) S_{\lambda_l, X_l}(t, t_{ij}) \right) \\ &= \prod_{\{l=l_1, l_2\}} \left(\sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{Ns_l^2} \varepsilon(t_{ij}) S_{\lambda_l}(t, t_{ij}) \right), \quad \forall l_1, l_2 \in \{0, \dots, k\}. \end{aligned} \quad (\text{B.9})$$

From Lemma 1, (B.2), (B.4), assumptions A1-A4, and Lemma A, the expectation of the left hand side of (B.9) is derived as

$$\begin{aligned} & E \left[\prod_{\{l=l_1, l_2\}} \left(\hat{\beta}_{l(s)}(t; \lambda) - \sum_{\{i,j\}} \frac{X_{il}}{Ns_l^2} \left(\beta_l(t_{ij}) X_{il} - \sum_{l_3 \neq l} B_{l_3(s)}(t_{ij}; \lambda) X_{il_3} \right) S_{\lambda_l, X_l}(t, t_{ij}) \right) \right] \\ &= E \left[\prod_{\{l=l_1, l_2\}} \left(\hat{\beta}_{l(s)}(t; \lambda) - \int_0^1 \beta_l(s) G_{\lambda_l}(t, s) f(s) ds (1 + o_p(1)) \right. \right. \\ &\quad \left. \left. + \sum_{l_3 \neq l} \frac{E[X_l X_{l_3}]}{E[X_l^2]} \int_0^1 B_{l_3(s)}(s; \lambda) G_{\lambda_l}(t, s) f(s) ds (1 + o_p(1)) \right) \right] \\ &= E \left[\prod_{\{l=l_1, l_2\}} \left(\sum_{l_3=0}^k \frac{E[X_l X_{l_3}]}{E[X_l^2]} (B_{l_3(s)}(t; \lambda) + \frac{\lambda_l}{f(t)} \beta_{l_3}^{(4)}(t)) (1 + o_p(1)) \right) \right] \\ &= E \left[\prod_{\{l=l_1, l_2\}} \left(\sum_{l_3=0}^k \frac{E[X_l X_{l_3}]}{E[X_l^2]} (\hat{\beta}_{l_3(s)}(t; \lambda) - E[\hat{\beta}_{l_3(s)}(t; \lambda)]) (1 + o_p(1)) \right) \right] \end{aligned}$$

$$= \sum_{\{l_3, l_4\}}^k \left(\frac{E[X_{l_1} X_{l_3}] E[X_{l_2} X_{l_4}]}{E[X_{l_1}^2] E[X_{l_2}^2]} \right) Cov \left(\hat{\beta}_{l_3(s)}(t; \boldsymbol{\lambda}), \hat{\beta}_{l_4(s)}(t; \boldsymbol{\lambda}) \right) (1 + o(1)). \quad (\text{B.10})$$

Taking expectation of the right hand side of (B.9),

$$E \left[\prod_{\{l=l_1, l_2\}} \left(\sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{N s_l^2} \varepsilon(t_{ij}) S_{\lambda_l, X_l}(t, t_{ij}) \right) \right] = V_I + V_{II} + V_{III}, \quad (\text{B.11})$$

where

$$V_I = E \left[\sum_{i=1}^n \sum_{j=1}^{m_i} \left(\frac{X_{il_1} X_{il_2}}{N^2 s_{l_1}^2 s_{l_2}^2} \right) \varepsilon^2(t_{ij}) S_{\lambda_{l_1}, X_{l_1}}(t, t_{ij}) S_{\lambda_{l_2}, X_{l_2}}(t, t_{ij}) \right],$$

$$V_{II} = E \left[\sum_{i=1}^n \sum_{\{j_1 \neq j_2\}} \left(\frac{X_{il_1} X_{il_2}}{N^2 s_{l_1}^2 s_{l_2}^2} \right) \varepsilon(t_{ij_1}) \varepsilon(t_{ij_2}) S_{\lambda_{l_1}, X_{l_1}}(t, t_{ij_1}) S_{\lambda_{l_2}, X_{l_2}}(t, t_{ij_2}) \right],$$

and

$$V_{III} = E \left[\sum_{\{i_1 \neq i_2\}} \sum_{\{j_1, j_2\}} \left(\frac{X_{i_1 l_1} X_{i_2 l_2}}{N^2 s_{l_1}^2 s_{l_2}^2} \right) \varepsilon(t_{i_1 j_1}) \varepsilon(t_{i_2 j_2}) S_{\lambda_{l_1}, X_{l_1}}(t, t_{i_1 j_1}) S_{\lambda_{l_2}, X_{l_2}}(t, t_{i_2 j_2}) \right].$$

From assumption A4, Lemma 1, and Lemma A, it can be derived that

$$\begin{aligned} V_I &= \frac{E[X_{l_1} X_{l_2}]}{N E[X_{l_1}^2] E[X_{l_2}^2]} \int_0^1 \sigma^2(t) G_{\lambda_{l_1}}(t, s) G_{\lambda_{l_2}}(t, s) f(s) ds (1 + o(1)) \\ &= \frac{1}{2\sqrt{2}N} (f(t))^{-3/4} \left(\frac{\lambda_{l_1 l_2} E[X_{l_1} X_{l_2}]}{E[X_{l_1}^2] E[X_{l_2}^2]} \right) \sigma^2(t) (1 + o(1)), \end{aligned} \quad (\text{B.12})$$

and

$$\begin{aligned} V_{II} &= \left(\sum_{i=1}^n \left(\frac{m_i}{N} \right)^2 - 1/N \right) \left(\frac{E[X_{l_1} X_{l_2}]}{E[X_{l_1}^2] E[X_{l_2}^2]} \right) \int_0^1 \int_0^1 \rho(s_1, s_2) G_{\lambda_{l_1}}(t, s_1) \\ &\quad \cdot G_{\lambda_{l_2}}(t, s_2) f(s_1) f(s_2) ds_1 ds_2 (1 + o(1)) \\ &= \left(\sum_{i=1}^n \left(\frac{m_i}{N} \right)^2 \right) \left(\frac{E[X_{l_1} X_{l_2}]}{E[X_{l_1}^2] E[X_{l_2}^2]} \right) \rho(t, t) (1 + o(1)). \end{aligned} \quad (\text{B.13})$$

Since $\varepsilon_i(t)$ is a mean zero stochastic process and is independent of the covariates \mathbf{X} , it implies that

$$V_{III} = 0 \quad (\text{B.14})$$

Substituting (B.12), (B.13), and (B.14) into (B.11), we can get

$$\begin{aligned}
& E \left[\prod_{\{l=l_1, l_2\}} \left(\sum_{i=1}^n \sum_{j=1}^{m_i} \frac{X_{il}}{N s_l^2} \varepsilon_i(t_{ij}) S_{\lambda_l, X_l}(t, t_{ij}) \right) \right] \\
&= \left(\frac{E[X_{l_1} X_{l_2}]}{E[X_{l_1}^2] E[X_{l_2}^2]} \right) \left(\frac{\lambda_{l_1 l_2}}{2\sqrt{2}N} (f(t))^{-3/4} \sigma^2(t) + \left(\sum_{i=1}^n (m_i/N)^2 \right) \rho(t, t) \right) \\
&\quad \times (1 + o(1)). \tag{B.15}
\end{aligned}$$

From (B.10) and (B.15), the variance of $\hat{\beta}_{(s)}(t; \lambda)$ in (23) is then obtained.

REFERENCES

- Abramovich, F. and Grinshtein, V. (1999). Derivation of equivalent kernel for general spline smoothing: a systematic approach. *Bernoulli*. **5**, 359-379.
- Chiang, C.-T., Rice, J. A. and Wu, C. O. (2001). Smoothing Spline Estimation for Varying Coefficient Models With Repeatedly Measured Dependent Variable. *Journal of the American Statistical Association*. **96**, 605-619.
- Fan, J. Q. and Zhang, J. T. (2000). *Functional linear models for longitudinal data*. *Journal of the Royal Statistical Society*. **B62**, 303-322.
- Hastie, T. J. and Tibshirani, R. J. (1993). Varying coefficient model. *Journal of the Royal Statistical Society*. **B55**, 757-796.
- Hoover, D. R., Rice, J. A., Wu, C. O. and Yang, L.-P. (1998). Nonparametric smoothing estimates of time-varying coefficient models with longitudinal data. *Biometrika*. **85**, 809-822.

- Kaslow, R. A., Ostrow, D. G., Detels, R., Phair, J. P., Polk, B. F. and Rinaldo, C. R. (1987). The multicenter AIDS cohort study: Rationale, organization and selected characteristics of the participants. *American Journal of Epidemiology*. **126**,310-318.
- Nychka, D. (1995). Splines as local smoothers. *Ann. Statist.* **23**, 1175-1197.
- Lin, X. and Carroll, R. J. (2000). Nonparametric function estimation for clustered data when the predictor is measured without/with error. *Journal of the American Statistical Association*. **95**, 520-543.
- Rice, J. A. and Silverman, B. W. (1991). Estimating the mean and covariance structure nonparametrically when the data are curves. *Journal of the Royal Statistical Society*. **B53**, 233-243.
- Umbricht-Schneider, A., Montoya, I. D., Hoover D. R., Demuch, K. L., Chiang, C.-T. and Preston, K. L. (1999). Naltrexone shortened opioid detoxification with buprenorphine. *Drug and Alcohol Dependence*. **56**, 181-190.
- Wahba, G. (1975). Smoothing noisy data with spline functions. *Numer. Math.* **24**, 383-393.
- Wu, C. O. and Chiang, C.-T. (2000). Kernel Estimation and Its Applications of a Varying Coefficient Model with Longitudinal Data. *Statistica Sinica*. **10**, 433-456.
- Wu, C. O., Chiang, C.-T. and Hoover, D. R. (1998). Asymptotic Confidence Regions for Kernel Smoothing of a Varying Coefficient Model with Longitudinal Data. *Journal of the American Statistical Association*. **93**, 1388-1402.

- Wu, C. O., Yu K. F. and Chiang, C.-T. (2000). A Two-Step Smoothing method for Varying Coefficient Models With Repeated Measurements. *Annals of the Institute of Statistical Mathematics*. **52**, 519-543.
- Zeger, S. L. and Diggle, P. J. (1994). Semiparametric models for longitudinal data with application to CD4 cell numbers in HIV seroconverters. *Biometrics*. **50**, 689-699.

Table 1: The real curve $\beta_0(t)$, the averages of 500 estimated curves $\hat{\beta}_{0(c)}(t; 1)$ and $\hat{\beta}_{0(s)}(t; 1)$, and the standard errors of 500 simulation estimates at nine time points.

Time	3.0	6.0	9.0	12.0	15.0	18.0	21.0	24.0	27.0
$\beta_0(t)$	4.517	5.509	6.451	7.321	8.096	8.759	9.292	9.682	9.920
$m(\hat{\beta}_{0(c)}(t; 1))$	4.513	5.499	6.448	7.323	8.103	8.766	9.291	9.660	9.901
(s.d.)	0.096	0.111	0.127	0.140	0.156	0.177	0.187	0.188	0.201
$m(\hat{\beta}_{0(s)}(t; 1))$	4.525	5.509	6.449	7.317	8.091	8.747	9.270	9.652	9.913
(s.d.)	0.013	0.011	0.010	0.011	0.010	0.010	0.010	0.010	0.012

Table 2: The real curve $\beta_1(t)$, the averages of 500 estimated curves $\hat{\beta}_{1(c)}(t; 1)$ and $\hat{\beta}_{1(s)}(t; 1)$, and the standard errors of 500 simulation estimates at nine time points.

Time	3.0	6.0	9.0	12.0	15.0	18.0	21.0	24.0	27.0
$\beta_1(t)$	-0.45	-0.69	-0.93	-1.14	-1.33	-1.49	-1.63	-1.72	-1.78
$m(\hat{\beta}_{1(c)}(t; 1))$	-0.44	-0.68	-0.92	-1.15	-1.35	-1.51	-1.64	-1.71	-1.77
(s.d.)	0.176	0.204	0.233	0.255	0.283	0.319	0.335	0.338	0.364
$m(\hat{\beta}_{1(s)}(t; 1))$	-0.46	-0.70	-0.92	-1.14	-1.32	-1.48	-1.60	-1.69	-1.79
(s.d.)	0.017	0.015	0.015	0.015	0.015	0.014	0.014	0.015	0.017

Table 3: The real curve $\beta_2(t)$, the averages of 500 estimated curves $\hat{\beta}_{2(c)}(t; 1)$ and $\hat{\beta}_{2(s)}(t; 1)$, and the standard errors of 500 simulation estimates at nine time points.

Time	3.0	6.0	9.0	12.0	15.0	18.0	21.0	24.0	27.0
$\beta_2(t)$	0.104	0.148	0.181	0.207	0.225	0.237	0.245	0.248	0.250
$m(\hat{\beta}_{2(c)}(t; 1))$	0.103	0.146	0.181	0.207	0.225	0.235	0.243	0.248	0.251
(s.d.)	0.022	0.027	0.030	0.031	0.033	0.037	0.041	0.042	0.047
$m(\hat{\beta}_{2(s)}(t; 1))$	0.103	0.146	0.181	0.207	0.225	0.237	0.245	0.248	0.250
(s.d.)	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002




Figure 1: The simultaneous smoothing splines $\hat{\beta}_{l(s)}(t; \lambda_l)$ (solid curve) and the component-wise smoothing splines $\hat{\beta}_{l(c)}(t; \lambda_l)$ (dotted curve) with the corresponding 95% bootstrap confidence intervals.




Figure 2: The simultaneous smoothing splines $\hat{\beta}_{l(s)}(t; \lambda_l)$ (solid curve) and the component-wise smoothing splines $\hat{\beta}_{l(c)}(t; \lambda_l)$ (dotted curve) with the corresponding 95% bootstrap confidence intervals.

出席國際會議心得報告

在 2001 年喬治亞州所舉行之美國統計年會，吾參與了統計人之年度盛會並報告了此計劃主題之相關成果，此外，在此會議其間，吾與研究合作者有更進一步之討論及作更深入研究方向之釐定，並聆聽多場統計精英之最近研究成果分享。對於此類大型國際會議之參與，確可增進對不同領域發展之深入瞭解並擴展新視野，希望爾後有機會參與更多此類型之國際會議。