



Two New Species of *Encarsia* (Hymenoptera: Aphelinidae) from Taiwan

L. Y. Chou¹, Y. S. Su¹, K. C. Chou¹, and C. C. Ko²

1. Department of Applied Zoology, Taiwan Agricultural
Research Institute, Wufeng, Taichung, Taiwan, R.O.C.

2. Department of Plant Pathology and Entomology,
National Taiwan University, Taipei, Taiwan, R.O.C.

(Accepted for publication: May 10, 1996)

ABSTRACT

Chou, L. Y., Su, Y. S., Chou, K. C., and Ko, C. C. 1996. Two new species of *Encarsia* from Taiwan. Plant Prot. Bull. 38: 137-142.

Two new species of the genus *Encarsia* Foerster, *E. abundantia* Chou & Su and *E. collecta* Chou & Su, from Taiwan are described and illustrated.

(Key words: *Encarsia*, Aphelinidae, Aleyrodidae, Taiwan)


INTRODUCTION

The species of the genus *Encarsia* Foerster are parasitoids of whiteflies, scale insects and aphids. It contains about 230 species worldwide, including 21 species in Taiwan⁽¹⁻⁵⁾. In this paper 2 new *Encarsia* species are described from Taiwan. The authorship of the name of new species are L.Y. Chou and Y.S. Su. The terminology used in this paper follows Hayat⁽³⁾. The

specimens used in this study are deposited in the Insect Collection of the Taiwan Agricultural Research Institute, Taichung, Taiwan, R.O.C.

***Encarsia abundantia* Chou & Su sp. n.
(Figs. 1-5)**

Female: Head and mesosoma brown to dark brown, side lobe of mesoscutum, axillae, apical scutellum, and median part of metanotum and propodeum yellow to pale



Figs. 1-5. *Encarsia abundantia* Chou & Su sp. n. ♀ : 1, middle tibia and basitarsus; 2, second valvifer (A) and third valvula (B); 3, mesonotum; 4, antenna; 5, fore wing (scale 0.5 mm).

brown; gaster largely yellow to pale brown, except each side of terga I-VI brown; 3rd valvula pale brown to brown. Antenna yellow, scape basally and F6 more or less brown. Fore wing hyaline. Legs pale yellow; fore and hind coxae dark

brown, middle coxa slightly infuscated at base.

Frontovertex 1.3-1.6 times as long as dorsal eye width; mandible tridentate; mouth fossa 0.48-0.57 times as long as frontovertex; eye with pale brown setae;

setae on frontovertex dark brown. Antenna as in Fig. 4; F1 1.3-1.6 times as long as pedicel; F1 3.5-4.5 times as long as wide, 1.1-1.3 times as long as F2, and subequal in width to F2; F2 subequal in length to F3; F3 1.1-1.2 times as long as F4.

Mesonotum as in Fig. 3; mid lobe of mesoscutum with 17-28 dark brown setae; side lobe with 2 pale brown setae; axillae with 1 dark brown seta; distance between anterior pair of scutellar setae shorter than that between posterior pair; scutellar sensilla separated by a distance less than width of one sensillum; fore wing as in Fig. 5; basal cell with 1-2 setae; marginal vein with 7-9 setae; submarginal vein with 2 setae; marginal fringe about 0.20 times as long as fore wing width. Middle tibial spur 0.70-0.86 times as long as middle basitarsus (Fig. 1); hind tibia slightly longer than middle tibia.

Petiole plus gaster about 2 times as long as mesosoma; terga II-VI with 1+1, 1+1, 1+1, 2+2+2, 1+2+1, respectively; tergum VII with 5-9 setae. Ovipositor extends from base of gaster and exerted at apex; 3rd valvula 0.72-0.86 times as long as 2nd valvifer (Fig. 2); ovipositor 2.1-2.4 times as long as middle tibia (Figs. 1, 2).

Male: Unknown.

Material examined: Holotype ♀, TAICHUNG: Wufeng, 23.XI.1994, ex *Dialeurodes tetrastigmae* on *Tetrastigma formosanum* (K.C. Chou). Paratypes: TAICHUNG: Wufeng, 3 ♀, 12.IV.1994, ex *D. kirkaldyi* on *Jasminum sambac* (K.C. Chou); 1 ♀, 24.X.1994, ex *D. kirkaldyi* on *Hiptage benghalensis* (K.C. Chou); 1 ♀, 7. XII.1994, 1 ♀, 25.IV.1995, & 1 ♀, 9. VIII. 1995, ex *D. tetrastigmae* on *Tetrastigma*

formosanum (K.C. Chou). NANTOU: Chunglio, 1 ♀, 30.IX.1994, ex *D. citri* on *Citrus ponk* (K.C. Chou); Lienhuachih, 2 ♀, 14.IV.1994, ex *D. citri* on *Persea* sp. (K.C. Chou); Shuili, 1 ♀, 6.IX.1995, ex *D. dioscoreae* on *Dioscorea* sp. (K.C. Chou).

Distribution: Taiwan.

Hosts: *Dialeurodes citri* (Ashmead), *D. dioscoreae* Takahashi, *D. kirkaldyi* (Kotinsky), *D. tetrastigmae* Takahashi (Aleyrodidae).

Remarks: This species differs from other *Encarsia* species by the mid lobe of mesoscutum with 17-28 setae; 3rd valvula 0.72-0.86 times as long as 2nd valvifer; ovipositor 2.1-2.4 times as long as middle tibia.


Etymology: *Abundantia* means plenty in Latin, refers to its mesoscutum with rather more setae.

Encarsia collecta Chou & Su sp. n.
(Figs. 6-10)

Female: Body brown to dark brown; side lobe of mesoscutum, median part of metanotum and propodeum, petiole, terga I-II and apex of tergum VII yellow to pale brown; 3rd valvula pale brown to brown. Antenna yellow. Fore wing hyaline. Legs yellow.

Frontovertex 4.0-4.2 times as long as dorsal eye width; mouth fossa about 0.30 times as long as frontovertex; eye with pale brown setae; setae on frontovertex brown. Antenna as in Fig. 9; F1 about 0.57 times as long as pedicel; F1 1.0-1.1 times as long as wide, about 0.42 times as long as F2, and 0.78-0.80 times as wide as F2; F2 1.0-1.1 times as long as F3; F3 subequal in length to F4.

Mesonotum as in Fig. 8; mid lobe of



Figs. 6-10. *Encarsia collecta* Chou & Su sp. n. ♀ : 6, middle tibia and basitarsus; 7, second valvifer (A) and third valvula (B); 8, mesonotum; 9, antenna; 10, fore wing (scale 0.5 mm).

mesoscutum longitudinally reticulate, with 11-12 setae; side lobe with 3 dark brown setae; axillae with 1 dark brown seta; distance between anterior pair of scutellar setae more than that between posterior pair; scutellar sensilla separated by a distance more than width of one sensillum. Fore wing as in Fig. 10; basal cell with 3

setae; marginal vein with 10-11 setae; marginal fringe 0.23-0.25 times as long as fore wing width. Middle tibial spur about 0.90 times as long as middle basitarsus; hind tibia distinctly shorter than middle tibia.

Petiole plus gaster about 1.6 times as long as mesosoma. Terga II-VI with 1+1,

1+1, 1+1, 1+2+1, 1+2+1 respectively, tergum VII with 4 setae. Ovipositor extends from tergum IV; 3rd valvula 0.31-0.36 times as long as 2nd valvifer (Fig. 7); ovipositor 1.0-1.1 times as long as middle tibia (Fig. 6, 7).

Male: Unknown.

Material examined: Holotype ♀, TAIPEI: Fushan, 3.VI.1994, ex *Parabemisia myricae* on *Engelhardtia roxburghiana* (K.C. Chou). Paratype: TAIPEI: Mucha, 1 ♀, 7.I.1995, ex *P. myricae* on *Maesa japonica* (K.C. Chou).

Distribution: Taiwan.

Host: *Parabemisia myricae* (Kuwana) (Aleyrodidae).

Remarks: This species differs from other *Encarsia* species by the reticulate sculpture on mesoscutum rather narrow and long.

Etymology: *Collecta* means narrowed in Latin, refers to its reticulate sculpture on mesoscutum rather narrow.

ACKNOWLEDGEMENTS

The authors wish to express their cordial thanks to Mr. Yen-sheue Tseng for the identification of host plants, to Mr. Chen-yu Wong and Miss Chia-hsin Liu for their help in field collection, to Miss Chia-hsin Liu for mounting the specimens. This

study was financially supported (83AST-1.3-FAD-24(7), 84AST-1.3-FAD-15 and 85AST-1.2-FAD-23(1A)) in part by the Council of Agriculture, Executive Yuan, Republic of China.

LITERATURE CITED

1. Chou, K. C., Su, Y. S., Chou, L. Y., and Ko, C. C. 1996. New records of Aphelinidae (Hymenoptera) from Taiwan. J. Agric. Res. China 45: 195-202.
2. Chou, L. Y., and Lin, F. C. 1993. Notes on two parasitoids of *Bemisia tabaci* (Homoptera: Aleyrodidae) from Taiwan. Chinese J. Entomol. 13: 265-267. (in Chinese).
3. Hayat, M. 1989. A revision of the species of *Encarsia* Foerster (Hymenoptera: Aphelinidae) from India and the adjacent countries. Oriental Ins. 19: 163-310.
4. Huang, J. 1994. Systematic studies on Aphelinidae of China (Hymenoptera: Chalcidoidea). Chongqing Publishing House, Chongqing. 348pp. (in Chinese).
5. Silvestri, F. 1930. Contributo alla conoscenza delle specie orientali del genere *Prospaltella* (Hym., Chalcididae). Boll. Lab. Zool. Portici 25: 49-68.

摘 要

周樑鎰¹、蘇怡秀¹、周根清¹、柯俊成² 1996 臺灣恩蚜小蜂屬(膜翅目：蚜小蜂科)二新種 植保會刊 38: 137-142. (¹臺中縣霧峰鄉臺灣省農業試驗所應用動物系；²臺北市國立臺灣大學植物病蟲害學系)

恩蚜小蜂屬(*Encarsia*)隸屬膜翅目蚜小蜂科，目前全世界已知之種類約有 230 種；而臺灣有記錄者僅有 21 種。本文描述臺灣產之 2 新種，並輔以性狀圖說明。

(關鍵詞：恩蚜小蜂屬、蚜小蜂科、粉蝨科、臺灣)