

## Molecular Cloning and Expression of a Phase I Flagellin Gene from *Salmonella enterica* serovar Choleraesuis

Gan-Nan Chang<sup>(1)</sup>, Chen-Shan Chen<sup>(2)</sup> and Kuo-Chieh Ho<sup>(2,3)</sup>

(Manuscript received 10 December, 2006; accepted 16 March, 2007)

**ABSTRACT:** P56, a protein with a molecular weight of 56 kDa, was isolated from the cell-free supernatant of *Salmonella enterica* serovar Choleraesuis CH12440 (*Salmonella* Choleraesuis CH12440) culture. Comparison of the N-terminal amino acid sequence of P56 with the sequences in GenBank indicated that it was flagellin, the product of *fliC* gene and the subunit of flagellar filament. The gene encoding P56 was cloned and expressed. Although the gene displayed a 99.2 % nucleotide sequence homology and a 98.6 % amino acid sequence homology with other *S. Choleraesuis* phase-1-c flagellin genes, there was an insert of 36 nucleotides absent in other phase-1-c flagellin genes. The expressed P56 protein was toxic to macrophage cell Raw264.7, and caused the programmed death of the cells *in vitro*.

**KEY WORDS:** Cytotoxin, Flagellin, *Salmonella enterica* serovar Choleraesuis, Programmed death.

### INTRODUCTION

*Salmonella enterica* serovar Typhi (*S. Typhi*), *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) and *Salmonella enterica* serovar Choleraesuis (*S. Choleraesuis*) are three major pathogenic *Salmonella* species. They are etiological agents of typhoid fever, diarrhea and abdominal pain, and septicemia, enteritis and pneumonia, respectively (Nnalue, 1990; Salyers and Whitt, 1994; Chang and Tsai, 1996). *S. Choleraesuis* normally is a swine pathogen and occasionally causes systemic infection in human (Chang and Tsai, 1996). The endotoxin is considered a major virulence factor of *Salmonella* spp. and is the cause of death in people with systemic infection. Enterotoxin has also been reported for a large number of *Salmonella* serotypes and, at least in part, responsible for the loss of electrolytes and fluid from small intestine (Giannella et al., 1975; Caprioli et al., 1982; Singh et al., 1985).

In the past decade, the cytotoxins produced by enteric pathogens have been increasingly investigated. These toxins have been referred to as verotoxins because they are cytotoxic to Vero cells (Konowalchuk et al., 1977). Some cytotoxins are lethal to mammalian cells, such as Shiga toxin

produced mainly by *Shigella dysenteriae* serotype 1 (Bartlett et al., 1986) and the closely related Shiga-like toxin (SLT) produced by *Escherichia coli* (Cleary et al., 1985; Marques et al., 1986). A similar cytotoxin is produced by *Campylobacter jejuni* (Guerrant et al., 1987). Although evidence has evolved regarding the role of cytotoxins produced by toxigenic organisms in the pathogenesis of particular diseases, very limited data are available concerning the cytotoxin produced by *Salmonella* spp. We previously isolated a cytotoxin from the cell-free supernatant of *S. Choleraesuis* culture, named P56. It caused degeneration and necrosis of mouse macrophage cell line P388-D1 *in vitro*, and hepatomegaly, splenomegaly and pneumonia in ICR mice following intravenous injection (Chang and Tsai, 1996). In this study, we demonstrated that cytotoxin P56 was flagellin, the subunit of flagellar filament.

Flagella are thin, hair-like and rigid appendages of bacteria, and are bacterial locomotive structures. A flagellum has three basic parts: The outmost and longest part is a filament which consists of around 20,000 protein subunits of a single protein called flagellin (FliC) with a molecular weight of 50 to 60 kDa. The flagellin is made within the cell and then passed along the hollow core of the helical filament to be added to the distal end. The filament is attached to a slightly wider and shorter hook which arises from the basal body. The basal body anchors the flagellum to the cell wall and plasma membrane, and is the motor of the flagellum (Kondoh and Hotani, 1974; Macnab, 1996).

1. Department of Biotechnology, Tajen University, 20, Wei-Shin Rd., Shin-II Tsun, Yan-Pu Hsiang, Pingtung 907, Taiwan.
2. Department of Life Science and Graduate Institute of Plant Biology, National Taiwan University, 1, Sec. 4, Roosevelt Rd., Taipei 106, Taiwan.
3. Corresponding author. Tel: 886-2-33662508; Email: kch@ntu.edu.tw



Flagella have been recognized as one of the potential virulence factors of microorganisms. However, most suspicion has been focused on motility necessary for colonization and the aid of virulence factor secretions (Penn and Luke, 1992; Josenhans and Suerbaum, 2002; Hirano et al., 2003). In this communication, we reported the cloning and characterization of the gene encoding P56. The expressed P56 was directly involved in the pathogenesis of bacteria.

## MATERIALS AND METHODS

### Bacterial strain

*Salmonella* Choleraesuis CH12440 was isolated from lungs, hilus lymph node of lungs, spleen, liver, and gallbladder of pigs naturally infected with systemic septicemic salmonellosis (Chang and Tsai, 1996).

### Cell line

Murine macrophage cell line RAW264.7 was purchased from American Type Culture Collection (Manassas, VA) and cultured in RPMI 1640 medium supplemented with 10 % fetal calf serum.

### Amino acid determination

The purified P56 protein showed a single band on a 9% SDS-polyacrylamide gel (SDS-PAGE) was used for amino acid sequence determination. The N-terminal amino acid sequence of protein was determined by Edman degradation using an ABI Procise model 491 protein sequencer (Applied Biosystem, USA) according to the method described by Tam et al. (1998).

### Bacterial genomic and plasmid DNA isolation

Genomic DNA was isolated using the procedure described by Roussel and Chabbert (1978) with some modifications. A 300 mL overnight culture in LB broth (Luria-Bertani broth: 10g bacto-tryptone, 5g bacto-yeast extract, 10 g NaCl per liter) was harvested at 4,000xg. Pellet was suspended in 4 mL of TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0), then 100  $\mu$ L lysozyme (10 mg/mL in TE) and 100  $\mu$ L RNAase A (10 mg/mL in water) were added. The solution was incubated at 37°C for 30 min. Incubation was continued for 1 hr after addition of 1.5 mL proteinase k/sarkosine (5 mg proteinase k and 0.1 g sarkosine/mL TE). After centrifugation at 25,000xg, 20 min, 4 °C, the supernatant was transferred to a new tube and extracted with phenol/chloroform/isoamylalcohol (25:24:1) twice, and then chloroform/isoamylalcohol (24:1) two

times. The aqueous layer was transferred to a new tube and mixed with 0.04 volumes of 5 M NaCl and 2.5 volume of 100% alcohol. The DNA was precipitated at 12,000xg, 4°C, 20 min after chilling at -20°C for 1 hr. The pellet was rinsed with 75% alcohol, air dried and then resuspended in 1.5 mL TE.

Plasmid DNA was prepared by alkaline extraction procedure (Sambrook et al., 1989). For sequencing, plasmid DNA was isolated by QIAprep Spin Miniprep Kit (QIAGEN, Germany).

### DNA probe preparation

After P56 was identified by partial protein sequencing as flagellin, two oligonucleotides (SC1: 5'-TGACCCAGAATAACCTG-3' and SC2: 5'-GAGTCGAGGTCAGACTG-3') were made according to the conserved regions of phase-1 and phase-2 flagellin genes of several *S. Choleraesuis* and *S. Typhimurium* strains (Joys, 1985; Vanegas and Joys, 1995; Wei and Joys, 1985). A 295-bp DNA fragment, named fragment FL295 was amplified on the *S. Choleraesuis* CH12440 genomic DNA using this pair of oligonucleotides in a PCR reaction. The PCR reaction was carried out as described previously (Ho et al., 1992). The fragment was cloned into pUC18 at *Sma*I site and sequenced to confirm the existence of the *fliC* sequence.

### Genomic library construction and screening

Library construction and screening were done as described previously (Ho and Chang, 2000; Ho et al., 2001). The *S. Choleraesuis* CH12440 DNA was partially digested with *Eco*RI and cloned into  $\lambda$ zap (ZAP Express System, Stratagene). The library contained  $1.1 \times 10^5$  plaque-forming unit (pfu) with greater than 95% recombinants determined in the presence of isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) and 5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactoside (Xgal). The library was screened for the *fliC* gene by plaque hybridization using  $^{32}$ P-labeled fragment FL295 as probe. The positive plaque areas were selected and re-screened until a single, isolated plaque could be picked up. A clone named  $\lambda$ SC170 was obtained. The recombinant  $\lambda$  DNA was then converted into the phagemid pSC170.

### Construction of flagellin gene in expression vector

In order to express P56 protein, a PCR DNA fragment containing the coding region of *fliC* was amplified on pSC170 DNA using oligonucleotides of Sal-B and Sal-MS (Sal-B: 5'-GCGGGATCCATGGCA CAAGTAATCAACAC-3' and Sal-MS: 5'-TCC CCCGGGTAAACGCAGTAAAGAGAGGAC-3'). The underlined sequences were the recognition site for



*Bam*HI and *Sma*I, respectively.), and cloned into expression vector pGEX-6P-1 (Amersham Biosciences Ltd) to generate a recombinant plasmid pSC15. The cloned DNA of pSC15 was subjected to sequencing to insure its entirety. The recombinant plasmid was used to transform *E. coli* DH5 $\alpha$ .

#### Expression of the recombinant protein

The recombinant protein was expressed as follows: 2.5 mL of an overnight-culture were added to a 47.5 mL LB broth containing 50  $\mu$ g/mL of ampicillin. The bacteria grew at 37°C until the OD<sub>600</sub> of the culture was 0.7 to 0.9, then an IPTG solution was added to a final concentration of 1 mM. The cultivation was continued for 2 hr. The bacteria were harvested at 10,000xg, 4°C, 10 min and resuspended in 1 mL cold lysis buffer (50 mM NaH<sub>2</sub>PO<sub>4</sub>/300 mM NaCl, pH 8.0) containing 1 mg lysozyme, and then incubated on ice for 30 min. The cells were sonicated on ice for 15 sec each with cooling pause of 15 sec in between at 30 W (Ultrasonic processor, grade 5, Heat systems, NY) until the lysate became transparent and yellowish. The bacterial debris was removed at 10,000xg, 4°C, 30 min. The recombinant proteins were further purified by Glutathione Sepharose 4B column according the instruction of manufacture (Amersham Biosciences Ltd) and/or then cleaved by PreScission protease. Proteins were analyzed on a 12% SDS-PAGE. Proteins were stored at -70°C if they were not used immediately.

#### Purification of cytotoxin, preparation of antibodies, Western blot analysis and DNA sequence determination and cytotoxicity test

These experiments were performed as described previously (Ho et al., 2003).

#### Nucleotide sequence Accession numbers

The nucleotide sequence of *fliC* was deposited in GenBank (accession no. AF159459).

## RESULTS

#### Cloning the gene encoding protein P56

In order to clone the gene coding for P56, the purified protein was subjected to the amino acid analysis by Edman degradation using an ABI Procise model 491 protein sequencer. A sequence of AlaGlnValIleAsnThrAsnSerLeuSerLeuLeuThrGlnAsn from the N-terminus was obtained and demonstrated to be a part of amino acid sequence of flagellin, the product of *fliC* gene and the protein subunit of flagellar filament by comparison with the proteins deposited in GenBank.

A DNA fragment, named fragment FL295, containing 295 nucleotides of *fliC* was amplified in a PCR reaction on *S. Choleraesuis* CH12440 genomic DNA using a pair of primers (SC1 and SC2) designed according to the conserved regions of bacterial *fliC* genes. The fragment FL295 was used as a probe to screen a  $\lambda$ <sub>ZAP</sub> *S. Choleraesuis* genomic library, and a positive clone ( $\lambda$ SC170) containing a 3,787-bp insert DNA was obtained.

Comparison of the nucleotide sequence of the insert DNA with sequences of the genes in GenBank showed that it had a high homology with the sequences of *fliD*, *fliC*, *fliU* and *fliV* genes of *Salmonella enterica* serovar Muenchen (Doll and Frankel, 1993; Ho and Chang, 2000).

#### Characteristics of *fliC* gene


The *fliC* gene contained 1,506 nucleotides encoding a 501-amino acid flagellin protein (FliC) with a predicted molecular weight of 52 kDa and a pI of 4.7. The gene had an ATG as a start codon which was preceded by AGGA Shine-Dalgarno sequence, and flagellar operon-specific -10 (GCCGATAC) and -35 (TAAA) promoter consensus sequences (Kutsukake et al., 1990). Comparison with the *S. Choleraesuis* phase-1-c flagellin genes published by Wei and Joys (1985), the genes had a 99.2 % similarity in nucleotide sequence and 98.6 % in amino acid sequence. The high homology indicated our *fliC* was a phase-1 flagellin gene. However, there was a region of 36 nucleotides absent in other *S. Choleraesuis* phase-1-c flagellin genes (Fig. 1).

#### Expression of the recombinant proteins

*E. coli* DH5 $\alpha$  was transformed with plasmid pSC15 to over-express the recombinant proteins in the presence of IPTG. The cell lysate was analyzed on a 12% SDS-PAGE. Figure 2 showed that a protein with a molecular weight of approximate 56 kDa (named as P56A, lane 4) was detected in the lysate of pSC15-transformed cells, but not in that of pGEX-6P-1 transformed ones. Western blot analysis shown in Fig. 3 indicated that expressed protein P56A could be stained by the antibody raised against purified P56.

#### Cytotoxicity assay for the P56 and P56A proteins

The cytotoxicity of P56 and P56A were tested on mouse macrophage cell line Raw264.7 by the MTT method. The macrophage cells were cultured in RPMI 1640 medium supplemented with 10 % fetal calf serum at 37°C for 24 hr, and then different amounts of P56 or P56A were added. Figure 4 showed that the survival rate decreased significantly in cells treated with P56 and P56A.



```

1---fliD---[CAT]938GCCTTCTTCCTTTTTGATTGCAACACAGTAGTTA571
Met SD
AGCGCGGTTATCGGCAATCTGGAGGCAAAAGTTAATGATAAATTTTGC617
-10 -35
AAAAAATATGCGCGGAATAATGATGCATAAAGCGGCTATTTTCGCC663
CCTAAGAAAAAAGATCGGGGGAAGTGA AAAATTTTCTAAAGTTTCGAA709
-35
ATTCAGGTGCCGATACAAAGGGTTACGGTGAGAAACCGTGGGCAACA755
-10 fliC
GCCCAATAACATCAAGTTGTAATTGATAAGGAAAAGATC794 [ATG] GCA800
Met ala2
CAA GTC ATT AAT ACA AAC AGC CTG TCG CTG TTG ACC836
gln val ile asn thr asn ser leu ser leu leu thr14
CAG AAT AAC CTG AAC AAA TCC CAG TCT GCT CTG GGT872
gln asn asn leu asn lys ser gln ser ala leu gly26
ACC GCT ATC GAG CGT CTG TCT TCC GGT CTG CGT ATC908
thr ala ile glu arg leu ser ser gly leu arg ile40
AAC AGC GCG AAA GAC GAT GCG GCA GGT CAG GCG ATT1844
asn ser ala lys asp asp ala ala gly gln ala ile54
GCT AAC CGT TTC ACC GCG AAC ATC AAA GGT CTG ACT980
ala asn arg phe thr ala asn ile lys gly leu thr66
CAG GCT TCC CGT AAC GCT AAC GAC GGT ATT TCT ATT1016
gln ala ser arg asn ala asn asp gly ile ser ile78
GCG CAG ACC ACT GAA GGC GCG CTG AAC GAA ATC AAC1052
ala gln thr thr glu gly ala leu asn glu ile asn90
AAC AAC CTG CAG CGT GTG CGT GAA CTG GCG GTT CAG1088
asn leu gln arg val arg glu leu ala val gln102
ACT GCT AAC AGC ACC AAC TCC CAG TCT GAC CTC GAC1124
thr ala asn ser thr asn ser gln ser asp leu asp114
TCC ATC CAG GCT GAA ATC ACC CAG CGT CTG AAC GAA160
ser ile gln ala glu ile thr gln arg leu asn glu126
ATC GAC CGT GTA TCC GGT CAG ACT CAG TTC AAC GGC1196
ile asp arg val ser gly gln thr gln phe asn gly150
GTG AAA GTC CTG GCG CAG GAC AAC ACT CTG ACC ATC1232
val lys val leu ala gln asp asn thr leu thr ile168
CAG GTT GGT GCC AAC GAC GGT GAA ACT ATC GAT ATC1268
gln val gly ala asn asp gly glu thr ile asp ile182
GAT CTG AAG CAG ATC AAC TCT CAG ACC CTG GGC CTA1304
asp leu lys gln ile asn ser gln thr leu gly leu174
GAT ACG CTG AAT GTG CAG AAA AAA TAT GAT GTG AGC1340
asp thr leu asn val gln lys tyr asp val ser186
GAT ACT GCT GTA GCT TCC TAT TCC GAC TCG AAA1376
asp thr ala val ala ala ser tyr ser asp ser lys198
CAG AAT ATT GCT GTT CCT GAT AAA ACA GCT ATT ACT1412
gln asn ile ala val pro asp lys thr ala ile thr210
GCA AAA ATT GGT GCA GCA ACC AGT GGT GGT GCT GGT1448
ala lys ile gly ala ala thr ser gly gly ala gly222
ATA AAA GCA GAT ATT AGC TTT AAA GAT GGC AAG TAT1484
ile lys ala asp ile ser phe lys asp gly lys tyr234
TAC GCG ACT GTC AGT GGA TAC GAT GAT GCC GCA GAT1520
tyr ala thr val ser gly tyr asp asp ala ala asp246
ACA GAT AAA AAT GGA ACC TAT GAA GTC ACT GTT GCC1556
thr asp lys asn gly thr tyr glu val thr val ala258

GCA GAT ACA GGA GCA GTT ACT TTT GCG ACT ACA CCA1592
ala asp thr gly ala val thr phe ala thr thr pro270
ACA GTG GTT GAC TTA CCA ACT GAT GCA AAA GCA GTT1628
thr val val asp leu pro thr asp ala lys ala val282
TCA AAA GTT CAA CAG AAT GAT ACT GAA ATA GCA GCA1664
ser lys val gln gln asn asp thr glu ile ala ala294
ACA AAT GCG AAA GCT GCA TTA AAA GCT GCA GGA GTT1700
thr asn ala lys ala ala leu lys ala ala gly val306
GCA GAT GCA GAA GCT GAT ACA GCT ACT TTA GTG AAA1736
ala asp ala glu ala asp thr ala thr leu val lys318
ATG TCT TAT ACA GAT AAT AAT GGC AAA GTT ATT GAT1772
met ser tyr thr asp asn asn gly lys val ile asp330
GGT GGG TTC GCA TTT AAG ACC TCC GGT GGT TAT TAT1808
gly gly phe ala phe lys thr ser gly gly tyr tyr342
GCA GCA TCT GTT GAT AAA TCT GGC GCA GCT AGC TTG1844
ala ala ser val asp lys ala ser gly ala ala ser354
AAA GTT ACT AGC TAC GTT GAC GCT ACC ACT GGT ACC1880
lys val thr ser tyr val asp gln thr thr gly thr366
GAA AAA ACT GCT GCG AAT AAA TTA GGT GGC GCA GAC1916
glu lys thr ala ala asn lys leu gly gly ala asp378
GGT AAA ACC GAA GTT GTT ACT ATC GAC GGT AAA ACC1952
gly lys thr glu val val thr ile asp gly lys thr390
TAC AAT GCC AGC AAA GCC GCT GGC CAC AAC TTC AAA1988
tyr asn ala ser lys ala ala gly his asn phe lys402
GCA CAG CCA GAG CTG GCG GAA GCG GCT GCT ACA ACC2024
ala gln pro glu leu ala glu ala ala thr thr414
ACT GAA AAC CCG CTG CAG AAA ATT GAT GCT GGT TTG2060
thr glu asn pro leu gln lys ile asp ala glu426
GCG CAG GTG GAT GCG CTG CGT TCT CAG CTG GAT GCG2096
val gln val asp ala leu arg ser asp leu gly ala438
GTT CAG AAC CGT TTC AAC TCC GCT ATC ACC AAC CTG2132
thr ser val leu ala gln ala asn gln val pro gln450
AAC GTC CTC TCT TTA CTG CGT TAA TCCGGCGATTGATTC2315
asn val leu ser leu leu arg450
-35
ACCGACACGTGGTACACAATCAAGGCAGCGAAAGCTGCCTTTTTTAA2362
-10
TTCCGGAGCCTGTGTAATG2381---fliU---[TAG]3584TTTTGTTCG3593
SD Met
GGGTGGGCTTTTTCGCGCATCCCGTTTATCCTATCTGGCTGATTTATT3640
CTGCTTTTGAGCCGCTAAAAAGGCGCTACAGGTTACATAAGTGAAA3687
TAACCCCTCTTTTATAGCCTTATTCCTTCGATAGAACCCTCTGTAGA3734
AACGGATAATCATGCCGATAACTCATTTAACCGAGGGGCTGTTATCG3781
TGAATTC3788

```

Fig. 1. The nucleotide sequence of flagellar genes in pSC170. Complete nucleotide and deduced amino acid sequences of *fliC* were shown. ORFs of *fliD* and *fliU* were displayed by broken lines. Numbering of the nucleotide sequence commenced with the *EcoRI* site of insert DNA. Putative -10 and -35 regions of promoters, and Shine-Dalgarno (SD) sequences of each gene were underlined and indicated. The initiation and termination codons were boxed. The sequence of 36 nucleotides in *fliC*, absent from the same genes of other *Salmonella* spp. was underlined (nucleotide 1681 to 1716). The nucleotides of *fliC* in pSC15 different from those in pSC170 were C<sup>806</sup>→A, T<sup>809</sup>→C, T<sup>812</sup>→C, and A<sup>1089</sup>→T.




Fig. 2. Expression of FliC from pSC15 in *E. coli* DH5 $\alpha$ . Proteins were extracted from cells by a buffer containing SDS, separated on a 12% SDS-PAGE and then stained with Coomassie blue. Lane M, protein size markers (kDa). Lane 1, proteins (50  $\mu$ g) from pGEX-6P-1 transformants. Lane 2, proteins (50  $\mu$ g) from pSC15 transformants. Lane 3, fusion protein of GST and P56A purified from pSC15 transformants by a glutathione sepharose 4B column, (20  $\mu$ g). Lane 4, P56A, purified fusion protein of GST and P56A cleaved by PreScission protease (20  $\mu$ g). Expression of proteins was induced by 1mM IPTG for 2 hr.




Fig. 3. Western blot analysis of expressed proteins. Lane M, protein size markers (kDa). Lane 1, cell lysate of *E. coli* DH5 $\alpha$ . Lane 2, cell lysate of *E. coli* DH5 $\alpha$  harboring pGEX-6P-1. Lane 3, cell lysate of *E. coli* harboring pSC15. Expression of proteins was induced by 1mM IPTG for 2 hr. The antibody was prepared against purified P56 from culture supernatant of *S. Choleraesuis*.






Fig. 4. Cytotoxic effect of P56 and P56A on mouse macrophage Raw264.7 cell line. The Raw264.7 cells were cultured in RPMI 1640 medium containing different amounts of P56 and P56A. The effect of protein on the cell viability was determined by the MTT method. P56 protein was isolated from culture supernatant. P56A was isolated from the purified fusion proteins (GST-P56A) cleaved by PreScission protease. The experiments were performed three times in triplicate.

The fusion protein of P56A and glutathione S-transferase (GST) had the same toxicity of P56A (data not shown).

To further characterize the mechanism of cell death resulting from P56, the DNA isolated from P56-treated and untreated Raw 264.7 cells were electrophoresed on a 1% agarose gel. DNA fragmentation in the ladder pattern was noted in P56-treated cells but not clearly identified in untreated ones as shown in Fig. 5. The result suggested that the cell death caused by P56 was involved apoptosis.

## DISCUSSION

The production of toxin is one of major virulence factors involved in the pathogenesis of *Salmonella* spp. The enterotoxins are reported to be associated with some *S. Typhimurium* strains and the symptoms of typhoid fever probably result from the endotoxin, lipopolysaccharide (LPS) of the outer membrane-mediated release of cytokines. However, considerable tissue destruction observed during *Salmonella* infection cannot be explained by damage due to endotoxin or enterotoxin (Salyers and Whitt, 1994; O'Brien and Holmes, 1996). In fact, *Salmonella* cytotoxins have long been suspected. In 1983, Koo and Peterson reported a heat-labile cytotoxin that inhibited protein synthesis in cell-free extract of *Salmonella*. A membrane-bound cytotoxin of *S. Typhimurium* was reported by Reitmeyer et al. (1986). The production of cytotoxins could account for the tissue damage (Reitmeyer et al., 1986). Libby

et al. (1990) cloned a gene encoding cytotoxin with haemolysin activity from *S. Typhimurium*. Kita et al. (1993) reported a cytotoxic protein of 32 kDa isolated from L-form *S. Typhimurium*, with an isoelectric point (pI) of 6.4. This toxin was thermolabile and sensitive to trypsin. It could stimulate macrophages to produce a significant amount of tumor necrosis factor alpha (TNF-alpha) and had a cytolytic effect at concentrations higher than 0.7 µg/mL. Rumeu et al. (1997) characterized another outer membrane-bound, heat-stable cytotoxin produced by *S. enterica* serovar Eenteritidis (*S. Eenteritidis*) and *S. Typhimurium*, that inhibited protein synthesis in culture cells as well as in the intestinal mucosa.




Fig. 5. DNA fragmentation of Raw 264.7 cells. DNA was isolated from Raw 264.7 cells treated (lane 1) or untreated (lane 2) with P56 at a concentration of 80 µg/mL for 72 hr, and then electrophoresed on a 1% agarose gel.

Previously, we have isolated a cytotoxin, P56, with a molecular weight of 56 kDa and a pI of 6.2 from the cell-free supernatant of the *S. Choleraesuis* CH12440 culture. This cytotoxin caused hepatomegaly, splenomegaly and interstitial pneumonia in mice. It also caused degeneration and necrosis of mouse macrophage cell line P388-D1 *in vitro*. A negative Shwartzman's reaction indicated that it was not an LPS endotoxin (Chang and Tsai, 1996). P56 can be detected as early as 3 hr in the supernatant of the culture. The secretion of this protein increased by a factor of 3.2 when the cells were grown in BHI supplemented with 10 mM CaCl<sub>2</sub>, compared to that in the medium without CaCl<sub>2</sub> (Ho et al., 2003).

In this communication, the gene encoding P56 was cloned and characterized. The partial amino acid sequence of P56 indicated that it was the protein flagellin of *S. Choleraesuis* CH12440. A fluorescence confocal microphotograph showed that the periplasm and flagella of *S. Choleraesuis* were stained with the antibody raised against P56 (data not shown). This result agreed with that flagellin is secreted into the periplasm where it is exported to assemble the flagellum.

Flagella are reported not only necessary for motility but also important for the virulence and invasion of *Vibrio cholerae* (Freter and Jones, 1976; Attridge and Rowley, 1983) and *Vibrio vulnificus* (Lee et al., 2004). Drake and Montie (1988) demonstrated that Fla<sup>-</sup> variants of *Pseudomonas aeruginosa* had a diminished virulence and immotile Fla<sup>+</sup> variants lost virulence in animal models. In *C. jejuni*, the flagella played an important role in attachment and invasion (Wassenaar et al., 1991). It was reported that two flagellar genes, *fliA* and *flgM* involved in regulation of the flagellar operon might be involved in regulating the expression of virulence genes in *S. Typhimurium* (Carsiotis et al., 1989; Gillen and Hughes, 1991; Schmitt et al., 1994) and that flagella could help *S. Typhimurium* to survive within murine macrophages (Weinstein et al., 1984). Analysis of nonflagellated *Salmonella* strains revealed a correlation between the ability to induce TNF- $\alpha$  and the expression of the phase-1 filament subunit protein FliC. Furthermore, *Salmonella* FliC could restore the TNF- $\alpha$  inducing phenotype in *E. coli*, which otherwise lacked the activity (Ciacci-Woolwine et al., 1998). Flagellar formation was also suggested to contribute to the virulence of *Erwinia carotovora* subsp. *carotovora* significantly in plants (Matsumoto et al., 2003).

The overlap between motility and the secretion of virulence proteins has been noted in many organisms. There are significant homologies between several flagellar proteins and proteins involved in the expression of virulence genes, the export of virulence proteins, or both, in plant and animal pathogens. The type III secretion mechanism of bacteria exports many virulence effector proteins involved in subversion of eukaryotic cells. The system requires more than 20 proteins, and its core components are homologous to proteins essential for the assembly of flagella (Harshey and Toguchi, 1996; Bennett and Hughes, 2000).

Although motility and flagella were suggested to play role in bacterial virulence, the role of the flagellum in pathogenesis has not yet been proved directly. Our data provide the evidence that flagellum is directly involved in the pathogenesis of bacteria.

Note: The whole genome of a local *S. Choleraesuis* isolate with the insert of 36 nucleotides in *fliC* was sequenced and published after our work was completed (Chiu et al., 2005).

#### ACKNOWLEDGMENTS

We thank Dr. M. F. Tam, Researcher of the Institute of Molecular Biology, Academia Sinica, Taiwan for determining the N-terminal sequences of the purified P56. We also thank Dr. Te-Chang Lee, Researcher of the Institute of Biomedical Science, Academia Sinica, Taiwan for doing DNA autosequencing. Finally, thanks should go to M. Conrad of University of North Carolina at Chapel Hill for his critical reading and suggestions. This work was partially supported by grant NSC 92-2621-B-002-010 from National Science Council of Taiwan to K.-C. Ho.

#### LITERATURE CITED

- Attridge, S. R. and D. Rowley. 1983. The role of the flagellum in the adherence of *Vibrio cholerae*. *J. Infect. Dis.* **147**: 864-872.
- Bartlett, A. V., D. Prado III, T. G. Cleary and L. K. Pickering. 1986. Production of Shiga toxin and other cytotoxins by serogroups of *Shigella*. *J. Infect. Dis.* **154**: 996-1002.
- Bennett, J. C. and C. Hughes. 2000. From flagellum assembly to virulence: the extended family of type III export chaperones. *Trends Microbiol.* **8**: 202-204.



- Caprioli, A., G. D'Agnolo, V. Falbo, L. G. Roda and M. Tomasi. 1982. Isolation of *Salmonella wien* heat-labile enterotoxin. *Microbiologica* **5**: 1-10.
- Carsiotis, M., B. A. Stocker, D. L. Weinstein and A. D. O'Brien. 1989. A *Salmonella typhimurium* virulence gene linked to *flg*. *Infect. Immun.* **57**: 3276-3280.
- Chang, G.-N. and S.-S. Tsai. 1996. Characterization of a cytotoxin "P65" produced by *Salmonella choleraesuis*: (I) Histopathological study on the toxicity of P65 in ICR mice. *J. Chin. Soc. Vet. Sci.* **22**: 331-339. (in Chinese, with English abstract)
- Chiu, C.-H., P. Tang, C. Chu, S. Hu, Q. Bao, J. Yu, Y.-Y. Chou, H.-S. Wang and Y.-S. Lee. 2005. The genome sequence of *Salmonella enterica* serovar Choleraesuis, a highly invasive and resistant zoonotic pathogen. *Nucl. Acids Res.* **33**: 1690-1698.
- Ciacci-Woolwine, F., I. C. Blomfield, S. H. Richardson and S. B. Mizel. 1998. *Salmonella* flagellin induces tumor necrosis factor alpha in a human promonocytic cell line. *Infect. Immun.* **66**: 1127-1134.
- Cleary, T. G., J. J. Mathewson, E. Faris and L. K. Pickering. 1985. Shiga-like cytotoxin production by enteropathogenic *Escherichia coli* serogroups. *Infect. Immun.* **47**: 335-337.
- Doll, L. and G. Frankel. 1993. *fliU* and *fliV*: two flagellar genes essential for biosynthesis of *Salmonella* and *Escherichia coli* flagella. *J. Gen. Microbiol.* **139**: 2415-2422.
- Drake, D. and T. C. Montie. 1988. Flagella motility and invasive virulence of *Pseudomonas aeruginosa*. *J. Gen. Microbiol.* **134**: 43-52.
- Freter, R. and G. W. Jones. 1976. Adhesive properties of *Vibrio cholerae*: nature of the interaction with intact mucosal surface. *Infect. Immun.* **14**: 246-256.
- Giannella, R. A., R. E. Gots, A. N. Charney, W. B. Greenough and S. B. Formal. 1975. Pathogenesis of *Salmonella*-mediated intestinal fluid secretion: activation of adenylate cyclase and inhibition by indomethacin. *Gastroenterology* **69**: 1238-1245.
- Gillen, K. and K. T. Hughes. 1991. Molecular characterization of *flgM*, a gene encoding a negative regulator of flagellin synthesis in *Salmonella typhimurium*. *J. Bacteriol.* **173**: 6453-6459.
- Guerrant, R. L., C. A. Wanke and R. A. Pennie. 1987. Production of unique cytotoxin by *Campylobacter jejuni*. *Infect. Immun.* **55**: 2526-2530.
- Harshey, R. M. and A. Toguchi. 1996. Spinning tails: homologies among bacterial flagellar systems. *Trends Microbiol.* **4**: 226-231.
- Hirano, T., T. Minamino, K. Namba and R. M. Macnab. 2003. Substrate specificity classes and the recognition signal for *Salmonella* Type III flagellar export. *J. Bacteriol.* **185**: 2485-2492.
- Ho, K.-C. and G.-N. Chang. 2000. The *fliU* and *fliV* genes are expressed as a single ORF in *Salmonella choleraesuis*. *Can. J. Microbiol.* **46**: 1149-1152.
- Ho, K.-C., D.-C. Kuo and G.-N. Chang. 2003. Biphasic modulation of mouse macrophage cell growth by the *Salmonella choleraesuis* cytotoxin P56. *Taiwania* **48**: 224-231.
- Ho, K.-C., V. E. Quarmby, F. S. French and E. M. Wilson. 1992. Molecular cloning of rat prostate transglutaminase complementary DNA: The major androgen-regulated protein DPI of rat dorsal prostate and coagulating gland. *J. Biol. Chem.* **267**: 12660-12666.
- Ho, K.-C., C.-C. Tsai and T.-L. Chung. 2001. Organization of ribosomal genes from a loofah witches' broom phytoplasma. *DNA Cell Biol.* **20**: 112-122.
- Josenhans, C. and S. Suerbaum. 2002. The role of motility as a virulence factor in bacteria. *Int. J. Med. Microbiol.* **291**: 605-614.
- Joys, T. M. 1985. The covalent structure of the phase-1 flagellar filament protein of *Salmonella typhimurium* and its comparison with other flagellins. *J. Biol. Chem.* **260**: 15758-15761.
- Kita, E., N. kamikaidou, A. Nakano and S. Kashiba. 1993. Isolation of a cytotoxin from L-form *Salmonella typhimurium*. *FEMS Microbiol. Lett.* **103**: 179-184.
- Kondoh, H. and H. Hotani. 1974. Flagellin from *Escherichia coli* k12: polymerization and molecular weight in comparison with *Salmonella* flagellin. *Biochim. Biophys. Acta* **336**: 117-139.
- Konowalchuk, J., J. I. Speirs and S. Stavric. 1977. Vero response to a cytotoxin of *Escherichia coli*. *Infect. Immun.* **18**: 775-779.
- Koo, F. C. W. and J. W. Peterson. 1983. Cell-free extracts of *Salmonella* inhibit protein synthesis and cause cytotoxicity in eukaryotic cells. *Toxicon* **21**: 309-320.
- Kutsukake, K., Y. Ohya and T. Iino. 1990. Transcriptional analysis of the flagellar regulon of *Salmonella typhimurium*. *J. Bacteriol.* **172**: 741-747.



- Lee, J.-H., J. B. Rho, K. J. Park, C. B. Kim, Y.-S. Han, S. H. Choi, K.-H. Lee and S. J. Park. 2004. Role of flagellum and motility in pathogenesis of *Vibrio vulnificus*. *Infect. Immun.* **72**: 4905-4910.
- Libby, S. J., W. Goebel, S. Muir, J. G. Songer and F. Heffron. 1990. Cloning and characterization of a cytotoxin gene from *Salmonella typhimurium*. *Res. Microbiol.* **141**: 775-783.
- Macnab, R. M. 1996. Flagella and motility. In: Neidhardt, F. C., R. Curtiss III, J. L. Ingraham, E.C.C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter and H. E. Umbarger (eds.), *Escherichia coli* and *Salmonella typhimurium*: cellular and molecular biology, 2nd ed. ASM Press, Washington, D.C., USA. pp. 123-145.
- Marques, L. R. M., M. A. Moore, J. G. Wells, I. K. Wachsmuth and A. D. O'Brien. 1986. Production of Shiga-like toxin by *Escherichia coli*. *J. Infect. Dis.* **154**: 338-341.
- Matsumoto, H., H. Muroi, M. Umehara, Y. Yoshitake and S. Tsuyumu. 2003. Peh production, flagellum synthesis, and virulence reduced in *Erwinia carotovora* subsp. *carotovora* by mutation in a homologue of *cytR*. *Mol. Plant-Microbe Interact.* **16**: 389-397.
- Mosmann, T. 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **65**: 55-63.
- Nnalue, N. A. 1990. Mice vaccinated with a non-virulent, aromatic-dependent mutant of *Salmonella choleraesuis* die from challenge with its virulent parent but survive challenge with *Salmonella typhimurium*. *J. Med. Microbiol.* **31**: 225-233.
- O'Brien, A. D. and R. K. Holmes. 1996. Protein toxins of *Escherichia coli* and *Salmonella*. In: Neidhardt, F. C., R. Curtiss III, J. L. Ingraham, E.C.C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter and H. E. Umbarger (eds.), *Escherichia coli* and *Salmonella typhimurium*: cellular and molecular biology, 2nd ed. ASM Press, Washington, D.C., USA. pp. 2788-2802.
- Penn, C. W. and C. J. Luke. 1992. Bacterial flagellar diversity and significance in pathogenesis. *FEMS Microbiol. Lett.* **79**: 331-336.
- Reitmeyer, J. C., J. W. Peterson and K. J. Wilson. 1986. *Salmonella* cytotoxin: a component of the bacterial outer membrane. *Microbiol. Pathog.* **1**: 503-510.
- Roussel, A. F. and Y. A. Chabbert. 1978. Taxonomy and epidemiology of gram-negative bacterial plasmids studied by DNA-DNA filter hybridization in formamide. *J. Gen. Microbiol.* **104**: 269-276.
- Rumeu, M. T., M. A. Suarez, S. Morales and R. Rotger. 1997. Enterotoxin and cytotoxin production by *Salmonella enteritidis* strains isolated from gastroenteritis outbreaks. *J. Appl. Microbiol.* **82**: 19-31.
- Salyers, A. A. and D. D. Whitt. 1994. Bacterial pathogenesis: A molecular approach. ASM Press, Washington, D. C., USA. pp. 229-243.
- Sambrook, J., E. F. Fritsch and T. Maniatis. 1989. Molecular cloning: A laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY., USA. pp. 125-130.
- Schmitt, C. K., S. C. Darnell, V. L. Tesh, V. A. D. Stocker and A. D. O'Brien. 1994. Mutation of *flgM* attenuates virulence of *Salmonella typhimurium*, and mutation of *fliA* represses the attenuated phenotype. *J. Bacteriol.* **176**: 368-377.
- Singh, S. P., V. D. Sharma and I. P. Singh. 1985. Isolation of an enterotoxic factor from culture supernatant of *Salmonella saintpaul*. *FEMS Microbiol. Lett.* **26**: 301-304.
- Tam, M. F., C.-H. Hsieh, S.-P. Tsai and T.-C. S. Tam. 1998. Amino acid sequence of glutathione S-transferase rGSTM5\* from rat testis. *Biochem. J.* **333**: 735-739.
- Vanegas, R. A. and T. M. Joys. 1995. Molecular analyses of the phase-2 antigen complex 1.2 of *Salmonella* spp. *J. Bacteriol.* **177**: 3863-3864.
- Wassenaar, T. M., N. M. C. Bleumink-Pluym and M. A. M. van der Zeijst. 1991. Inactivation of *Campylobacter jejuni* flagellin genes by homologous recombination demonstrates that *flaA* but not *flaB* is required for invasion. *The EMBO J.* **10**: 2055-2061.
- Wei, L.-N. and T. M. Joys. 1985. Covalent structure of three phase-1 flagellar filament proteins of *Salmonella*. *J. Mol. Biol.* **186**: 791-803.
- Weinstein, D. L., M. Carsiotis, C. R. Lissner and A. D. O'Brien. 1984. Flagella help *Salmonella typhimurium* survive within murine macrophages. *Infect. Immun.* **46**: 819-825.



## 豬霍亂沙門氏桿菌第一型鞭毛素基因的選殖與表現

張甘楠<sup>(1)</sup>、陳真珊<sup>(2)</sup>、何國傑<sup>(2,3)</sup>

(收稿日期：2006年12月10日；接受日期：2007年3月16日)

### 摘 要

P56 是從豬霍亂沙門氏桿菌 CH12440 (*Salmonella Choleraesuis* CH12440) 培養基上清液純化出來，分子量為 56 kDa 的蛋白質。將 P56 的 N 端胺基酸定序並與基因資料庫 (GenBank) 比對，發現它是一種鞭毛素(flagellin)。此蛋白為 *fliC* 基因的產物，是構成鞭毛絲狀體的次單元。我們將 P56 蛋白的基因選殖於大腸桿菌並表現其蛋白。雖然這個基因與其他 *S. Choleraesuis* 第一型(phase-1-c)鞭毛素基因之核苷酸序列有 99.2% 的相似性，胺基酸序列有 98.6% 的相似性，但是在 P56 基因中有一連續 36 個核苷酸是其他 *S. Choleraesuis* 第一型鞭毛素基因所沒有。這個選殖基因表現出來的蛋白質對小鼠巨噬細胞 Raw264.7 具有毒性並會造成細胞的凋亡。

關鍵詞：細胞毒素、鞭毛素、豬霍亂沙門氏桿菌、細胞凋亡。

---

1. 私立大仁科技大學生物科技系暨研究所，907 屏東縣鹽埔鄉新二村維新路 20 號，臺灣。

2. 國立臺灣大學生命科學系及植物科學研究所，106 台北市羅斯福路 4 段 1 號，臺灣。

3. 通信作者。Tel: 886-2-33662508; Email: kch@ntu.edu.tw

