

Didymoplexis micradenia: A Newly Recorded Orchid (Orchidaceae) in Taiwan

Tien-Chuan Hsu⁽¹⁾ and Shih-Wen Chung^(2,3)

(Manuscript received 1 June, 2007; accepted 5 September, 2007)

ABSTRACT: *Didymoplexis micradenia* (Rchb. f.) Hemsl. was recently found to be a new record of *Didymoplexis* Griffith in Taiwan. It is similar to *D. pallens* Griff. except that it is characterized by a smaller flower, very short column foot, and lip apical margin that is minutely denticulate. The description and illustration of *D. micradenia* and a key of *Didymoplexis* in Taiwan are provided in the present work.

KEY WORDS: Orchidaceae, *Didymoplexis micradenia*, new record, Taiwan, Taxonomy.

INTRODUCTION

Didymoplexis (Orchidaceae) is a small genus containing approximately 20 species found in tropical Africa, Madagascar, Southeast Asia, India, Pacific Islands, North Australia, and north to the Ryukyu Islands and Taiwan (Su, 2000; Pridgeon et al., 2005). Like other genera classified within Gastrodieae (Pridgeon et al., 2005), *Didymoplexis* species are all small saprophytic, achlorophyllous plants and emerge aboveground only when flowering and fruiting. They are often neglected by field investigators because of their inconspicuous growth habits and limited flowering seasons. Previously only the type species of *Didymoplexis*, *D. pallens*, was recognized in Taiwan (Su, 2000; Huang, et al., 2003). Some specimens recently collected by the authors were identified to be the new record *D. micradenia*. It is similar to *D. pallens* but differs in floral morphology. In this study we provide the description, line drawing, color photos and distribution map of the new record *D. micradenia*, along with a key to distinguish the genus *Didymoplexis* in Taiwan.

Key to the species of *Didymoplexis* in Taiwan

1. Column with short foot ca. 2 mm long; lip orbitriangular when expanded, more than 6 mm wide, margin nearly rounded or a little undulate. *D. pallens*
1. Column foot very short, less than 1 mm long; lip obovate when expanded, less than 5 mm wide, apical margin minute denticulate. *D. micradenia*

1. Institute of Ecology and Evolutionary Biology, National Taiwan University, 1, Sec. 4, Roosevelt Road, Taipei 106, Taiwan.
2. Division of Forestry Biology, Taiwan Forestry Research Institute, 53, Nanhai Road, Taipei 105, Taiwan.
3. Corresponding author. Tel: 886-2-23039978; Email: biflora@gmail.com


TAXONOMIC TREATMENT

Didymoplexis micradenia (Rchb.f.) Hemsl., J. Linn. Soc. Bot. 20: 311, 1883; Lewis & Cribb, Orch. Vanuatu 1989: 24 — Type: Ovalau, Fiji, *Seemann* 610 (holo. W; iso, K) 小鬼蘭 Figs. 1-3

Epiphanes micradenia Rchb. f. Seem. Fl. Vit. 295, 1868.

Didymoplexis minor J. J. Sm., Bull. Inst. Bot. Buit. 7: 1, 1900; Smith, Die Orchideen von Java. f. LIII; Comber, Orch. Java 1990: 83.

Leafless and achlorophyllous terrestrial herbs. Plants 6-30 cm high. Rhizome subterranean, pale brown, fleshy, fusiform or moniliform, 1-5 cm long, 4-7 mm thick. Roots few, arising from apex of rhizome. Peduncle pale brown to whitish, 5-15 cm long, 1.5-2 mm in diam., glabrous, with 3-5 scarious scales. Raceme 2-5 cm, 8-15 flowered; bract ovate, 1.5 mm long, acuminate; pedicel and ovary 4-6 mm long, glabrous. Flowers whitish, tinged with pale red, not opening widely; sepals and petals connate at base into floral tube; dorsal sepal 7-8 mm long, connate with petal for 1/2 to 2/3 their length, petals shorter, lateral sepals 5-7 mm long, connate for 2/3 to 4/5 their length, whole structure united with petals ca. 1/3 its length, free apex reflexed, 2-lobed, lobules semiorbicular. Lip white, obovate, 4-6 mm long, erect on both sides, 4-5 mm wide when flattened, base cuneate, apex obtuse, minute irregular denticulate in apical margin, disc with longitudinal raised keel of brownish warty papillae. Column white, clavate, ca. 5 mm long, apex enlarged, with 2 truncate rhombic wings, base expanded into inconspicuous short foot less than 1 mm long; anther oval or orbicular, ca. 0.8 mm long, white; pollinia 4, 2






Fig. 1. Distribution map of *Didymoplexis micradenia* (arrowhead / solid circle) and *D. pallens* (gray colored area / open triangle) in the world (A) and in Taiwan (B).





Fig. 2. *Didymoplexis micradenia* (Rchb.f.) Hemsl. A: Habits in different growth phase. B, C: Varied views of flowers. D: Column and lip. E: Lip. F: Connate dorsal sepal and petals. G: Connate lateral sepals. H: Various views of anther. I: Pollinia.






Fig. 3. *Didymoplexis micradenia* (Rchb.f.) Hemsl. A: Habitat and growth habit in fruit. B: Growth habit. C & D: Varied views of flowers. E: Ventral view of column. F: Side view of column. G: Flowers of *D. micradenia* (left) and *D. pallens* (right).

pairs, white, soft, ellipsoid, connected by broad viscidium detached from rostellum. Capsules cylindrical, 1.8-2.2 cm long, triangular in cross section; pedicel lengthening 10-25 cm long in fruit, ca. 2 mm in diam., fleshy.

Specimens examined: Nantou County, Lugu Hsiang, Xidicheng, ca. 300 m in elevation, 2007 May 18, *T. C. Hsu* & *S. W. Chung* 801 (TAIF); Pingtung County, Manjhou Hsiang, Jiutuo, ca. 100m in elevation, 2007 May 26, *T. C. Hsu* & *S. W. Chung* 821 (TAIF); Taitung County, Lanyu Hsiang, Tianchi, mountain ridge, ca. 220 m in elevation, 2004 Apr 13, *S. W. Chung* 8327 (TAIF); Xiaotianchi, ca. 200 m in elevation, 2006 Apr 14, *T. C. Hsu* 469

(TAIF); same loc., 2007 May 25, *T. C. Hsu* 821 (TAIF); Lanyu Lighthouse, ca. 200 m in elevation, 2006 Apr 14, *T. C. Hsu* 470 (TAIF); same loc., 2005 Mar 19, *C. L. Yeh* & *C. R. Yeh* 4032, 4033 (PPI). From roadway to Tianchi, trailside, ca. 150 m in elevation, 2006 Apr 15, *T. C. Hsu* 474 (TAIF); same loc., 2005 Mar 19, *C. L. Yeh* & *C. R. Yeh* 4034, 4035 (PPI); Dongqingxi, ca. 300m in elevation, 2007 May 24, *T. C. Hsu* 815 (TAIF)

Distribution: Samoa, Vanuatu, New Caledonia, Fiji, Tonga, Niue, Palau, West Java and Vietnam; Taiwan, scattered on central and southern lowlands, and the southeast islet of Lanyu.



臺灣大學學術
期刊資料庫

Phenology: Flowering observed from late March to May. Fruiting from April to June.

Notes: In the past *Didymoplexis pallens* Griff., which is distributed in the central to southern part of the main island (Su, 2000), has been the only reported *Didymoplexis* in Taiwan. Recently, few photographs have been taken and specimens collected for the *Didymoplexis* species in Lanyu, but most researchers regarded them as *D. pallens* (Yeh et al., 2006). After thoroughly studying the fresh flowering *Didymoplexis* samples collected in Lanyu, we found that some characteristics differ from those of *D. pallens* in Taiwan. The Lanyu species exhibits a smaller growth habit and flower, narrower and more connate lateral sepal, narrower lip with erose apical margin, and very short column foot. These characters almost agree with the description and line drawing of a Java species *D. minor* (Smith, 1908; Comber, 1990).

Lewis and Cribb (1989) treated *D. minor* as a synonym of *D. micradenia* (Rchb. F.)Hemsl. The description of *D. micradenia* (Hemsley, 1938; Lewis and Cribb, 1989; Cribb and Whistler, 1996) showed similar floral morphology with *D. minor*, except for the almost entirely fused lateral sepal. These differences may be regarded as normal variation between populations. We follow this treatment here and our discovery extends the northern distributional range of *D. micradenia* into Taiwan.

In 2007 we found *D. micradenia* populations again in central and southern Taiwan that differ from the Lanyu populations which grow in the wet rain forests. *Didymoplexis micradenia* in the main island was found mixed with *D. pallens* in seasonal drought forests, or under bamboo plantations. Despite overlapping habitats and flower season, these two species can be clearly distinguished by floral morphology mentioned above, and no transition forms or hybrids were found in the field. Additionally, *D. pallens* has not been found in Lanyu to date. The fruit stalks of *D. micradenia* elongate rapidly after pollination, which is also observed in other *Didymoplexis* and some *Gastrodia* species in Orchidaceae.

ACKNOWLEDGMENTS

We are thankful to Mr. C. R. Yeh for kindly providing useful information and Mr. S. K. Yu for his enthusiastic assistance in field work. We are indebted to Dr. Robert Lauri, Rancho Santa Ana Botanic Garden, for reviewing English writing.

LITERATURE CITED

- Cribb, P. and W. A. Whistler. 1996. Orchids of Samoa. Royal Botanic Gardens, Kew. London, UK. pp. 46-48.
- Hemsley, W. B. 1838. On the synonymy of the orchidaceous genus *Didymoplexis* Griffith, and the elongation of the pedicels of *D. pallens* after flowering. Proceedings of the Linnean Society of London **20**: 309-311.
- Lewis, B. A. and P. Cribb. 1989. Orchids of Vanuatu. Royal Botanic Gardens, Kew. London, UK. p. 47.
- Lin, T.-P. 1987. Native Orchids of Taiwan, vol. 3. Southern Materials Center, Taipei, Taiwan. pp. 59-61
- Pridgeon, A. M., P. J. Cribb, M. W. Chase and F. N. Rasmussen. 2005. Genera Orchidacearum, Vol. 4, Epidendroideae (part 1). Oxford University Press, Oxford, UK. pp. 435-437
- Yang, Y.-P. and H.-Y. Liu. 2002. Manual of Taiwan Vascular Plants, vol. 6. Council of Agriculture, Taipei, Taiwan. p. 387.
- Yeh, C.-L., C.-R. Yeh and C.-S. Leou. 2006. An Observation on the *Corymborkis veratrifolia* (Reinw.) B1. (Orchidaceae) from Lanyu, Taiwan. *Taiwania* **51**: 53-57.
- Smith, J. J. 1908. Die Orchideen von Java. Figuren-Atlas. Leiden, Netherlands. p. 43
- Su, H.-J. 2000. In: Huang, T.-C. et al. (eds.), Orchidaceae. Flora of Taiwan, 2nd ed. **5**: 851-854. Editorial Committee, Dept. Bot., NTU, Taipei, Taiwan.



臺灣新紀錄蘭科植物：小鬼蘭

許天銓⁽¹⁾、鐘詩文^(2,3)

(收稿日期：2007年6月1日；接受日期：2007年9月5日)

摘 要

近年我們在臺灣中南部與蘭嶼發現了臺灣蘭科的新紀錄種：小鬼蘭 *Didymoplexis micradenia* (Rchb. f.) Hemsl.。本種近似吊鐘鬼蘭 *D. pallens* Griff.，但由唇瓣較窄且末端具細齒緣、蕊柱足部不顯著等特徵可清楚分辨。本文並提供描述、圖片與臺灣鬼蘭屬之檢索表。

關鍵詞：蘭科、小鬼蘭、新記錄、分類、臺灣。

1. 行政院農業委員會林業試驗所生物組，100 台北市南海路 53 號，臺灣。

2. 國立臺灣大學生態學與演化生物學研究所，106 台北市羅斯福路 4 段 1 號，臺灣。

3. 通信作者。Tel: 886-2-23039978; Email: biflora@gmail.com

