

Anadromy and the dispersal of an invasive fish species (*Oncorhynchus mykiss*) in Eastern Quebec, as revealed by otolith microchemistry

Thibault I, Hedger RD, Dodson JJ, Shiao J-C, Iizuka Y, Tzeng W-N. Anadromy and the dispersal of an invasive fish species (*Oncorhynchus mykiss*) in Eastern Quebec, as revealed by otolith microchemistry. Ecology of Freshwater Fish 2010. © 2010 John Wiley & Sons A/S

Abstract – Rainbow trout (*Oncorhynchus mykiss*) is invading rivers bordering the St. Lawrence Estuary (Quebec, Canada). Some rivers in Eastern Quebec support self-sustaining populations while adult vagrants are frequently captured in rivers where no reproduction has been confirmed. We hypothesised that the development of anadromy has promoted the species dispersal. Otolith Sr:Ca analyses revealed that although all fish captured in the upstream stocking region were freshwater residents, both anadromous and freshwater resident phenotypes were found downstream in Eastern Quebec. The proportion of fish exhibiting the anadromous life cycle increased with the distance from the stocking zone. Eastern Quebec steelhead migrated to sea at the same age but at a larger size than steelhead within their native range. Age at first reproduction was similar to that observed in native populations. The development of the anadromous life cycle enables this species to colonise new rivers following long-distance migrations along the St. Lawrence Estuary corridor.

**I. Thibault¹, R. D. Hedger²,
J. J. Dodson¹, J.-C. Shiao³,
Y. Iizuka⁴, W.-N. Tzeng⁵**

¹Département de biologie, Québec-Océan, Université Laval, Québec, 1045 Avenue de la Médecine, Québec, Canada G1V 0A6, ²Norwegian Institute for Nature Research, NO-7485 Trondheim, Norway, ³Institute of Oceanography, College of Science, National Taiwan University, Taipei, Taiwan 106, ROC, ⁴Institute of Earth Sciences, Academia Sinica, 128, Section 2, Academia Road, Nankang, Taipei, Taiwan 115, ROC, ⁵Institute of Fisheries Science, College of Life Science, National Taiwan University, Taipei, Taiwan 106, ROC

Key words: exotic fish invader; hierarchical filter; dispersion capacity; anadromy; freshwater residency; otolith microchemistry

Isabel Thibault, 1045 Avenue de la Médecine, Département de biologie, Québec-Océan, Université Laval, Québec, Québec, Canada G1V 0A6; Tel.: +1 418 656 2131X7862, Fax: +1 418 656 2339; e-mail: isabel.thibault.2@ulaval.ca

Accepted for publication February 10, 2010

Introduction

Once established in a new habitat, the success of an exotic invader will depend on the species' capacity to disperse in that habitat and to increase its population size (Theoharides & Dukes 2007). This is the last of the hierarchical filters which determine the success of biological invasions (Kolar & Lodge 2001; Colautti et al. 2006; Moyle & Marchetti 2006; Theoharides & Dukes 2007). Characteristics associated with invaders capable of surpassing this last filter are numerous, including long-distance dispersal capacities, small body size, high fecundity, asexual reproduction, rapid growth and early maturity (Kolar & Lodge 2001; Moyle & Marchetti 2006; Theoharides & Dukes 2007).

Rainbow trout (*Oncorhynchus mykiss*, Walbaum 1792) is one of the most widely introduced fish species in the world (GISD 2005). Stocking of this salmonid,

originating from the Northern Pacific Ocean, has occurred throughout the world since 1874 with varying levels of success (MacCrimmon 1971; Burgner et al. 1992). Like several of its relatives of the Salmonidae family, rainbow trout may adopt two life-cycle forms: the anadromous phenotype, with a growth phase in saltwater, or the freshwater resident phenotype. Despite a prevailing tendency toward philopatry (Burgner et al. 1992), the anadromous phenotype may nevertheless migrate into new streams, hence increasing the species' capacity to disperse. Anadromy also implies larger size at reproduction for males and females. Larger body size is generally related to higher fitness, particularly among females who produce more and larger eggs with increasing body size (DuBois et al. 1989; Seamons et al. 2007 and references therein). However, the few naturally spawning populations of rainbow trout that have been established after an introduction event are usually

composed of freshwater residents. The introduced self-sustaining population in the Santa Cruz River (Southern Patagonia) is the only one known to have developed an anadromous run (Pascual et al. 2001; Riva Rossi et al. 2004). The reason for the dominance of residency among introduced populations is not clear. It may be related to the phenotype of the introduced populations, which is usually resident. It may also be that the costs associated with anadromy in recently founded populations are higher than for residency, resulting in selection against migrants.

In Eastern Canada, stocking of rainbow trout has been conducted for several years in the upper St. Lawrence River system, from the Great Lakes to the vicinity of Quebec City, leading to the establishment of some self-sustaining populations in Eastern Quebec outside the area where stocking and farming of rainbow trout are permitted (Thibault et al. 2009). The first such population was discovered in the 1980s and reproduces in two rivers, Du Gouffre and Malbaie, located in the Charlevoix region, on the north shore of the St. Lawrence Estuary. In 2007–2008, young of the year (0+) and other juveniles were also observed in two rivers (Matane and Les Méchins) located downstream, on the south shore of the St. Lawrence Estuary. Other than these established populations, adults have been frequently captured during many years in streams all along the estuary. Given that no reproduction has been detected in these rivers, captures are believed to be composed of vagrant fish. Genetic analyses revealed that such vagrants mainly originate from a population founded in Lake Ontario, and also from the recently established population in Charlevoix (Thibault et al. 2009). These findings suggest the presence of anadromous individuals in the population of the Malbaie and Du Gouffre rivers, which contradicts the belief that introduced self-sustaining populations of rainbow trout only support the freshwater resident form (Pascual et al. 2001; Riva Rossi et al. 2004).

The main objective of this study was to demonstrate that anadromy is involved in the colonisation process of Eastern Quebec by rainbow trout. We aimed to determine if the anadromous phenotype is present in recently naturalised populations in Eastern Quebec, which are believed to consist of mainly freshwater residents. We also tested the hypothesis that all vagrants should be anadromous. To determine the occurrence of anadromy, both direct (i.e., individual migratory behaviour) and indirect (maternal phenotype) observations were used. Our second objective was to characterise the anadromous phenotype in Eastern Quebec rivers. We thus determined the age and size at migration and at reproduction for steelhead (anadromous rainbow trout), and compared growth rates between the anadromous and resident phenotypes. Finally, we compared these characteristics with

what is observed in populations found within their native range.

Method

Sampling and measurements

In 2005 and 2006, we obtained more than 200 rainbow trout from different sites in Quebec waters, including rivers, lakes, creeks and the St. Lawrence River and Estuary. Most rainbow trout (77%) were provided by recreational fishers, whereas the remaining specimens came from commercial fisheries, the monitoring of dams and surveys performed by biologists and river managers. The majority of fish were captured from June to August. As fishing effort varied between sample locations, our sample did not reflect the real distribution and abundance of the species.

Fish were weighed, measured (fork length), and sexed. As biological information was mainly provided by fishers, some data were missing (especially mass) and their precision is somewhat uncertain. When possible, size and mass were measured again and sex was confirmed in the laboratory. A tissue sample was collected to determine if fish captured outside the stocking area originated from a domestic sterile strain used for stockings (triploid fish – 3N) or a wild population [refer to Thibault et al. (2009) for more details on the method]. Sagittal otoliths were extracted, cleaned with distilled water, dried in air and conserved in plastic Eppendorf tubes until otolith microchemistry analyses were performed (see section ‘Otolith Sr:Ca analysis and interpretation’). For the purpose of this study, only 140 specimens were kept for the otolith microchemistry analyses. Our selection criteria were: (i) the capture location, in order to represent different habitats, favouring fish captured to the east of Quebec City, (ii) the fish size, as larger individuals were believed to be older and therefore more likely to present a more complete life cycle, and (iii) the fish sex, to ensure a sample with an almost equal sex ratio. Ten farmed rainbow trout (3+ years of age) provided by a governmental hatchery (*Lac-des-Écorces*, LDE) were also analysed.

Otolith Sr:Ca analysis and interpretation

Samples preparation for electron probe micro-analysis of otoliths was performed in National Taiwan University. Quantitative chemical analyses of strontium (Sr) and calcium (Ca) in otoliths were carried out by an electron probe micro analyser (EPMA: JEOL JXA-8900R, JEOL, Akishima, Tokyo, Japan) and a wavelength dispersive spectrometer (WDS) at the Institute of Earth Sciences, Academia Sinica in Taipei. Secondary- and backscattered electron images were used to guide

the analysis on target positions located along an axis from the core of the otolith to the edge at 10- μ m intervals. Beam conditions were 15 kV for the acceleration voltage and 3 nA for the current, and a $5 \times 4 \mu$ m rectangular scanning beam was used. The Sr $L\alpha$ peak position was measured for 80 s and each of the upper and lower baselines for 20 s. The peak concentration of Ca $K\alpha$ was measured for 20 s and each of the upper and lower baselines for 10 s. The beam power density of 2.25 Wm^{-2} was within the range of 0.5–3 Wm^{-2} as suggested by Gunn et al. (1992), and the counting time was sufficient to ensure effective measurement.

The measured X-ray intensities were corrected by the PRZ [phi-rho-z, Reed (1993)] method using the standard calibration of natural and synthetic chemical-known standard minerals; a strontianite [(Sr0.95Ca0.15)CO₃; NMNH-R10065 from National Museum of Natural History, Smithsonian Institution, Washington, DC, USA] for Sr and a synthesised aragonite (CaCO₃) for Ca. Detection limits based on 3 σ of standard calibration were less than 400 ppm for Sr.

As the otolith Sr:Ca ratio increases with salinity (Zimmerman 2005), it is possible to distinguish periods when the fish grew in freshwater or the sea. Among the 150 otoliths selected for the analysis (140 captures + 10 farmed specimens), some broke during

the preparation or presented vaterite inclusions (Tzeng et al. 2007; Jessop et al. 2008), and so were discarded. Interpretation of the otolith Sr:Ca life-history transects was possible for 125 specimens, representing 29 different watercourses (and the LDE government hatchery, $N = 8$), distributed over three sectors: (i) South-western Quebec (SWQ), including six stocked creeks and lakes in the upper part of the St. Lawrence River, (ii) Quebec City (QC), comprising ten rivers located within 70 km of Quebec City, at the downstream limit of the stocking area, and (iii) Eastern Quebec (EQ), consisting of 13 rivers located outside the stocking area (Fig. 1). Following analysis, one specimen was discarded (see below).

Among fish captured inside the stocking area, 41 were *a priori* known to be freshwater residents. Twenty-one were farmed fish that either came directly from the LDE government hatchery ($N = 8$) or were captured in lakes shortly after stocking events (SWQ, $N = 13$). Another 20 specimens were caught upstream of two impassable waterfalls (QC, $N = 13$) or came from watercourses located far away from the nearest marine habitat (SWQ, $N = 7$) (Fig. 1). All these fish were used to establish the threshold demarcating fresh from marine (or brackish) water habitats (thereafter called FW-threshold). Their mean otolith Sr:Ca values varied from 0.327 to 3.684×10^{-3} , with an average of




Fig. 1. Capture locations of the 116 rainbow trout (*Oncorhynchus mykiss*) analysed for otolith microchemistry (the eight fish from the *Lac-des-Écorces* hatchery are not presented). Numbers in the circles represent the sample sizes. Specimens ($N = 33$) used for determination of the threshold between freshwater and saltwater signatures are shown in white. The area where stocking and farming of rainbow trout is allowed is shown in dark grey. Quebec City is represented by an asterisk. SWQ: South-western Quebec region, QC: Quebec City region (encircled), and EQ: Eastern Quebec region.

1.517×10^{-3} . Mean Sr:Ca + 1 SD was $0.790-3.724 \times 10^{-3}$, with an average of 2.528×10^{-3} (Fig. 2). More than 90% of all Sr:Ca values were less than or equal to 3×10^{-3} . To be conservative, we decided to set the FW-threshold at 3.5×10^{-3} , which included 96% of the freshwater observations.

The phenotype of the 84 remaining rainbow trout (QC and EQ sectors) was determined by two methods. The first method consisted of visually interpreting the otolith Sr:Ca transect of each individual from a point situated $100 \mu\text{m}$ from the otolith's primordium to the otolith edge. Fish were classified as anadromous (A) – or steelhead – if the two-point smoothed average curve exceeded the FW-threshold of 3.5×10^{-3} anywhere along the otolith transect from $100 \mu\text{m}$ outward. Fish were classified as freshwater resident rainbow trout (FW) when all values of the two-point smoothed average were less than the FW-threshold anywhere along the otolith transect from $100 \mu\text{m}$ outward. Note that when only one point was located above or under the FW-threshold, it was not taken into account. At distances less than $100 \mu\text{m}$ from the primordium, Sr:Ca values represented the maternal contribution (C. Donohoe, pers. comm., Institute of Marine Science, UCSC). If a gravid anadromous female does not stay in a freshwater spawning river for a long period before egg deposition, then a marine signal transmitted to the offspring otolith through the egg cytoplasm may be observed near the otolith core (Rieman et al. 1994; Volk et al. 2000; Zimmerman & Reeves 2000, 2002; Donohoe et al. 2008). Fish exhibiting high Sr:Ca values between 0 and $100 \mu\text{m}$ were therefore classified as the offspring of steelhead. Note that it was not possible to determine the maternal form for two rainbow trout because the Sr:Ca transect did not begin at the otolith primordium.

Despite the fact that visual interpretation of Sr:Ca patterns is common in studies using Sr:Ca otolithom-

etry (Tzeng et al. 2003; Gillanders 2005; Daverat et al. 2006; e.g., Tzeng et al. 2000; Zimmerman & Reeves 2000), the method is quite subjective. More quantitative approaches have been recently proposed to avoid any reader bias (e.g., Hedger et al. 2008). To validate results obtained with the conventional method, we also employed a quantitative approach based on nonparametric smoothing to interpret the otolith Sr:Ca transects. Otolith sequences were smoothed using generalised additive models (GAMs). GAMs were used for modelling the relationship between Sr:Ca ratio and distance from the otolith's primordium because there was no *a priori* reason to choose between the forms of the relationship. Use of GAMs enabled this relationship to be determined from the data (instead of being arbitrarily imposed). GAMs were fitted using the *gam(mgcv)* package of R (Wood 2006), using penalised regression splines. The degree of smoothness of the model term was estimated as part of fitting using generalised cross validation. GAMs were fitted separately to the maternal and offspring parts of the transect sequence. Determination of the transition from maternal to offspring parts of the sequence was based on the fact that the maternal signature could have been a mix of freshwater and saltwater Sr:Ca ratios, but that the initial offspring signature must have been a freshwater Sr:Ca ratio due to spawning in freshwater. The procedure for defining the break was as follows: (i) a distance of $100 \mu\text{m}$ was defined as the initial break-point between maternal and offspring histories, (ii) an offspring GAM was fitted to all parts of the sequence at distances greater than the break-point (thus excluding the maternal history), (iii) if the part of the offspring GAM nearest to the primordium showed a saltwater signature (predicted values greater than the FW-threshold), then an extra $10 \mu\text{m}$ was added to the break-point, (iv) steps 2–3 were repeated until the part of the offspring GAM




Fig. 2. Sr:Ca pattern (two-point smoothed averages) of 41 rainbow trout with a known freshwater residence phenotype. The full line corresponds to the mean Sr:Ca values, and the dashed line corresponds to the mean + 1 SD. The threshold of 3.5×10^{-3} , delineating freshwater and saltwater, is shown by a dotted horizontal line. The vertical line represents the distance from the otolith primordium beyond which the number of specimens is less than 30% of the sample.

nearest to the core showed a freshwater signal, and (v) a maternal GAM was fitted to all parts of the sequence nearer to the primordium than the part of the sequence used to determine the offspring GAM. Otolith histories (maternal or offspring) were defined as being anadromous if any value predicted by the GAM exceeded the FW-threshold.

Comparing interpretations obtained with qualitative and quantitative methods, we found 92% and 87% of concordance for fish phenotypes and maternal forms respectively. Discordant interpretations ($N = 18$) consisted mainly (83%) of anadromous categorisations according to the visual method as opposed to a freshwater residency classification obtained by the GAM method. Final categorisation followed the GAM's interpretation except in cases where obvious evidence (e.g., discordance between signal and capture location of the fish, abnormally low Sr:Ca values for an entire transect) supported the visual method result. Note that one fish classified as anadromous according to both methods was finally removed from the analysis as we suspected an edge effect (Jessop et al. 2002): it presented a sudden saltwater Sr:Ca signal at the very end of the otolith transect, despite being captured in freshwater.

Life history of established versus invading rainbows

Inside the stocking zone, where the majority of captured rainbow trout are believed to be cultured fish, all specimens should originate from a freshwater resident mother and present a freshwater residency phenotype. On the contrary, outside the boundaries of the stocking area, the FW-resident phenotype is expected only among established fish found in the two rivers where reproduction has been confirmed (i.e., Du Gouffre and Malbaie). All rainbows captured elsewhere, where the species is not believed to reproduce, were considered as vagrants. They thus should have experienced at least one marine phase representing the migration from their native stream to the river of capture using the St. Lawrence River (≥ 20 PSU in this area, Lecomte & Dodson 2004), a phase that should be reflected along the otolith transect by higher Sr:Ca values. Capture of putative vagrants with a solely freshwater signature would thus suggest that the species is reproducing in more rivers than documented to date. For both established fish and vagrants, the two maternal forms (resident or anadromous) are possible.

Age and size at migration and growth rate

Farmed strains are genetically selected to increase growth rates, and rearing conditions in hatcheries are designed to favour growth. Therefore, growth rates of farmed fish differ to those of fish originating from

naturalised populations. Because fish found inside the stocking area limits probably originated in a hatchery, only fish from Eastern Quebec (Charlevoix, Bas-St-Laurent and Gaspesia regions) were kept for size and growth analyses. The age of each remaining specimen ($N = 51$) was determined by two readers from otolith photographs (obtained after microprobe analyses and otoliths etched with 0.05 N HCl) by counting annuli. Given that the locations of the microprobe Sr:Ca measurements were visible on the otolith photos, it was possible to identify the approximate position of each annulus on the Sr:Ca transects, and therefore determine the age of fish at migration in the case of anadromous fish.

To estimate size at migration of steelhead, we back-calculated fish size to the age at habitat switch, as determined from otolith Sr:Ca ratios, according to the 'body proportional hypothesis' (BPH) technique (Francis 1990; Jessop et al. 2004; Brisson-Bonenfant 2006). The BPH equation is as follows:

$$\log_{10}L_i = \left[\frac{(c + d\log_{10}O_i)}{(c + d\log_{10}O_c)} \right] \log_{10}L_c \quad (1)$$

where c is the intercept and d is the slope of the body length–otolith radius (BL–OR) regression ($\log_{10}BL = 1.33 \times \log_{10}OR + 2.35$; $N = 93$, $R^2 = 0.82$), L_i and O_i are body length and otolith radius at age i , and L_c and O_c are body length and otolith radius at capture. Otolith radius and distances between annuli and the primordium were measured along the longest radius along the ventral side with image analysis software Image-Pro Express version 6.0 (Media Cybernetics Inc., Bethesda, MD, USA).

The size-at-age relationship was determined for both anadromous and resident fish in order to compare growth rates between life-history forms. Dates of capture varied among specimens, and current year growth was visible for some fish. To avoid bias, we back-calculated the size of each fish to their last visible annulus (i.e., to the spring preceding capture), instead of using size at capture.

Results

General characteristics of sampled fish

Specimens used for Sr:Ca analyses (excluding fish from the LDE hatchery) showed a wide range of sizes and masses (Table 1). The overall sex ratio in our sample was 1.6, with a bias toward females. As expected, this bias was much more pronounced inside the stocking area (SWQ and QC) because when strains used for stocking are not triploid (and hence sterile), they often consisted of *all-female* lineages. Rainbow trout captured outside the stocking area, in Eastern Quebec (EQ), were all believed to be vagrants, except

Table 1. Biological characteristics of 116 rainbow trout captured in Quebec's streams in 2005–2006 and used for otolith microchemistry analyses, including 33 specimens selected to determine the threshold between freshwater and saltwater signatures, but excluding the specimen eliminated because of otolith edge effect.

Sector of capture	Sex	N	Fork length (mm)		Mass† (g)	
			Mean	Range	Mean	Range
South-western Quebec (SWQ) (inside stocking area)	F	10	321	229–406	513 (9)	294–1361
	M	3	274	250–323	223 (2)	220–225
	I	1	229	–	?	–
	3N	6	287	210–559	504	100–2381
Quebec City (QC) (inside stocking area)	F	17	245 (16)	119–495	89 (10)	20–291
	M	7	179	124–260	69	24–113
	I	5	229 (3)	229	135 (3)	135
	3N	16	270 (15)	232–308	174 (11)	113–340
Eastern Quebec (EQ) (outside stocking area)	F	24	343	176–334	596 (21)	63–1364
	M	22	286	173–585	325 (20)	54–1361
	I	5	235	90–317	279 (2)	227–331

†Some fish were weighed after thawing.

Numbers in parentheses correspond to the N used for calculating means when some data were missing.

F: female, M: male, I: sex not determined, 3N: triploid farmed fish.

those coming from Du Gouffre and Malbaie Rivers, where natural reproduction is documented. In our sample, the oldest rainbow trout was 9-years old, whereas the youngest one was 1-year old (Table 2). On average, fish from the established population were older than those thought to be vagrants.

Life history patterns: fish phenotype and maternal form

Four main otolith Sr:Ca life history patterns were observed: (i) FW-resident phenotype with a FW-resident maternal form (FW/FW, Fig. 3a, b), (ii) FW-resident phenotype with an anadromous maternal form (A/FW, Fig. 3c, d), (iii) anadromous phenotype with a FW-resident maternal form (FW/A, Fig. 3e, f), and (iv) anadromous phenotype with an anadromous maternal form (A/A, Fig. 3g, h). Fish sampled in the stocking area showed the same life history phenotype as their mother whereas this was not the case outside the stocking area in Eastern Quebec (Table 3). Among fish sampled in Eastern Quebec, 38% of FW-residents originated from an anadromous mother, whereas 42% of anadromous trout originated from a FW-resident mother.

Occurrence of anadromy varied according to the capture location (Table 3 and Fig. 4). Inside the stocking area, that is, the Quebec City region, the

FW-resident phenotype dominated the sample, and no specimen originated from a steelhead mother. In contrast, outside the stocking area (Eastern Quebec), the signal of anadromy, expressed either by the specimen's phenotype or the maternal form, dominated the sample. Based on a Chi-square analysis of the incidence of the saltwater signal as a function of distance from the stocking zone (Charlevoix, Bas-St-Laurent, Gaspesia), the proportion of fish having experienced saltwater increased downstream ($\chi^2 = 8.87$, $N = 51$, d.f. = 2, $P = 0.01$). In the more upstream Charlevoix region, where natural reproduction is documented in two rivers, all specimens but one exhibited a FW-resident phenotype, but six (out of 20) appeared to be the offspring of anadromous mothers.

Most (74%) of the rainbow trout showing either an anadromous phenotype or an anadromous maternal origin were captured in the regions bordering the downstream St. Lawrence Estuary (Bas-St-Laurent and Gaspesia). In these two regions, although we initially surmised no reproduction in these rivers and therefore expected only the anadromous phenotype, 18 rainbow trout showed a FW-resident life cycle, including two juveniles of 1- and 2-years old.

Characteristics of the anadromous phenotype

Age at habitat change

Thirteen rainbow trout, all found in Eastern Quebec (i.e., outside the stocking area), experienced at least one migration into saltwater. Characteristics of their life cycle are detailed in the Supporting Information. Briefly, most of them (62%) migrated to the marine habitat at 3 years of age. Among the seven specimens that experienced at least one complete marine phase, three first migrated upstream at 4-years old, whereas it occurred one year later for a fourth individual. Three

Table 2. Age of 51 rainbow trout found in Eastern Quebec (EQ), according to their region of capture.

Region	N	Mean	Range
Charlevoix			
<i>Du Gouffre and Malbaie R.</i>	16	5.5	4–9
<i>Other</i>	5	2.8	2–3
Bas-St-Laurent	10	3.3	1–7
Gaspesia	20	3.5	3–6

Contribution of anadromy to dispersion capacity




Fig. 3. Four otolith Sr:Ca life-history patterns observed in 83 rainbow trout (*Oncorhynchus mykiss*) captured in Eastern Quebec, as obtained by the visual interpretation method using the two-point smoothed averages (left panels), and the quantitative interpretation method using GAMs (right panels). Sr:Ca ratios between 0 and $\sim 100 \mu\text{m}$ are associated with the maternal history, whereas Sr:Ca ratios between $100 \mu\text{m}$ and the otolith edge represent the growing habitats experienced by the fish throughout its life time. Full circles represent the estimated position of each annulus. A-mother: anadromous maternal form, FW-mother: freshwater resident maternal form. The threshold of 3.5×10^{-3} , delineating between freshwater and saltwater, is shown by a dotted horizontal line.

specimens (female, male and unknown sex) were probably multispawners according to their age at change of habitat. All steelhead but two returned to freshwater during the growing season following the year of the seaward migration.

Size at migration and growth rate

Average fork length (\pm SD) at the age of the first downstream migration (excluding the youngest migrant NO1, see supporting information) was $265 \pm 67 \text{ mm}$ ($N = 12$, range: 168–363 mm). When fish first returned to freshwater, they measured $303 \pm 89 \text{ mm}$ on average ($N = 7$, range: 157–390 mm). When removing younger

specimens (NO1 and PC2) that might have been immature at that time, the mean fish length increased to $347 \pm 52 \text{ mm}$ on average ($N = 5$, range: 258–390 mm). Growth rate (length-at-age) observed for steelhead (excluding NO1) was the same as that for FW-residents [two-factor ANCOVA (age, form, age \times form): d.f. = 36, $F = 28.74$, $P > 0.05$, Fig. 5]. However, when size was back-calculated to the most common age for the first freshwater-to-sea migration, that is, at the age of 3 years, future steelhead tended to be larger than future FW-residents, but this difference was not significant ($268 \pm 61 \text{ mm}$ vs. $252 \pm 73 \text{ mm}$, Student's t -test: d.f. = 33, t -value = 0.39, $P = 0.70$). It must be

Table 3. Life-history pattern of 83 rainbow trout captured in the Quebec City sector and in three different regions of Eastern Quebec (EQ), according to their sex. For two specimens, maternal form was not possible to determine.

Region	Sex	Maternal form/Specimen's phenotype						N
		FW/FW	A/FW	FW/A	A/A	?/FW	?/A	
Quebec City	F	11	0	0	0	0	0	32
	M	4	0	0	0	0	0	
	I	17	0	0	0	0	0	
Charlevoix (EQ)	F	8	3	0	0	1	0	21
	M	5	2	1	0	0	0	
	I	0	1	0	0	0	0	
Bas-St-Laurent (EQ)	F	2	2	2	1	0	0	10
	M	0	1	0	0	0	0	
	I	2	0	0	0	0	0	
Gaspesia (EQ)	F	1	2	1	1	0	0	20
	M	5	3	1	3	0	1	
	I	0	0	0	2	0	0	
Total	–	55	14	5	7	1	1	83

FW, freshwater resident; A, anadromous; F, female; M, male; I, sex not determined (mainly triploid sterile fish).

noted however that all back-calculations were based on fish size in spring (at the time of annulus formation). Thus, estimation of size at migration does not take into account the fish's growth during summer, even if it changed habitat later during the growing season.

Discussion

Rainbow trout is slowly but surely invading Quebec streams, from the upper part of the St. Lawrence River where stocking is allowed, towards rivers located further downstream, outside the limits of the stocking area. The invasion process has led to the establishment of some self-sustaining populations in Eastern Québec, such as in two rivers of Charlevoix (Du Gouffre and Malbaie), where it forms a genetically distinct population (Thibault et al. 2009). As success of biological invasions usually depends on the presence of long-distance migrants (Begon et al. 1996), anadromy could have promoted the species' dispersion as steelhead can reach streams located far from their native river. Otolith microchemistry analysis performed on specimens captured inside and outside the stocking area revealed that although all fish in the Quebec City region were freshwater residents, both phenotypes (anadromous and freshwater resident) were found in Eastern Quebec. Moreover, the proportion of fish exhibiting the anadromous life cycle increased with the distance from the stocking zone.

Anadromy in the newly established population of Charlevoix

All rainbows captured in the Du Gouffre and Malbaie Rivers, where the species reproduces, exhibited a freshwater resident phenotype. However, 30% of them

were produced by anadromous mothers, indicating the presence of an anadromous contingent in this population. The presence of anadromy (as revealed by the maternal form) increases the probability of population dispersal. Furthermore, our results support previous genetic analyses (Thibault et al. 2009) and contradict the belief that anadromy among introduced self-sustaining populations of rainbow trout only exists in the Santa Cruz River (Pascual et al. 2001; Riva Rossi et al. 2004). Moreover, our sampling in Du Gouffre and Malbaie Rivers was not representative of the real spatial and temporal distribution of the species. Specimens were caught during and after the reproduction period, mainly in the streams, and less so at the rivers' mouths. Therefore, capture of resident fish was more probable than anadromous fish that usually stay in the river only for the spawning period. Thus, the proportion of steelhead is probably underestimated. Predominance of freshwater resident rainbows in rivers supporting reproduction could also be related to sport fishing pressure. Rainbow trout has been fished in Du Gouffre and Malbaie Rivers for many years. Steelhead, being larger than residents, are targeted by fishers, so a decrease in the relative abundance of anadromous trout could be expected, as has been observed in a native population until the adoption of a *catch-and-release* policy (Kostow 2003).

Anadromy and freshwater residency among vagrants

As predicted, anadromy (revealed by fish phenotype and maternal forms) was predominant among vagrants captured in Bas-St-Laurent and Gaspesia regions, and was even probably slightly underestimated because of the capture of a few specimens too young to have undergone smoltification. The invasion process thus appears to be supported by long-distance migrants. However, the presence of freshwater residents was unexpected in these regions, as all the fish were believed to have migrated from populations established upstream in the St. Lawrence River system. The presence of nonmigratory fish in these rivers (especially in Gaspesia) and in the Noire River (next to the Malbaie R.), suggests that either there were illegal stocking events, or that the species is reproducing in these streams. Evidence suggests that punctual stockings have no influence on the rainbow trout colonisation process (Thibault et al. 2009). Thus the spread of established populations may be greater than previously indicated.

Age and size at smoltification

Most of the sampled steelhead exhibited a typical anadromous life cycle, with a first migration towards the marine habitat at approximately 3-years old and




Fig. 4. Proportion of the anadromy and freshwater (FW)-residency forms in 83 rainbow trout (*Oncorhynchus mykiss*) captured in the Quebec City and Eastern Quebec regions, as determined from the otolith Sr:Ca patterns: (a) phenotype of analysed specimens, (b) maternal form. Anadromy is shown in black, whereas FW-residency is shown in white. Rivers where natural reproduction has been documented are labelled. The area where stocking and farming of rainbow trout is allowed is shown in dark grey. Approximate limit of the freshwater is indicated by a dashed line. 1: Quebec City region, 2: Charlevoix, 3: Bas-St-Laurent, and 4: Gaspesia.

measuring ~ 265 mm. The few younger migrants captured would constitute an exception (see below). The age at smoltification observed in Eastern Quebec is similar to that observed in juvenile steelhead within their native range where they usually spend 2 or 3 years in freshwater before migration (Burgner et al. 1992; Busby et al. 1996). However, Eastern Quebec

steelhead appear to leave freshwater at a larger size than North American west coast steelhead (smolt size ~ 160 mm, Burgner et al. 1992). A great variation was found in size at migration, suggesting that migration is more related to age than to size. This contradicts general observations for *O. mykiss* and other salmonids indicating that smolt outmigration is related




Fig. 5. Relationship between body length (fork length) and age for 37 anadromous and freshwater (FW)-resident rainbow trout (*Oncorhynchus mykiss*) captured in Eastern Quebec regions in 2005 and 2006.

to a critical size (Brisson-Bonenfant 2006 and references therein; Burgner et al. 1992). However, our results might be biased by small sample size and/or errors associated with the back-calculation method.

Purpose of habitat changes

All anadromous rainbow trout were found in rivers where no reproduction was expected. Therefore, we assumed that when they undertook their first seaward migration, they moved far away downstream of their native stream, never to return, and after a time spent in saltwater they finally entered into a new (non-native) river. The period spent in the marine habitat should not be considered only as a growing period before reproduction but also as an opportunity to move between rivers. We cannot be certain that migration into non-native rivers is solely for the purpose of reproduction. For example, a specimen captured in the Noire River (located approximately 30 km downstream of Malbaie River), that experienced precocious down- and upstream migrations, completed its marine phase before the age of two years (see Supporting Information). It appears unlikely that this fish was mature enough to reproduce. We rather suggest that it might have quickly migrated out of its native river (probably the Malbaie River), simply to switch rivers without necessarily spawning.

Age at reproduction

For almost all anadromous fish sampled in this study, the first return to freshwater occurred at 4 years of age. Despite the fact that the first upstream migration might not always be for the purpose of reproduction, it seems reasonable to affirm that the age at reproduction for most steelhead in Eastern Quebec is similar to steelhead within their native range, spawning at approximately 4-years old (range: 3–5) (Busby et al.

1996; Kostow 2003; Seamons et al. 2007). However, it should be noted that this conclusion is based on only a few individuals ($N = 5$) that had time to enter into the river for spawning before being captured.

Multispawning of steelhead

If repeat spawning is frequent among freshwater residents, then steelhead multispawners are believed to be generally few in populations of the temperate zone of the North American west coast, with an incidence usually (but not exclusively) less than 10% (Burgner et al. 1992; Busby et al. 1996; Kostow 2003). In our study, three of the 13 fish exhibiting the anadromous life cycle were probably multispawners. Despite our small sample size, we think that the prevalence of multispawners in Eastern Quebec is probably higher than what has been reported in almost all coastal native populations (Narum et al. 2008a), but certainly less than what was observed in the naturalised population of the Santa Cruz River (60%, Pascual et al. 2001). Multispawning can increase the success of invaders as this reproductive mode accelerates effective population growth and ensures genetic variability (Riva-Rossi et al. 2007; Narum et al. 2008a). This impact is enhanced when multispawners are anadromous females, which are known to be much more fecund than resident females (Kostow 2003).

Determination of the phenotype

Although we are unable to identify the life cycle of the fathers of our sampled fish, our results show that the maternal form is not necessarily expressed by its progeny. Within our sample in Eastern Quebec, we found resident rainbow trout with an anadromous maternal origin (38%) and steelhead with a resident maternal origin (42%). This is contrary to what was found by Zimmerman & Reeves (2000) and Donohoe et al. (2008), using a similar approach. However, direct spawning observations, pedigree analyses and breeding experiments have previously demonstrated that both resident and anadromous parents can produce either freshwater resident or steelhead offspring (see references in Kostow 2003). Supporting these observations, population structure analyses conducted on sympatric resident and anadromous forms have shown a lack of strong genetic divergence between life-history forms (Pascual et al. 2001; Docker & Heath 2003; Narum et al. 2004; Riva Rossi et al. 2004; Heath et al. 2008). Many environmental factors associated with a particular geographic area, such as habitat productivity, migratory hazards, habitat connectivity or freshwater mortality, have been proposed to explain the occurrence of the two life history strategies (Kostow 2003 and

references therein; Narum et al. 2008b). However, as we found both forms in the same streams, it appears that the adoption of residency or anadromy is the expression of conditional, alternative developmental pathways under the control of threshold traits (Brockman & Taborsky 2008).

We found no difference in growth rate (length-at-age relationship) and size at the age of migration between anadromous and freshwater resident specimens. Based on previous studies on salmonids, we expected that future steelhead would be, just before migration, smaller than future residents (for a same age) (Thériault & Dodson 2003 and references therein), but would thereafter experience a faster growth than freshwater residents (MacCrimmon & Gots 1972; Thorpe 1994; Pascual et al. 2001). The absence of significant differences between the two life-history forms may have been related to purely methodological problems: (i) the number of anadromous specimens was low, (ii) sizes provided by fishers may have been incorrectly measured or (iii) annuli on otolith's pictures may have been wrongly positioned, or estimated distances between the otolith core and annuli may have been erroneous due to picture precision and unequal otolith grinding. Excluding growth, size-at-age and maternal form, other individual traits, such as sex or condition factor, could have played a role in the adoption of the phenotype. In this study, we found no relation between sex and a particular life history strategy, as the sex ratio was of 1:1 among anadromous fish. As the mass was not available for many specimen, it was not possible to investigate a potential role of condition factor in the adoption of a phenotype.

Sampling bias and implications for extrapolation

As the invasion process is still in progress and the number of rainbow trout in many rivers is low, our sampling effort was not systematic but was rather based on voluntary angler returns. As such, some caution is required in applying our conclusions to the situation as a whole. Nevertheless, the presence of consistent patterns within the different river categories (only freshwater residents in all rivers located upstream inside the stocking area (SWQ and QC), dominance of freshwater residents in established populations and dominance of anadromy in rivers of the Gaspesia and Bas-St-Laurent regions) indicates that our conclusions would probably apply in the case of increased sample sizes and locations. Nevertheless, future studies could focus the sampling efforts on a smaller number of rivers harbouring an abundance of rainbow trout and located at various distances from the established populations and stocking area to validate these preliminary conclusions.

Concluding remarks

As expected, otolith microchemistry analyses demonstrated that the invasion process by rainbow trout in Eastern Quebec is promoted by the presence of anadromous individuals that are able to colonise new rivers following long-distance migrations along the St. Lawrence Estuary corridor. In the newly established population of Charlevoix, anadromy was revealed by the maternal signature, indicating that the Santa Cruz River (Patagonia) is not the only introduced rainbow trout population that has developed an anadromous run. Surprisingly, some specimens with a freshwater resident phenotype were found among putative vagrants, suggesting the existence of unknown self-sustaining populations in Bas-St-Laurent and/or Gaspesia regions. These findings reveal the need for a more rigorous monitoring of the rivers in these regions, in order to monitor and mitigate the establishment of new rainbow trout populations in rivers supporting indigeneous salmonids such as Atlantic salmon (*Salmo salar*) and brook charr (*Salvelinus fontinalis*).

Acknowledgements

This work was funded by grants from the Natural Sciences and Engineering Research Council of Canada, Centre Interuniversitaire de Recherche sur le Saumon Atlantique (CIRSA), the Fédération Québécoise de la Faune and the Réseau Aquaculture de Québec. We thank our collaborators that helped to collect rainbow trout samples: Ministère des Ressources naturelles et de la faune, Québec Pêche, Fédération québécoise pour le saumon Atlantique, Fédération des gestionnaires de rivières à saumon du Québec, Simon Blanchet, and all ZECs, organisations, sport and commercial fishermen that participated in the sampling campaigns. A special thank to Véronique Thériault and Simon Blanchet for their implication in age determination. We are greatfull to Christopher Donohoe for his advice on the maternal Sr:Ca signature interpretation, and to Rémy Pouliot for his comments on earlier versions of the manuscript.

References

- Begon, M., Harper, J.L. & Townsend, C.R. 1996. Ecology. Individuals populations and communities, 3rd edn. Blackwell Science, Cambridge, p. 1068.
- Brisson-Bonenfant, C. 2006. Stratégie conditionnelle et migration chez le saumon atlantique (*Salmo salar*) juvénile. M.Sc. Thesis. Faculté des Sciences et de Génie, Département de biologie, Université Laval, Québec, 94 pp.
- Brockman, H.J. & Taborsky, M. 2008. Alternative reproductive tactics and the evolution of alternative allocation phenotypes. In: Oliveira, R.F., Taborsky, M. & Brockman, H.J., eds. Alternative Reproductive tactics: an integrative approach. Cambridge University Press, Cambridge, pp. 25–51.
- Burgner, R.L., Light, J.T., Margolis, L., Okazaki, T., Tautz, A. & Ito, S. 1992. Distribution and origins of steelhead trout (*Oncorhynchus mykiss*) in offshore waters of the North

- Pacific ocean. International North Pacific Fisheries Commission, Bulletin 51: 73.
- Busby, P.J., Wainwright, T.C., Bryant, G.J., Lierheimer, L.J., Waples, R.S., Waknitz, F.W. & Lagomarsino, I.V. 1996. Status review of West Coast steelhead from Washington, Idaho, Oregon and California. NOAA Tech. Memo., NMFS-NWFSC-27. US Dept. of Commerce, 261 pp.
- Colautti, R.I., Grigorovich, I.A. & MacIsaac, H.J. 2006. Propagule pressure: a null model for biological invasions. *Biological Invasions* 8: 1023–1037.
- Daverat, F., Limburg, K.E., Thibault, I., Shiao, J.-C., Dodson, J.J., Caron, F., Tzeng, W.-N., Iizuka, Y. & Wickström, H. 2006. Phenotypic plasticity of habitat use by three temperate eel species *Anguilla anguilla*, *A. japonica* and *A. rostrata*. *Marine Ecology Progress Series* 308: 231–241.
- Docker, M.F. & Heath, D.D. 2003. Genetic comparison between sympatric anadromous steelhead and freshwater resident rainbow trout in British Columbia, Canada. *Conservation Genetics* 4: 227–231.
- Donohoe, C.J., Adams, P.B. & Royer, C.F. 2008. Influence of water chemistry and migratory distance on ability to distinguish progeny of sympatric resident and anadromous rainbow trout (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences* 65: 1160–1175.
- DuBois, R.B., Plaster, S.D. & Rasmussen, P.W. 1989. Fecundity of spring- and fall-run steelhead from two western Lake Superior tributaries. *Transactions of the American Fisheries Society* 118: 311–316.
- Francis, R.I.C.C. 1990. Back-calculation of fish length: a critical review. *Journal of Fish Biology* 36: 883–902.
- Gillanders, B.M. 2005. Otolith chemistry to determine movements of diadromous and freshwater fish. *Aquatic Living Resources* 18: 291–300.
- GISD. 2005. Global Invasive Species Database. *Oncorhynchus mykiss*. Available from: <http://www.issg.org/database/species/ecology.asp?si=103&fr=1&sts=sss&lang=EN> (Accessed 26th August 2009).
- Gunn, J.S., Harrowfield, I.R., Proctor, C.H. & Thresher, R.E. 1992. Electron microprobe analysis of fish otoliths - evaluation of techniques for studying age and stock discrimination. *Journal of Experimental Marine Biology and Ecology* 158: 1–36.
- Heath, D.D., Bettles, C.M., Jamieson, S., Stasiak, I. & Docker, M.F. 2008. Genetic differentiation among sympatric migratory and resident life history forms of rainbow trout in British Columbia. *Transactions of the American Fisheries Society* 137: 1268–1277.
- Hedger, R.D., Atkinson, P.M., Thibault, I. & Dodson, J.J. 2008. A quantitative approach for classifying fish otolith strontium:calcium sequences into environmental histories. *Ecological Informatics* 3: 207–217.
- Jessop, B.M., Shiao, J.-C., Iizuka, Y. & Tzeng, W.N. 2002. Migratory behaviour and habitat use by American eels *Anguilla rostrata* as revealed by otolith microchemistry. *Marine Ecology Progress Series* 233: 217–229.
- Jessop, B.M., Shiao, J.C., Iizuka, Y. & Tzeng, W.N. 2004. Variation in the annual growth, by sex and migration history, of silver American eels *Anguilla rostrata*. *Marine Ecology Progress Series* 272: 231–244.
- Jessop, B.M., Shiao, J.C., Iizuka, Y. & Tzeng, W.N. 2008. Prevalence and intensity of occurrence of vaterite inclusions in aragonite otoliths of American eels *Anguilla rostrata*. *Aquatic Biology* 2: 171–178.
- Kolar, C.S. & Lodge, D.M. 2001. Progress in invasion biology: predicting invaders. *Trends in Ecology and Evolution* 16: 199–204.
- Kostow, K. 2003. Factors that influence evolutionarily significant unit boundaries and status assessment in a highly polymorphic species, *Oncorhynchus mykiss*, in the Columbia Basin. Information Report #2003-04. Oregon Department of Fish and Wildlife and NOAA Fisheries, 122 pp.
- Lecomte, F. & Dodson, J.J. 2004. Role of early life-history constraints and resource polymorphism in the segregation of sympatric populations of an estuarine fish. *Evolutionary Ecology Research* 6: 631–658.
- MacCrimmon, H.R. 1971. World distribution of rainbow trout (*Salmo gairdneri*). *Journal of Fisheries and Research Board of Canada* 28: 663–704.
- MacCrimmon, H.R. & Gots, B.L. 1972. Rainbow trout in the Great Lakes. Ministry of Natural Resources, Ontario, 66 pp.
- Moyle, P.B. & Marchetti, M.P. 2006. Predicting invasion success: freshwater fishes in California as a model. *BioScience* 56: 515–524.
- Narum, S.R., Contor, C., Talbot, A. & Powell, M.S. 2004. Genetic divergence of sympatric resident and anadromous forms of *Oncorhynchus mykiss* in the Walla Walla River, U.S.A. *Journal of Fish Biology* 65: 471–488.
- Narum, S.R., Hatch, D., Talbot, A.J., Moran, P. & Powell, M.S. 2008a. Iteroparity in complex mating systems of steelhead *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Biology* 72: 45–60.
- Narum, S.R., Zendt, J.S., Graves, D. & Sharp, W.R. 2008b. Influence of landscape on resident and anadromous life history types of *Oncorhynchus mykiss*. *Canadian Journal of Fisheries and Aquatic Sciences* 65: 1013–1023.
- Pascual, M.A., Bentzen, P., Riva-Rossi, C.M., Mackey, G., Kinnison, M.T. & Walker, R. 2001. First documented case of anadromy in a population of introduced rainbow trout in Patagonia, Argentina. *Transactions of the American Fisheries Society* 130: 53–67.
- Reed, S.J.B. 1993. *Electron microprobe analysis*, 2nd edn. Cambridge University Press, Cambridge, 326 pp.
- Rieman, B.E., Myers, D.L. & Nielsen, R.L. 1994. Use of otolith microchemistry to discriminate *Oncorhynchus nerka* of resident and anadromous origin. *Canadian Journal of Fisheries and Aquatic Sciences* 51: 68–77.
- Riva Rossi, C.M., Lessa, E.P. & Pascual, M.A. 2004. The origin of introduced rainbow trout (*Oncorhynchus mykiss*) in the Santa Cruz River, Patagonia, Argentina, as inferred from mitochondrial DNA. *Canadian Journal of Fisheries and Aquatic Sciences* 61: 1095–1101.
- Riva-Rossi, C., Pascual, M.A., Babaluk, J.A., García-Asorey, M. & Halden, N.M. 2007. Intra-population variation in anadromy and reproductive life span in rainbow trout introduced in the Santa Cruz River, Argentina. *Journal of Fish Biology* 70: 1780–1797.
- Seamons, T.R., Bentzen, P. & Quinn, T.P. 2007. DNA parentage analysis reveals inter-annual variation in selection: results from 19 consecutive brood years in steelhead trout. *Evolutionary Ecology Research* 9: 409–431.
- Theoharides, K.A. & Dukes, J.S. 2007. Plant invasion across space and time: factors affecting nonindigenous species

- success during four stages of invasion. *New Phytologist* 176: 256–273.
- Thériault, V. & Dodson, J.J. 2003. Body size and the adoption of a migratory tactic in brook charr. *Journal of Fish Biology* 63: 1144–1159.
- Thibault, I., Bernatchez, L. & Dodson, J.J. 2009. The contribution of newly-established populations to the dynamics of range expansion in a one-dimensional fluvial-estuarine system: the case of rainbow trout (*Oncorhynchus mykiss*) in Eastern Quebec. *Diversity and Distributions* 15: 1060–1072.
- Thorpe, J.E. 1994. Salmonid fishes and the estuarine environment. *Estuaries* 17: 76–93.
- Tzeng, W.-N., Wang, C.H., Wickström, H. & Reizenstein, M. 2000. Occurrence of the semi-catadromous European eel *Anguilla anguilla* in the Baltic Sea. *Marine Biology* 137: 93–98.
- Tzeng, W.-N., Iizuka, Y., Shiao, J.C., Yamada, Y. & Oka, H.P. 2003. Identification and growth rates comparison of divergent migratory contingents of Japanese eel (*Anguilla anguilla*). *Aquaculture* 216: 77–86.
- Tzeng, W.-N., Chang, C.-W., Wang, C.-H., Shiao, J.-C., Iizuka, Y., Yang, Y.-J., You, C.-F. & Lozys, L. 2007. Misidentification of the migratory history of anguillid eels by Sr/Ca ratios of vaterite otoliths. *Marine and Freshwater Research* 348: 285–295.
- Volk, E.C., Blakley, A., Schroder, S.L. & Kuehner, S.M. 2000. Otolith chemistry reflects migratory characteristics of Pacific salmonids: Using otolith core chemistry to distinguish maternal associations with sea and freshwaters. *Fisheries Research* 46: 251–266.
- Wood, S.N. 2006. Generalized additive models: an introduction with R. Chapman and Hall/CRC, Boca Raton, FL, 416 pp.
- Zimmerman, C.E. 2005. Relationship of otolith strontium-to-calcium ratios and salinity: experimental validation for juvenile salmonids. *Canadian Journal of Fisheries and Aquatic Sciences* 62: 88–97.
- Zimmerman, C.E. & Reeves, G.H. 2000. Population structure of sympatric anadromous and nonanadromous *Oncorhynchus mykiss*: evidence from spawning surveys and otolith microchemistry. *Canadian Journal of Fisheries and Aquatic Sciences* 57: 2152–2162.
- Zimmerman, C.E. & Reeves, G.H. 2002. Identification of steelhead and resident rainbow trout progeny in the Deschutes River, Oregon, revealed with otolith microchemistry. *Transactions of the American Fisheries Society* 131: 986–993.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Appendix A. Age at migration for 13 anadromous rainbow trout captured in 2006 in Eastern Quebec regions, as revealed by otolith microchemistry. “+” refers to fish age during the growing season (late spring to early fall), “++” refers to fish age at fall, absence of “+” refers to fish age at early spring

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.