

Multi-component decaying behavior on high-b-value diffusion-weighted MRI

Wen-Chau Wu¹, Kuo-Hsiang Kao¹, Ping-Hong Lai² and Hsiao-Wen Chung¹

¹Department of Electrical Engineering, National Taiwan University, Taipei, Taiwan, R.O.C.

²Department of Radiology, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan, R.O.C.

Abstract-This study reported the multi-component decaying behavior on high-b-value diffusion-weighted MR images. Signal intensity was measured in different regions of the brain. Results showed that gray matter and cerebrospinal fluid exhibited single exponential decaying characteristics while multi-component decaying behavior was observed in some white matter areas with complicated neural fiber orientations.

I. INTRODUCTION

Diffusion-weighted MR imaging (DWI), in particular mapping of the diffusion tensors, has been proven useful in depicting neural fiber orientations by making use of the anisotropy of structure-restricted diffusion [1], [2]. Due to scan time consideration, mapping of the diffusion tensors is often performed with only two b values, on the order of 0 and 1000 sec/mm² in the human brain. On the other hand, during the development of many neurological disorders in human brain, alterations in the apparent diffusion coefficient (ADC) of water have been detected [3], [4]. So far, most studies performed with conventional MRI hardware report diffusion measurement in terms of a single ADC. In other words, the signal decay caused by water diffusion is assumed to be mono-exponential. With the advent of advances in hardware technology, high-b-value (>1000 sec/mm²) DWI is made feasible in clinical MR systems and it has been shown that the diffusion signal decay is no longer monoexponential under high b values [5].

In this study we report the investigation of the multi-exponential behavior observed on DWI with b values up to 3000 sec/mm². We provide an explanation for this behavior and address the cautions in its applications.

II. MATERIALS AND METHODS

A. Image Acquisition

Transaxial echo-planar diffusion-weighted images of the brain were acquired from healthy volunteers using a 1.5T system (General Electric Signa Horizon, Milwaukee, Wisconsin) with multiple b values of 0, 500, 1000, 1500, 2000, 2500, and 3000 sec/mm² (TR/TE=4700/118, in-plane resolution 1.2x2.5 mm²). Diffusion-weighted gradients were applied along three orthogonal directions (superio-inferior, antero-posterior, and right-left) for all b values. Signal intensity measured in different regions of the brain was fitted by both monoexponential and biexponential models as described in the next paragraph. Then, the signal intensity was plotted in logarithmic scale versus the b values. Any observed nonlinear behavior was recorded, and the ADC obtained with b=0 and 1000 was

compared to that obtained with b=0 and 2000.

B. Mathematical Models

1. Monoexponential:

$$S = S_0 \exp(-bD)$$

where D is the ADC, S_0 and S are the signal intensity without and with diffusion weight respectively. This model assumes monoexponential decaying behavior with the increase of b values.

2. Biexponential: Since the fiber orientation in brain tissue is complicated, the overall diffusion behavior of a region can be regarded as the contribution of several compartments determined by the fiber tracts. In biexponential model, we divided water diffusion into two components.

$$S = S_0 [f_1 \exp(-bD_1) + f_2 \exp(-bD_2)]$$

$$f_1 + f_2 = 1$$

where f_1 is the fraction of primary component, i.e. the fraction of principle orientation of fibers enclosed, D_1 is the primary ADC, f_2 is the fraction of residual component and D_2 is the mean ADC contributed by other minor diffusion compartments.

C. SNR simulation

To assess the influence of noise, simulated DWIs, superimposed with Gaussian noise, were generated with signal-to-noise ratio (SNR) 6, 4.5 and 2. Referring to the literature ADC range of human brain tissues, D was assigned 0.001 mm²/sec in monoexponential model while D_1 and D_2 were given 0.001 mm²/sec and 0.0001 mm²/sec respectively in biexponential model.

D. Curve fitting

Least-squares algorithm was adopted to find D, D_1 , D_2 , f_1 and f_2 . R^2 was calculated, where R is the multiple correlation coefficient. In different regions of the brain, the fitted ADCs were compared with the literature values. There must be discrepancies because more than one diffusion compartments were considered here. However, it is rational to use the literature CSF ADC (2.94+/-0.05 ($\times 10^{-3}$ mm²/sec)) as the upper limit of the ADCs obtained in other brain tissues. Moreover, we excluded any fitted ADC exceeding 1.5 times the literature value of the tissue.

III. RESULTS

Fig.1 and Fig.2 show the result of simulation. In both mono- and bi- exponential models, noise plays an

important role as b value exceeds 2500. In addition, when SNR drops to 2 (2.5 actually) or less, the simulated result begins to deviate from the ideal condition and this may lead to misinterpretation of mono- or multi-exponential decay.

The anatomical regions of interest are shown in Fig.3. Fig.4 shows the signal intensities of the diffusion-weighted images in cortex gray matter, corpus callosum and optical radiation, plotted in logarithmic scale versus the b values. The cortex gray matter and cerebrospinal fluid (not shown here) exhibited single exponential decaying characteristics. In the white matter, however, variant behaviors were found in different regions. It is seen that the corpus callosum and optical radiation, where there exist complicated neural fiber orientations within the chosen regions of interest, demonstrated non-linear multi-exponential decaying compartments along some directions. In the corpus callosum for example, ADC calculated from $b=0$ and 1000 (along left-right direction) was higher than that obtained with $b=0$ and 2000 by a factor of 2.




Fig. 1. Assessment of noise interference in curve-fitting in mono-exponential model ($D = 0.001 \text{ mm}^2/\text{sec}$). Simulated DWIs superimposed by Gaussian noise were generated with SNR 6 (upper), 4.5 (middle) and 2 (lower) respectively. The signal intensity was plotted in logarithmic scale versus the b values. Solid line is the ideal mono-decay and dotted line is

obtained from simulation.




Fig. 2. Assessment of noise interference in curve-fitting in biexponential model ($D_1 = 0.001 \text{ mm}^2/\text{sec}$, $D_2 = 0.0001 \text{ mm}^2/\text{sec}$). Simulated DWIs superimposed by Gaussian noise were generated with SNR 6 (upper), 4.5 (middle) and 2 (lower) respectively. The signal intensity was plotted in logarithmic scale versus the b values. Solid line is the ideal biexponential decay and dotted line is obtained from simulation.




Fig. 3. The anatomical regions of interest: (a) cortex gray matter, (b) corpus callosum and (c) optical radiation.




Fig. 4. The DWI signal intensities in cortex gray matter (upper), corpus callosum (middle) and optical radiation (lower) are plotted in logarithmic scale versus their corresponding b values. Diffusion-weighted gradients were applied along three orthogonal directions: antero-posterior (AP), right-left (RL) and superio-inferior (SI).

IV. DISCUSSIONS AND CONCLUSIONS

Diffusion is known to be anisotropic, with the preferential direction of movement along the white matter track (i.e., highest ADC). We thus attribute the multi-exponential behavior as arising from an inclusion of neural fibers along different orientations in a single region of interest. In cortex gray matter consisting of cell bodies, no noticeable multiple diffusion compartments (fiber tracts) exist. Hence, diffusion-induced signal decay is contributed by only one component, i.e. monoexponential or linear in logarithmic scale. In areas such as the corpus callosum where the fiber directions are complicated, DWI with low b values presumably reflects the fastest decaying component (i.e., fibers along the direction of diffusion-sensitizing gradients). Therefore, the existence of slow ADC compartments can only be visible on DWI with high b

values. The findings from this study further suggest that the interpretation of white matter track should be taken with care if mapping of the diffusion tensor were accomplished with only low b values.

REFERENCE

- [1] P. J. Basser, S. Pajevic, C. Pierpaoli, J. Duda and A. Aldroubi, "In vivo fiber tractography using DT-MRI data," *Magn. Reson. Med.*, vol. 44, pp. 625-632, 2000.
- [2] D. K. Jones, A. Simmons, S. C. Williams and M. A. Horsfield, "Non-invasive assessment of axonal fiber connectivity in the human brain via diffusion tensor MRI," *Magn. Reson. Med.*, vol. 42, pp. 37-41, 1999.
- [3] S. Warach, D. Chien, W. Li, M. Ronthal and R. R. Edelman, "Fast magnetic resonance diffusion-weighted imaging of acute human stroke," *Neurology*, vol. 42, pp. 1717-1723, 1992.
- [4] H. B. W. Larsson, C. Thomsen, J. Frederiksen, M. Stubgaard and O. Henriksen, "In vivo magnetic resonance diffusion measurement in the brain of patients with multiple sclerosis," *Magn. Reson. Imaging*, vol. 31, pp. 7-12, 1992.
- [5] M. C. DeLano, T. G. Cooper, J. E. Siebert, M. J. Potchen and K. Kuppusamy, "High-b-value diffusion-weighted MR imaging of adult brain: image contrast and apparent diffusion coefficient map features," *AJNR Am. J. Neuroradiol.*, vol. 21, pp. 1830-1836, 2000.