

Synthesis of Organoplatinum Oligomers by Employing N-Donor Bridges with Predesigned Geometry: Structural and Photophysical Properties of Luminescent Cyclometalated Platinum(II) Macrocycles

Siu-Wai Lai, Michael C. W. Chan, Kung-Kai Cheung, Shie-Ming Peng,[†] and Chi-Ming Che^{*}

Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong

Received May 10, 1999

A series of luminescent di- and trimeric cyclometalated platinum(II) macrocycles, namely, [Pt(C–N)(N–N)]₂ (N–N = pyrazolate, pz (**2**); 7-azaindolate, aza (**3**); C–N = 2-(2'-thienyl)pyridyl, Thpy (**a**); 7,8-benzoquinolate, Bzqn (**b**); 2-phenylpyridyl, Phpy (**c**)) and [Pt(C–N)(N–N)]₃ (N–N = benzimidazolate, bzim (**4**); C–N = Thpy (**a**) and Bzqn (**b**)), are synthesized in high yields (70–90%). The rigid, predefined coordination geometry of the pz, aza, and bzim bridging ligands ensure the efficient and selective assembly of the respective cyclic oligomers. The stacking arrangements in the crystal structures of **1a** and **4a** reveal π – π interactions between 2-(2'-thienyl)pyridyl moieties. The lowest energy absorption band in the UV–vis spectra is assigned to metal-to-ligand charge transfer (MLCT) transitions. The 298 K fluid emissions of complexes **2–4a** bearing Thpy ligands are assigned to transitions with mixed MLCT and IL (intraligand) character. In crystalline form, the observed red-shift is attributed to excimeric emission arising from π -stacking in the solid state. In contrast, a number of Bzqn derivatives are nonemissive in CH₂Cl₂ at 298 K. The energy of the lowest energy UV–vis and emissive bands for Thpy oligomers are red-shifted relative to the Bzqn and Phpy congeners. Significantly, Thpy complexes and especially the μ -pyrazolate species **2a** ($\tau_0 = 15.5 \mu\text{s}$, $\phi_0 = 0.18$) emit with longer lifetimes and higher quantum yields.

Introduction

Multidentate nitrogen-donor ligands are extensively utilized for the assembly of cyclic supermolecules. Pyrazolyl,^{1,2} imidazolyl,³ and particularly pyridyl^{4–7} derivatives have been incorporated into the design and construction of novel oligometallic macrocycles and cage

complexes. Nevertheless, integration of photoluminescent moieties into these potential molecular receptors is rarely undertaken, although this can confer chemical-sensing applications.⁸ In this context, cyclometalated platinum(II) complexes bearing substituted pyridyl and 2,2'-bipyridyl ligands are promising candidates because they display emissive metal-to-ligand charge transfer (MLCT) excited states in solution,^{9–14} and these emissions are extremely environment-sensitive.^{15,16}

In this work, our intention was to fabricate platinum-containing macrocycles by connecting cyclometalated Pt-

^{*} Corresponding author. Fax: +852 2857 1586. E-mail: cmche@hkucc.hku.hk.

[†] Department of Chemistry, National Taiwan University, Taipei, Taiwan.

(1) Reviews: (a) Trofimenko, S. *Prog. Inorg. Chem.* **1986**, *34*, 115. (b) La Monica, G.; Ardzioia, G. A. *Prog. Inorg. Chem.* **1997**, *46*, 151.

(2) Pt examples: (a) Minghetti, G.; Banditelli, G.; Bonati, F. *J. Chem. Soc., Dalton Trans.* **1979**, 1851. (b) Goel, A. B.; Goel, S.; Vanderveer, D. *Inorg. Chim. Acta* **1984**, *82*, L9. (c) Jain, V. K.; Kannan, S.; Tiekink, E. R. T. *J. Chem. Soc., Dalton Trans.* **1993**, 3625. (d) Stobart, S. R.; Dixon, K. R.; Eadie, D. T.; Atwood, J. L.; Zaworotko, M. D. *Angew. Chem., Int. Ed. Engl.* **1980**, *19*, 931. Other metals: (e) Chong, K. S.; Rettig, S. J.; Storr, A.; Trotter, J. *Can. J. Chem.* **1979**, *57*, 3090. (f) Murray, H. H.; Raptis, R. G.; Fackler, J. P., Jr. *Inorg. Chem.* **1988**, *27*, 26. (g) Pinillos, M. T.; Tejel, C.; Oro, L. A.; Aprada, M. C.; Foces-Foces, C.; Cano, F. H. *J. Chem. Soc., Dalton Trans.* **1989**, 1133. (h) Jeffery, J. C.; Jones, P. L.; Mann, K. L. V.; Psillakis, E.; McCleverty, J. A.; Ward, M. D.; White, C. M. *Chem. Commun.* **1997**, 175.

(3) (a) Tiripicchio, A.; Tiripicchio Camellini, M.; Usón, R.; Oro, L. A.; Ciriano, M. A.; Pinillos, M. T. *J. Organomet. Chem.* **1982**, *224*, 207. (b) Chaudhuri, P.; Karpenstein, I.; Winter, M.; Butzlaff, C.; Bill, E.; Trautwein, A. X.; Flörke, U.; Haupt, H.-J. *J. Chem. Soc., Chem. Commun.* **1992**, 321. (c) Rüttimann, S.; Bernardinelli, G.; Williams, A. F. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 392. (d) Matsumoto, N.; Motoda, Y.; Matsuo, T.; Nakashima, T.; Re, N.; Dahan, F.; Tuchagues, J.-P. *Inorg. Chem.* **1999**, *38*, 1165. Related nucleobases: (e) Smith, D. P.; Baralt, E.; Morales, B.; Olmstead, M. M.; Maestre, M. F.; Fish, R. H. *J. Am. Chem. Soc.* **1992**, *114*, 10647. (f) Chen, H.; Olmstead, M. M.; Smith, D. P.; Maestre, M. F.; Fish, R. H. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 1514.

(4) (a) Fujita, M.; Oguro, D.; Miyazawa, M.; Oka, H.; Yamaguchi, K.; Ogura, K. *Nature* **1995**, *378*, 469. (b) Fujita, M. *Chem. Soc. Rev.* **1998**, *27*, 417.

(5) Stang, P. J.; Olenyuk, B. *Acc. Chem. Res.* **1997**, *30*, 502. (6) Slone, R. V.; Benkstein, K. D.; Bélanger, S.; Hupp, J. T.; Guzei, I. A.; Rheingold, A. L. *Coord. Chem. Rev.* **1998**, *171*, 221.

(7) (a) Leung, W. H.; Cheng, J. Y. K.; Hun, T. S. M.; Che, C. M.; Wong, W. T.; Cheung, K. K. *Organometallics* **1996**, *15*, 1497. (b) Schnebeck, R.-D.; Randaccio, L.; Zangrando, E.; Lippert, B. *Angew. Chem., Int. Ed.* **1998**, *37*, 119. (c) Hall, J. R.; Loeb, S. L.; Shimizu, G. K. H.; Yap, G. P. A. *Angew. Chem., Int. Ed.* **1998**, *37*, 121. (d) Schneider, R.; Hosseini, M. W.; Planeix, J.-M.; De Cian, A.; Fischer, J. *Chem. Commun.* **1998**, 1625.

(8) Slone, R. V.; Yoon, D. I.; Calhoun, R. M.; Hupp, J. T. *J. Am. Chem. Soc.* **1995**, *117*, 11813.

(9) Chassot, L.; Müller, E.; von Zelewsky, A. *Inorg. Chem.* **1984**, *23*, 4249.

(10) Sandrini, D.; Maestri, M.; Balzani, V.; Chassot, L.; von Zelewsky, A. *J. Am. Chem. Soc.* **1987**, *109*, 7720.

(11) (a) Chan, C. W.; Lai, T. F.; Che, C. M.; Peng, S. M. *J. Am. Chem. Soc.* **1993**, *115*, 11245. (b) Chan, C. W.; Cheng, L. K.; Che, C. M. *Coord. Chem. Rev.* **1994**, *132*, 87.

(12) (a) Cheung, T. C.; Cheung, K. K.; Peng, S. M.; Che, C. M. *J. Chem. Soc., Dalton Trans.* **1996**, 1645. (b) Lai, S. W.; Chan, M. C. W.; Cheung, T. C.; Peng, S. M.; Che, C. M. *Inorg. Chem.*, in press.

(II) luminophores with various bidentate N-donor linkers. The pz (Hpz = pyrazole), aza (Haza = 7-azaindole), and bzim (Hbzim = benzimidazole) ligands exhibit predefined yet different geometrical characteristics. Hence accurate design is anticipated to yield specific molecular architectures that are thermodynamically favored. A ligand-directed strategy for the synthesis of triplatinum macrocycles by employing bzim as a rigid nonlinear bridging motif has been communicated.¹⁷ We now describe the preparation, crystal structures, and photophysics of luminescent platinum(II) cyclic dimers (bearing pz and aza) and trimers (bearing bzim) plus their monomeric precursors. By varying the N-donor bridges and the cyclometalating ligands, namely, Thpy (HThpy = 2-(2'-thienyl)pyridine), Bzqn (HBzqn = 7,8-benzoquinoline), and Phpy (HPhpy = 2-phenylpyridine), we are able to assess trends and changes in emissive properties. Seminal spectroscopic studies by Gray and co-workers on d⁸-d⁸ binuclear species containing μ -pyrazolate ligands have appeared.^{18,19}

Experimental Section

General Procedures. The following chemicals were obtained from Aldrich Chemical Co. and used as received: K₂PtCl₄ (98%), 2-phenylpyridine (HPhpy), 7,8-benzoquinoline (HBzqn), sodium hydride, pyrazole (Hpy), benzimidazole (Hbzim), and 7-azaindole (Haza). 2-(2'-Thienyl)pyridine (HThpy) was purchased from Lancaster Chemical Co. [¹⁸⁵Pt(Phpy)Cl₂] (**1c**) was prepared by literature methods.²⁰ Synthesis of **4a,b** has been described previously.¹⁷ Dichloromethane for photophysics studies was washed with concentrated sulfuric acid, 10% sodium hydrogen carbonate, and water, dried by calcium chloride, and distilled over calcium hydride. All other solvents were purified according to conventional methods.²¹

Fast atom bombardment (FAB) mass spectra were obtained on a Finnigan Mat 95 mass spectrometer. ¹H (300 MHz) and ¹³C (126 MHz) spectra were recorded on DPX 300 and 500 Bruker FT-NMR spectrometers, respectively, with chemical shifts (in ppm) relative to tetramethylsilane. Elemental analysis was performed by the Institute of Chemistry at the Chinese Academy of Sciences, Beijing. Infrared spectra were recorded as KBr plates on a BIO RAD FT-IR spectrometer. UV-vis spectra were recorded on a Perkin-Elmer Lambda 19 UV/vis spectrophotometer.

Emission and Lifetime Measurements. Steady-state emission spectra were recorded on a SPEX 1681 Fluorolog-2 series F111AI spectrophotometer. Low-temperature (77 K) emission spectra for frozen and solid-state samples were recorded in 5 mm diameter quartz tubes, which were placed

in a liquid nitrogen Dewar equipped with quartz windows. Emission spectra were corrected for monochromator and photomultiplier efficiency and for xenon lamp stability. Details of emission quantum yield determinations by the method of Demas and Crosby²² have been given.¹¹ Emission lifetime measurements were performed with a Quanta Ray DCR-3 pulsed Nd:YAG laser system (pulse output 355 nm, 8 ns). The emission signals were detected by a Hamamatsu R928 photomultiplier tube and recorded on a Tektronix model 2430 digital oscilloscope. Errors for λ values (± 1 nm), τ ($\pm 10\%$), and ϕ ($\pm 10\%$) are estimated.

Synthesis. [Pt(Thpy)(HThpy)Cl]₂, 1a. [¹⁸⁵PtCl₄] was prepared in the organic phase of a phase-transfer metathesis reaction by mixing K₂PtCl₄ (0.20 g, 0.48 mmol) in H₂O (10 mL) with ¹⁸⁵PtCl₄ (0.58 g, 2.09 mmol) in CH₂Cl₂ (30 mL).²⁰ HThpy (0.32 g, 1.99 mmol) in CH₃OH (30 mL) was then added to the CH₂Cl₂ phase. The red solution was stirred at reflux for 48 h until an orange solution was obtained. Upon removal of solvent, the orange oily solid was washed with hexane (3 \times 10 mL) and cold acetone (10 mL). Recrystallization by diffusion of diethyl ether into a CH₂Cl₂ solution afforded orange crystals: yield 0.20 g, 74%. Anal. Calcd for C₁₈H₁₃N₂S₂PtCl: C, 39.17; H, 2.37; N, 5.08. Found: C, 39.25; H, 2.24; N, 5.04. FAB-MS: *m/z* 552 [M⁺], 516 [M⁺ - Cl]. IR (Nujol): ν = 1603, 1549, 1516 cm⁻¹. ¹H NMR (CD₂Cl₂): 6.07 (d, 1H, *J* = 4.8 Hz), 7.03 (m, 2H), 7.22–7.33 (m, 3H), 7.46 (d, 1H, *J* = 5.0 Hz), 7.71–7.98 (m, 4H), 9.15 (d with broad ¹⁹⁵Pt satellites, 1H, ³*J*_{HH} = 5.8 Hz, ³*J*_{PtH} = 47 Hz), 9.41 (d with broad ¹⁹⁵Pt satellites, 1H, ³*J*_{HH} = 5.3 Hz, ³*J*_{PtH} = 42 Hz). ¹³C{¹H} NMR (CD₂Cl₂): 117.5, 119.8, 124.0, 126.5, 127.4, 128.4, 129.7, 130.1, 130.4, 138.3, 139.8, 141.2, 145.4, 151.4, 155.1, 156.0, 163.4.

[Pt(Thpy)(pz)]₂, 2a. A mixture of pyrazole (0.03 g, 0.40 mmol) and excess NaH in THF (10 mL) under a N₂ atmosphere was stirred for 30 min until the evolution of hydrogen ceased. The solution was filtered, added to **1a** (0.20 g, 0.36 mmol) in CH₂Cl₂ (30 mL), and stirred under reflux for 30 h. The resultant orange solution was concentrated to 5 mL, and addition of diethyl ether afforded an orange solid. Recrystallization by diffusion of diethyl ether into a dichloromethane solution afforded orange crystals: yield 0.11 g, 72%. Anal. Calcd for C₂₄H₁₈N₆S₂Pt₂: C, 34.12; H, 2.15; N, 9.95. Found: C, 33.92; H, 1.94; N, 10.00. FAB-MS: *m/z* 844 [M⁺], 777 [M⁺ - pz]. IR (Nujol): ν = 1605, 1563, 1520 cm⁻¹. ¹H NMR (CD₂Cl₂): 6.45 (virtual t, 2H, H(4) of pz), 6.78–6.91 (m, 4H), 7.35–7.42 (m, 4H), 7.62–7.78 (m, 6H), 8.15 (d with v. broad ¹⁹⁵Pt satellites, 1H, ³*J*_{HH} = 5.7 Hz), 8.30 (d with v. broad ¹⁹⁵Pt satellites, 1H, ³*J*_{HH} = 6.0 Hz). ¹³C{¹H} NMR (CDCl₃): 105.7, 106.0, 106.1, 117.5, 117.6, 119.1, 119.4, 127.6, 127.8, 132.8, 133.0, 138.1, 138.4, 138.8, 138.9, 139.9, 140.1, 149.2, 149.3, 149.4, 150.1, 163.6, 163.9.

[Pt(Thpy)(aza)]₂, 3a. The procedure for **2a** was adopted using 7-azaindole (0.04 g, 0.38 mmol), excess NaH, and **1a** (0.20 g, 0.36 mmol) to afford red crystals: yield 0.13 g, 76%. Anal. Calcd for C₃₂H₂₂N₆S₂Pt₂: C, 40.68; H, 2.35; N, 8.89. Found: C, 40.43; H, 2.43; N, 8.63. FAB-MS: *m/z* 944 [M⁺], 827 [M⁺ - aza]. IR (Nujol): ν = 1607, 1559 cm⁻¹. ¹H NMR (CD₂Cl₂): 6.23 (d, 1H, *J* = 4.7 Hz), 6.33 (d, 1H, *J* = 3.8 Hz), 6.62 (t, 1H, *J* = 6.0 Hz), 6.75–7.35 (m, 10H), 7.53 (d, 1H, *J* = 4.8 Hz), 7.60–7.90 (m, 6H), 8.11 (d with v. broad ¹⁹⁵Pt satellites, 1H, *J* = 3.0 Hz), 9.71 (d with v. broad ¹⁹⁵Pt satellites, 1H, *J* = 4.8 Hz). ¹³C{¹H} NMR (DMSO-*d*₆): 99.6–101.1, 114.2–122.2, 125.1–133.2, 135.9–142.3, 146.3–146.8, 159.2, 159.6.

[¹⁸⁵Pt(Bzqn)Cl]₂, 1b. The procedure for **1a** was adopted using 7,8-benzoquinoline (0.36 g, 1.99 mmol), K₂PtCl₄ (0.20 g, 0.48 mmol), and ¹⁸⁵PtCl₄ (0.58 g, 2.09 mmol) to yield yellow crystals: yield 0.16 g, 50%. Anal. Calcd for C₂₉H₁₄N₂Cl₂Pt: C, 50.73; H, 6.46; N, 4.08. Found: C, 50.96; H, 6.45; N, 3.78. FAB-MS (–ve): *m/z* 444 [M⁻]. IR (Nujol): ν = 1621, 1592,

(13) (a) Tse, M. C.; Cheung, K. K.; Chan, M. C. W.; Che, C. M. *Chem. Commun.* **1998**, 2295. (b) Lai, S. W.; Chan, M. C. W.; Cheung, K. K.; Che, C. M. *Organometallics*, in press.

(14) (a) Zheng, G. Y.; Rillema, D. P.; DePriest, J.; Woods, C. *Inorg. Chem.* **1998**, *37*, 3588. (b) Zheng, G. Y.; Rillema, D. P. *Inorg. Chem.* **1998**, *37*, 1392.

(15) Liu, H. Q.; Peng, S. M.; Che, C. M. *J. Chem. Soc., Chem. Commun.* **1995**, 509.

(16) (a) Wu, L. Z.; Cheung, T. C.; Che, C. M.; Cheung, K. K.; Lam, M. H. W. *Chem. Commun.* **1998**, 1127. (b) Wong, K. H.; Chan, M. C. W.; Che, C. M. *Chem. Eur. J.*, in press.

(17) Lai, S. W.; Chan, M. C. W.; Peng, S. M.; Che, C. M. *Angew. Chem., Int. Ed.* **1999**, *38*, 669.

(18) Marshall, J. L.; Stobart, S. R.; Gray, H. B. *J. Am. Chem. Soc.* **1984**, *106*, 3027.

(19) Bailey, J. A.; Miskowski, V. M.; Gray, H. B. *Inorg. Chem.* **1993**, *32*, 369.

(20) Craig, C. A.; Garces, F. O.; Watts, R. J.; Palmans, R.; Frank, A. J. *Coord. Chem. Rev.* **1990**, *97*, 193.

(21) Perrin, D. D.; Armarego, W. L. F.; Perrin, D. R. *Purification of Laboratory Chemicals*, 2nd ed.; Pergamon: Oxford, 1980.

(22) Demas, J. N.; Crosby, G. A. *J. Phys. Chem.* **1971**, *75*, 991.

1569 cm^{-1} . ^1H NMR (CD_3CN): 0.91 (t, 12H, $J = 7.3$ Hz, $\text{CH}_3(\text{CH}_2)_3$), 1.28 (m, 8H, $\text{CH}_3(\text{CH}_2)_3$), 1.54 (m, 8H, $\text{CH}_3(\text{CH}_2)_3$), 3.06 (m, 8H, $\text{CH}_3(\text{CH}_2)_3$), 7.43–7.83 (m, 5H), 8.06 (d with broad ^{195}Pt satellites, 1H, $^3J_{\text{HH}} = 7.3$ Hz, $^3J_{\text{PtH}} = 43$ Hz), 8.42 (d, 1H, $J = 8.0$ Hz), 9.99 (d with broad ^{195}Pt satellites, 1H, $^3J_{\text{HH}} = 5.5$ Hz, $^3J_{\text{PtH}} = 47$ Hz). $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_3CN): 13.6; 20.3; 24.4; 59.4 ($\text{CH}_3(\text{CH}_2)_3$), 120.9, 122.3, 124.2, 127.5, 129.7, 129.8, 130.8, 134.3, 137.3, 140.0, 149.5, 156.6, 158.2.

[Pt(Bzqn)(pz)]₂, 2b. The procedure for **2a** was adopted using pyrazole (0.02 g, 0.29 mmol), excess NaH, and **1b** (0.18 g, 0.26 mmol). A yellow solution was obtained after stirring at reflux for 12 h under a N_2 atmosphere. Recrystallization by slow evaporation from an acetone/DMF mixture yielded yellow crystals: yield 0.08 g, 72%. Anal. Calcd for $\text{C}_{32}\text{H}_{22}\text{N}_6\text{Pt}_2$: C, 43.64; H, 2.52; N, 9.54. Found: C, 43.45; H, 2.78; N, 9.67. FAB-MS: m/z 879 [M^+], 812 [$\text{M}^+ - \text{pz}$]. IR (Nujol): $\nu = 1621, 1594, 1568$ cm^{-1} . ^1H NMR (CD_2Cl_2): 6.58 (virtual t, 2H, H(4) of pz), 7.33–7.57 (m, 10H), 7.74–7.80 (m, 4H), 7.94 (d, 2H, $J = 7.2$ Hz), 8.31 (t, 2H, $J = 6.6$ Hz), 8.67 (d with v. broad ^{195}Pt satellites, 1H, $J = 4.9$ Hz), 8.79 (d with v. broad ^{195}Pt satellites, 1H, $J = 5.3$ Hz). $^{13}\text{C}\{^1\text{H}\}$ NMR (DMSO- d_6): 106.3, 106.4, 106.5, 121.2, 121.3, 121.9, 122.0, 123.2, 123.4, 123.5, 126.4, 128.7, 128.8, 129.0, 129.2, 129.4, 130.5, 132.9, 138.1, 138.3, 139.1, 141.4, 141.5, 141.8, 142.6, 148.1, 148.3, 156.0, 156.1.

[Pt(Bzqn)(aza)]₂, 3b. The procedure for **2a** was adopted using 7-azaindole (0.04 g, 0.30 mmol), excess NaH, and **1b** (0.18 g, 0.26 mmol). An orange solution was obtained after stirring at reflux for 12 h under a N_2 atmosphere. Recrystallization by diffusion of diethyl ether into a dichloromethane solution yielded an orange microcrystalline solid: yield 0.09 g, 74%. Anal. Calcd for $\text{C}_{40}\text{H}_{26}\text{N}_6\text{Pt}_2$: C, 48.98; H, 2.67; N, 8.57. Found: C, 49.13; H, 2.43; N, 8.30. FAB-MS: m/z 979 [M^+], 862 [$\text{M}^+ - \text{aza}$]. IR (Nujol): $\nu = 1620, 1596, 1576$ cm^{-1} . ^1H NMR (CD_2Cl_2): 5.83 (d, 1H, $J = 7.4$ Hz), 6.67–7.36 (m, 15H), 7.46–7.81 (m, 5H), 8.10–8.25 (m, 3H), 8.36 (s, 1H), 9.12 (d with v. broad ^{195}Pt satellites, 1H, $J = 5.0$ Hz). $^{13}\text{C}\{^1\text{H}\}$ NMR (DMSO- d_6): 100.8–101.3, 101.7, 113.8, 114.4, 121.7–133.3, 135.6–140.2, 146.8, 151.6, 156.5.

[Pt(Phpy)(pz)]₂, 2c. The procedure for **2b** was adopted using pyrazole (0.02 g, 0.32 mmol), excess NaH, and **1c** (0.20 g, 0.30 mmol) to afford yellow crystals: yield 0.10 g, 80%. Anal. Calcd for $\text{C}_{28}\text{H}_{22}\text{N}_6\text{Pt}_2$: C, 40.39; H, 2.66; N, 10.09. Found: C, 40.69; H, 2.42; N, 10.00. FAB-MS: m/z 832 [M^+], 765 [$\text{M}^+ - \text{pz}$]. IR (Nujol): $\nu = 1607, 1584, 1565$ cm^{-1} . ^1H NMR (DMSO- d_6): 6.46 (virtual t, 2H, H(4) of pz), 6.89 (d, 1H, $J = 6.5$ Hz), 6.96–7.07 (m, 5H), 7.30 (m, 2H), 7.58 (d, 1H, $J = 2.2$ Hz), 7.69 (m, 4H), 7.80 (d, 1H, $J = 2.0$ Hz), 8.04 (m, 4H), 8.27 (d, 1H, $J = 5.8$ Hz), 8.41 (d, 1H, $J = 5.7$ Hz). $^{13}\text{C}\{^1\text{H}\}$ NMR (DMSO- d_6): 106.1, 106.2, 106.3, 119.3, 119.4, 122.8, 122.9, 123.1, 123.2, 123.6, 123.7, 129.1, 133.0, 133.1, 137.7, 137.9, 138.6, 138.7, 139.3, 144.3, 145.0, 145.2, 145.3, 148.5, 148.7, 166.8, 167.0.

[Pt(Phpy)(aza)]₂, 3c. The procedure for **2a** was adopted using 7-azaindole (0.04 g, 0.34 mmol), excess NaH, and **1c** (0.20 g, 0.30 mmol) to afford red crystals: yield 0.11 g, 79%. Anal. Calcd for $\text{C}_{36}\text{H}_{26}\text{N}_6\text{Pt}_2$: C, 46.35; H, 2.81; N, 9.01. Found: C, 46.46; H, 2.79; N, 9.14. FAB-MS: m/z 932 [M^+], 815 [$\text{M}^+ - \text{aza}$]. IR (Nujol): $\nu = 1607, 1585, 1557$ cm^{-1} . ^1H NMR (CD_2Cl_2): 5.93 (m, 1H), 6.03 (d, 1H, $J = 7.4$ Hz), 6.23 (m, 1H), 6.44–6.81 (m, 9H), 6.99 (d, 2H, $J = 7.4$ Hz), 7.15–7.75 (m, 8H), 7.91–7.99 (m, 2H), 8.47 (d with v. broad ^{195}Pt satellites, 2H, $J = 5.1$ Hz). $^{13}\text{C}\{^1\text{H}\}$ NMR (DMSO- d_6): 100.1, 100.2, 101.3, 113.4, 117.7, 121.3–124.4, 128.1–131.0, 138.1–140.4, 142.5, 144.7, 149.3, 159.9.

X-ray crystallography. Crystals of **1a** and **2a** were grown by vapor diffusion of diethyl ether into dichloromethane solutions. Crystal data and details of collection and refinement are summarized in Table 1.

For **1a**, diffraction experiments were performed on a Rigaku AFC7R diffractometer ($\lambda = 0.71073$ Å, $\omega - 2\theta$ scans). The structure was solved by Patterson methods, expanded by

Table 1. Crystal Data

	1a	2a
formula	$\text{C}_{18}\text{H}_{13}\text{N}_2\text{S}_2\text{PtCl}$	$\text{C}_{24}\text{H}_{18}\text{N}_6\text{S}_2\text{Pt}_2$
fw	551.98	844.75
color	orange	orange
crystal size, mm	$0.30 \times 0.20 \times 0.08$	$0.45 \times 0.25 \times 0.15$
crystal system	monoclinic	monoclinic
space group	$P2_1/n$ (No. 14)	$P2_1/c$
<i>a</i> , Å	9.078(2)	17.215(2)
<i>b</i> , Å	8.363(1)	9.204(1)
<i>c</i> , Å	23.562(7)	16.735(2)
β , deg	94.57(2)	116.466(8)
<i>V</i> , Å ³	1783.3(6)	2373.8(4)
<i>Z</i>	4	4
<i>D_c</i> (g cm ⁻³)	2.056	2.364
μ , cm ⁻¹	82.22	120.93
<i>F</i> (000)	1048	1567
<i>T</i> , K	301	295
$2\theta_{\text{max}}$, deg	50	50
no. of unique data	3381	4177
no. of obsd data	2666 [$I > 3\sigma(I)$]	2530 [$I > 2\sigma(I)$]
no. of variables	217	361
<i>R_w</i> ^a , <i>R_w</i> ^b	0.037, 0.045	0.031, 0.027
residual ρ , e Å ⁻³	+1.30, -0.94	+1.20, -0.62

$$^a R = \sum |F_o| - |F_c| / \sum |F_o|. \quad ^b R_w = [\sum w(|F_o| - |F_c|)^2 / \sum w|F_o|^2]^{1/2}.$$

Fourier methods (PATTY²³), and refined by full-matrix least-squares using the software package TeXsan²⁴ on a Silicon Graphics Indy computer. One formula unit constitutes a crystallographic asymmetric unit. All 24 non-H atoms were refined anisotropically, and 13 H atoms at calculated positions were not refined.

For **2a**, diffraction experiments were performed on a Nonius diffractometer ($\lambda = 0.71073$ Å, $\theta/2\theta$ scans). The structure was solved by direct methods and refined by least-squares treatment on F^2 using the NRCVAX program. One formula unit constitutes a crystallographic asymmetric unit. All 40 non-H atoms were refined anisotropically, and 26 H atoms at calculated positions were not refined. The two 2-(2'-thienyl)pyridyl groups are disordered with NC(1), NC(2), NC(3), and NC(4) having half carbon and nitrogen atom occupancy and S(1), S(2), S(2'), C(6), C(6'), C(7), C(7'), C(20), C(20'), C(21), and C(21') also having half occupancy.



Results and Discussion

Synthesis and Characterization. A general synthetic route to bi- and triplatinum macrocycles is illustrated in Scheme 1. The monomeric precursors **1a–c** (**a**, Thpy; **b**, Bzqn; **c**, Phpy) comprise two substitutionally labile ligands (HThpy and/or chloride). Treatment with stoichiometric amounts of deprotonated N-donor ligands in $\text{CH}_2\text{Cl}_2/\text{THF}$ under a nitrogen atmosphere afford cyclic dimers with μ -pz (denoted **2**) and μ -aza (denoted **3**) linkers, while trimers are formed with μ -bzim groups (denoted **4**). Syntheses of compounds based on the $[\text{Pt}(\mu\text{-pz})_2\text{Pt}]$ core have been reported.^{2a–c} All complexes are air- and moisture-stable at room temperature in solid and solution states.

These reactions to form macrocycles proceed in moderately high yields (70–90%). We have already established¹⁷ that the coordination geometry of the bzim ligand is suitable for the assembly of trimeric metallo-

(23) PATTY: Beurskens, P. R.; Admiraal, G.; Bosman, W. P.; Garcia-Granda, S.; Gould, R. O.; Smits, J. M. M.; Smykalla, C. The DIRDIF program system; Technical Report of the Crystallography Laboratory, University of Nijmegen: The Netherlands, 1992.

(24) TeXsan: Crystal Structure Analysis Package; Molecular Structure Corporation: The Woodlands, TX, 1985 and 1992.

Scheme 1. Synthetic Route toward Bi- and Trimetallic Macrocycles**Scheme 2. Ligand-Directed Strategy toward Trimetallic Macrocycles**

macrocycles, by coupling of three 150° edges (as for bzim) and three cis binding sites at square planar and octahedral metal centers (Scheme 2). Our unsuccessful attempts to synthesize $[\text{Pt}(\text{Phpy})(\text{bzim})]_3$, which afforded insoluble, intractable solids, are therefore surprising. Nevertheless, the intrinsic binding geometry of the bidentate pz and aza ligands evidently ensure the efficient and selective formation of dimeric Pt(II) complexes. The positive FAB mass spectra of the cyclic dimers and trimers reveal clusters corresponding to the molecular ion, while no other oligomeric species are detected. The ^1H NMR spectra for all complexes contain distinctive low-field doublets with broad ^{195}Pt satellites ($^3J_{\text{PtH}} \sim 40$ Hz) which are assigned to H_β of the cyclometalating ligands. Variable-temperature ^1H NMR studies (-60 to 40°C in CD_2Cl_2) were performed on **4a** to investigate the orientation of the bzim ligands in solution. No significant changes were detected, but the syn, anti, anti conformation in the solid state (see below) is unlikely to be maintained in solution.

Crystal Structures. In this work, the molecular structures of the mononuclear complex **1a** and μ -pyrazolate dimer **2a** (Figures 1 and 2, respectively) were




Figure 1. (Top) Perspective view of $[\text{Pt}(\text{Thpy})(\text{HThpy})\text{Cl}]$, **1a**, with 50% probability ellipsoids. (Bottom) Crystal packing diagram showing π - π stacking interactions.




Figure 2. Perspective view of $[\text{Pt}(\text{Thpy})(\text{pz})]_2$, **2a**, with 35% probability ellipsoids. Thpy ligands (NC(1-4), S(1'), S(2), C(6'), C(7'), C(20), C(21) atoms) are disordered.

determined by X-ray crystallography (Table 2). We shall also elaborate on the solid-state structure of **4a**.

The platinum atom in **1a** resides in an approximately square planar environment and is coordinated to monodentate (pyridyl N) and bidentate (pyridyl N and thienyl C) 2-(2'-thienyl)pyridyl ligands plus a chloride group (Figure 1). The Pt(1)-Cl(1) bond length of $2.402(2)$ Å is partially longer than that reported for $\text{Pt}(\text{bpy})\text{Cl}_2$ ($2.306(2)$ Å)²⁵ due to the greater trans influence of the

(25) Osborn, R. S.; Rogers, D. *J. Chem. Soc., Dalton Trans.* **1974**, 1002.

Table 2. Selected Bond Lengths (Å) and Angles (deg)



Complex 1a			
Pt(1)–Cl(1)	2.402(2)	Pt(1)–C(7)	1.990(9)
Pt(1)–N(1)	2.060(7)	S(1)–C(6)	1.73(1)
Pt(1)–N(2)	2.032(7)	S(1)–C(9)	1.71(1)
N(1)–Pt(1)–C(7)	80.4(3)	Cl(1)–Pt(1)–N(2)	89.3(2)
Cl(1)–Pt(1)–N(1)	97.3(2)	Cl(1)–Pt(1)–C(7)	177.3(3)
Complex 2a			
Pt(1)–NC(1)	1.995(9)	Pt(2)–N(6)	2.031(8)
Pt(1)–NC(2)	1.986(9)	Pt(2)–N(8)	2.043(8)
Pt(1)–N(5)	2.032(8)	S(2)–C(17)	1.799(15)
Pt(1)–N(7)	2.056(8)	S(2)–C(18)	1.72(1)
Pt(2)–NC(3)	2.017(9)	S(1')–C(5)	1.88(1)
Pt(2)–NC(4)	2.017(8)	S(1')–C(8)	1.804(14)
NC(1)–Pt(1)–NC(2)	79.1(4)	NC(4)–Pt(2)–N(8)	96.3(3)
NC(1)–Pt(1)–N(5)	96.6(4)	N(6)–Pt(2)–N(8)	88.3(3)
NC(2)–Pt(1)–N(7)	95.9(3)	Pt(1)–N(5)–N(6)	122.8(6)
N(5)–Pt(1)–N(7)	88.4(3)	Pt(2)–N(6)–N(5)	120.8(6)
NC(3)–Pt(2)–NC(4)	79.5(4)	C(5)–S(1')–C(8)	90.9(6)
NC(3)–Pt(2)–N(6)	95.9(4)	C(17)–S(2)–C(18)	92.5(6)

thienyl carbanion compared to the pyridyl N donor. The stacked arrangement of **1a** monomers (bottom of Figure 1) indicates π – π interactions between adjacent monodentate HThpy ligands, with interplanar separations of ca. 3.5 Å. The structure of the iodide congener has been reported.²⁶

In **2a**, two [Pt(Thpy)] moieties are bridged by two pyrazolate ligands in an exo-bidentate fashion (Figure 2). The central six-membered [Pt(μ -pz)₂Pt] core is non-planar, with mean Pt–N(pz) distances of 2.04 Å and N–Pt–N angles of ca. 88° between the pz groups. The metal atoms are positioned at the vertexes of a boat conformation like in previously reported Pt(II)^{2b,c} and Ir(I)²⁷ μ -pyrazolate dimers. The Pt(1)–Pt(2) separation of 3.4863(6) Å in **2a** is comparable to that in the mono-(pyrazolyl) species [Pt₂(tpy)₂(μ -pz)](ClO₄)₃ (3.432(1) Å),²⁸ but is longer than the corresponding metal–metal distance in [PtCl(PPh₂Me)(μ -3,5-Me₂pz)]₂ (3.170(1) Å).^{2c} Comparable bond lengths within the pyrazolate rings suggest substantial delocalization upon deprotonation. The bite angles of the cyclometalating Thpy ligands (79.1(4)° and 79.5(4)°) are comparable to that in **1a** (80.4(3)°).

The “two-up, one-down” orientation of the bzim ligands in the molecular structure of **4a**, which allows comparisons with the partial cone conformation in calixarenes²⁹ and the Pt(II)-incorporated analogue of calix[4]arene [Pt(en)(uracilate)]₄⁴⁺ (en = 1,2-diaminoethane),³⁰ has been described previously.¹⁷ From the partial crystal packing diagram (Figure 3), π – π interactions of ca. 3.5 Å between Thpy systems in adjacent molecules are apparent, and these are expected to influence the photophysical behavior of **4a** in the solid state (see below).

Absorption and Emission Spectroscopy. The UV–visible spectral data of the Pt(II) complexes are listed in Table 3. Their emission data in fluid solution

**Figure 3.** Partial packing diagram of [Pt(Thpy)(bzim)]₃, **4a** (π – π interactions indicated by asterisks).**Figure 4.** UV–vis absorption spectra of **1a** and **2a** in dichloromethane at 298 K.

and in crystalline state are given in Tables 4 and 5, respectively.

Complexes with Thpy Ligands. The UV–vis absorption spectra of **2–4a** in CH₂Cl₂ are comparable to that for the precursor **1a** (Figure 4 for **1a** and **2a**). For example, the spectrum of **4a** contains high-energy bands ($\lambda < 360$ nm) which are dominated by spin-allowed intraligand (¹IL: π (Thpy) \rightarrow π^* (Thpy)) with metal perturbation transitions.³¹ The moderately intense low-energy bands with λ_{\max} in the range 409–421 nm and the notably weaker, structured absorptions at λ_{\max} 480–554 nm ($\epsilon < 120$ dm³ mol^{–1} cm^{–1}) are assigned to the spin-allowed and spin-forbidden metal-to-ligand charge transfer (¹MLCT and ³MLCT: (5d)Pt \rightarrow π^* (Thpy)) transitions, respectively.

(26) Giordano, T. J.; Rasmussen, P. G. *Inorg. Chem.* **1975**, *14*, 1628.(27) Coleman, A. W.; Eadie, D. T.; Stodart, S. R.; Zaworotko, M. J.; Atwood, J. L. *J. Am. Chem. Soc.* **1982**, *104*, 922.(28) Bailey, J. A.; Gray, H. B. *Acta Crystallogr.* **1992**, *C48*, 1420.(29) Gutsche, C. D. *Calixarenes*; Royal Society of Chemistry: Cambridge, 1989.(30) Rauter, H.; Hillgeris, E. C.; Erxleben, A.; Lippert, B. *J. Am. Chem. Soc.* **1994**, *116*, 616.(31) Kvam, P.-I.; Puzyk, M. V.; Cotlyr, V. S.; Balashev, K. P.; Songstad, J. *Acta Chem. Scand.* **1995**, *49*, 645.

Table 3. UV–Vis Absorption Data in Dichloromethane at 298 K

complex	λ_{\max}/nm ($\epsilon/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$)
[Pt(Thpy)(HThpy)Cl], 1a	262 (22300), 300 (23900), 329 (sh, 17600), 406 (4600), 419 (4550), 509 (60), 553 (40)
[Pt(Thpy)(pz)] ₂ , 2a	293 (39400), 354 (26800), 410 (sh, 3800), 512 (60), 555 (40)
[Pt(Thpy)(aza)] ₂ , 3a	292 (42900), 354 (22800), 474 (sh, 2100)
[Pt(Thpy)(bzim)] ₃ , 4a	277 (47000), 337 (22000), 357 (18300), 409 (5250), 421 (4800), 480 (110), 512 (100), 554 (70)
[ⁿ Bu ₄ N][Pt(Bzqn)Cl ₂], 1b	280 (28700), 358 (4350), 374 (4400), 434 (1300)
[Pt(Bzqn)(pz)] ₂ , 2b	286 (35100), 370 (12300), 388 (sh, 8850), 421 (sh, 3150), 465 (sh, 250)
[Pt(Bzqn)(aza)] ₂ , 3b	285 (34400), 380 (sh, 8700), 410 (sh, 4000)
[Pt(Bzqn)(bzim)] ₃ , 4b	271 (70000), 278 (73000), 358 (14000), 372 (14000), 413 (4000), 469 (sh, 440)
[ⁿ Bu ₄ N][Pt(Phpy)Cl ₂], 1c	259 (23700), 287 (15200), 331 (6300), 379 (4200), 426 (sh, 780), 490 (120)
[Pt(Phpy)(pz)] ₂ , 2c	257 (45600), 284 (24500), 329 (12200), 359 (11000), 406 (3350), 446 (sh, 310)
[Pt(Phpy)(aza)] ₂ , 3c	266 (33600), 290 (36200), 345 (sh, 17500), 410 (sh, 3450), 459 (sh, 1400)


Table 4. Emission Data in CH₂Cl₂ (complex concentration 5 × 10⁻⁵ M)

complex	298 K: λ_{\max}/nm ; $\tau_0/\mu\text{s}$; ϕ_0	77 K: λ_{\max}/nm
[Pt(Thpy)(HThpy)Cl], 1a	556 (max), 577 (sh), 602, 628 (sh), 661 (sh); 12.0; 0.19	553 (max), 573 (sh), 600, 624 (sh), 654 (sh)
[Pt(Thpy)(pz)] ₂ , 2a	559 (max), 581 (sh), 606, 632 (sh), 664 (sh); 15.5; 0.18	557 (max), 576 (sh), 602, 624 (sh), 655 (sh)
[Pt(Thpy)(aza)] ₂ , 3a	561 (max), 579 (sh), 606, 638 (sh), 661 (sh); 3.1; 0.012	557 (max), 578 (sh), 602, 662 (sh)
[Pt(Thpy)(bzim)] ₃ , 4a	560 (max), 580 (sh), 606, 636 (sh), 661 (sh); 10.9; 0.016	557 (max), 575 (sh), 601, 633 (sh), 656 (sh)
[ⁿ Bu ₄ N][Pt(Bzqn)Cl ₂], 1b	nonemissive	494, 532, 572, 620 (sh)
[Pt(Bzqn)(pz)] ₂ , 2b	497, 535, 588 (sh); 1.6; 9.9 × 10 ⁻³	486, 522, 560
[Pt(Bzqn)(aza)] ₂ , 3b	nonemissive	497, 530, 565
[Pt(Bzqn)(bzim)] ₃ , 4b	nonemissive	nonemissive
[ⁿ Bu ₄ N][Pt(Phpy)Cl ₂], 1c	nonemissive	485 (max), 527, 557, 578, 604, 631 (sh), 668 (sh)
[Pt(Phpy)(pz)] ₂ , 2c	487, 522, 556, 600 (sh); 1.3; 0.039	485 (max), 515, 544, 596 (sh)
[Pt(Phpy)(aza)] ₂ , 3c	494, 520, 556, 601 (sh); 0.29; 4.5 × 10 ⁻⁴	489, 523 (max), 557, 605 (sh)

Table 5. Solid-State Emission Data

complex	298 K: λ_{\max}/nm	77 K: λ_{\max}/nm
[Pt(Thpy)(HThpy)Cl], 1a	563, 581, 606 (max), 632, 651	574 (max), 598, 625, 652 (sh), 690 (sh)
[Pt(Thpy)(pz)] ₂ , 2a	570 (sh), 591 (sh), 616 (max), 665	577 (max), 595, 624, 651 (sh)
[Pt(Thpy)(aza)] ₂ , 3a	556, 601 (max), 658	552 (max), 599, 677
[Pt(Thpy)(bzim)] ₃ , 4a	571, 617, 664 (max)	579 (max), 603 (sh), 630, 659 (sh), 691 (sh)
[ⁿ Bu ₄ N][Pt(Bzqn)Cl ₂], 1b	525 (sh), 542 (max), 591 (sh)	514 (max), 554, 597 (sh)
[Pt(Bzqn)(pz)] ₂ , 2b	533, 568 (max), 633	534 (max), 568, 615 (sh)
[Pt(Bzqn)(aza)] ₂ , 3b	550 (max), 592, 640 (sh)	517, 548 (max), 598, 632 (sh)
[Pt(Bzqn)(bzim)] ₃ , 4b	nonemissive	512 (max), 549, 593 (sh)
[ⁿ Bu ₄ N][Pt(Phpy)Cl ₂], 1c	502, 536, 566 (sh)	498 (max), 515, 526, 536, 558, 569
[Pt(Phpy)(pz)] ₂ , 2c	505, 539, 557 (sh)	506, 526, 549 (max), 596 (sh)
[Pt(Phpy)(aza)] ₂ , 3c	490, 524, 555	493, 528, 563

Structured emissions are observed for complexes **1–4a** in CH₂Cl₂ at room temperature, with peak maxima in the range 556–561 (sh 577–581) and 602–606 (sh 628–638, 661–664) nm (Table 4). Minor solvatochromic effects are detected for the emission energy, e.g., for **4a**, $\lambda_{\max} = 564$ (C₆H₆), 560 (CH₂Cl₂), 562 (CH₃CN), 567 nm (CH₃OH). Self-quenching of the emission has been detected for **1–4a**; in each case, a linear plot of $1/\tau$ against complex concentration was obtained. For example, k_q values of 3.5×10^7 and $3.0 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ have been obtained for **2a** and **4a**, respectively. The μ -pz complex **2a** emits with a longer intrinsic lifetime ($\tau_0 = 15.5 \mu\text{s}$) and a significantly higher quantum yield (0.18) than the μ -aza dimer **3a** ($\tau_0 = 3.1 \mu\text{s}$, $\phi_0 = 0.012$) and the trimer **4a** ($\tau_0 = 10.9 \mu\text{s}$, $\phi_0 = 0.016$). Indeed, the photoluminescent properties displayed by **2a** are substantially superior to those for previously reported binuclear Pt(II) luminophores, which are weakly emissive in solution at room temperature.¹⁹ At 77 K, emission bands are slightly blue-shifted to λ_{\max} 553–557 (sh 573–578) and 600–602 (sh 624–662) nm. With reference to earlier work^{31,32} and the absence of any major solvatochromic effects (<10 nm), we suggest that these emissions are more appropriately assigned to transitions with mixed MLCT and IL character.

**Figure 5.** Solid-state emission spectrum of **4a** ($\lambda_{\text{ex}} = 380$ nm) at 298 K (asterisk denotes instrumental artifact).

The solid-state emission spectrum of **4a** at 298 K (Figure 5) contains three bands at λ_{\max} 571, 617, and 664 (max) nm, all of which undergo nonexponential decay. Similarly, complexes **1–3a** display multiple emissions (including low-energy bands at 651–665 nm) in crystalline form at 298 K, and these are generally red-shifted from the solution-state emission (Table 5). This can be attributed to excimeric emission arising from weak π – π stacking of square planar Pt(II) species

(32) Maestri, M.; Sandrini, D.; Balzani, V.; Chassot, L.; Jolliet, P.; von Zelewsky, A. *Chem. Phys. Lett.* **1985**, *122*, 375.

in the solid state,³³ since these interactions are evident from the crystal structures of **1a** (bottom of Figure 1) and **4a** (Figure 3). Blue-shifts in emission maxima are exhibited upon cooling to 77 K. For example, the solid emission of **4a** at 77 K shows bands at λ_{max} 579 (sh 603, 630, 659, 691) nm, which are higher in energy than the 664 nm emission at 298 K. These solid-state emissions are proposed to originate from ³MLCT excited states.

Complexes with Bzqn and Phpy Ligands. The UV-vis absorption spectra of **2–4b** in CH₂Cl₂ are similar to that for **1b** and the bis(cyclometalated) derivative [Pt(Bzqn)₂].³² High-energy absorption bands with $\lambda_{\text{max}} < 390$ nm are ascribed to ¹IL transitions. The moderately intense bands at λ_{max} 410–421 nm and the significantly weaker low-energy absorptions ($\lambda_{\text{max}} > 450$ nm) are assigned to ¹MLCT and ³MLCT transitions, respectively.

In contrast to the Thpy complexes, a number of Bzqn derivatives are nonemissive in fluid solution. At room temperature, only the μ -pz species **2b** is luminescent and displays structured emission at 497, 535, and 588 (sh) nm ($\tau_0 = 1.6 \mu\text{s}$, $\phi_0 = 9.9 \times 10^{-3}$). A blue-shifted 77 K luminescence at 486, 522, and 560 nm is observed, and we tentatively assign the emission as mixed ³IL/³MLCT. The μ -aza complex **3b** shows structured emission at 77 K only, while the μ -bzim trimer **4b** is nonemissive at 298 and 77 K in solution. The solid-state emissions of **1–3b** are structured at 298 K, and blue-shifts are again observed upon cooling to 77 K.

The UV-vis absorption spectrum of the previously described **1c**²⁰ resembles those of **2c** and **3c** bearing the Phpy ligand, where ¹IL, ¹MLCT, and ³MLCT transitions ($\lambda < 370$, $\lambda_{\text{max}} = 406$ –410, and $\lambda > 440$ nm, respectively) can be assigned. While the precursor **1c** is nonemissive in solution at 298 K, **2c** and **3c** exhibit structured emission at λ_{max} 487–556 (sh 600) nm. A longer lifetime and greater quantum yield are again detected for the μ -pz complex **2c** ($\tau_0 = 1.3 \mu\text{s}$, $\phi_0 = 0.039$) compared to the μ -aza dimer **3c** ($\tau_0 = 0.29 \mu\text{s}$, $\phi_0 = 4.5 \times 10^{-4}$).

Concluding Remarks. Our systematic photophysical investigation has revealed several trends. First, complexes bearing the Thpy group and in particular μ -pyrazolate species display superior luminescent properties. Second, the energy of the emissions and lowest energy UV-vis bands for Thpy oligomers are generally red-shifted relative to the Bzqn and Phpy congeners. This can be illustrated by the 77 K emission spectra of **2a–c** in CH₂Cl₂ (Figure 6) and evidently reflects differences in the energies of the π^* orbitals for the cyclometalating ligands. The relatively low-energy ex-




Figure 6. Normalized emission spectra of **2a–c** ($\lambda_{\text{ex}} = 380$ nm) in dichloromethane at 77 K.

cited states for Thpy derivatives result in greater energy differences between the MLCT/IL and upper-lying non-emissive MC states. Hence such complexes, in particular [Pt(Thpy)(pz)]₂ (**2a**), exhibit improved emissive parameters compared to known Pt(II) binuclear complexes and are potentially interesting for photoinduced energy transfer and sensing applications.

We have described herein a simple yet efficient methodology for the selective synthesis of di- and trimeric platinum(II) macrocycles. Like the recently reported examples of molecular rectangles,³⁴ the trimetallic molecular pockets **4a** and **4b** have lower symmetry than tetrametallic square analogues.³⁵ While the cavity size of **4a** and **4b** pre-empts their employment as molecular hosts, this study has conceived a new direction for development using cyclometalated platinum(II) luminophores.

Acknowledgment. We are grateful for financial support from the University of Hong Kong and the Research Grants Council of the Hong Kong SAR, China [HKU 7298/99P].

Supporting Information Available: Tables of crystal data, atomic coordinates, calculated coordinates, anisotropic displacement parameters, and bond lengths and angles for **1a** and **2a**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM990342H

(34) (a) Benkstein, K. D.; Hupp, J. T.; Stern, C. L. *Inorg. Chem.* **1998**, *37*, 5404. (b) Woessner, S. M.; Helms, J. B.; Shen, Y.; Sullivan, B. P. *Inorg. Chem.* **1998**, *37*, 5406.

(35) Hunter, C. A. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 1079.

(33) Houlding, V. H.; Miskowski, V. M. *Coord. Chem. Rev.* **1991**, *111*, 145.