

Synthesis and Characterization of the W–Os Heterometallic Complex $\text{Os}_3(\text{CO})_{10}(\mu\text{-H})(\mu\text{-}\eta^3\text{-C(=CHPh)C}\equiv\text{CW(O)}_2(\text{C}_5\text{Me}_5))$: Evidence of Hydride Dislocation on a Triosmium Framework by Crystal Polymorphism

Te-Kun Huang,[†] Yun Chi,^{*,†} Shie-Ming Peng,^{*,†} and Gene-Hsiang Lee[‡]

Department of Chemistry, National Tsing Hua University, Hsinchu 30043, Taiwan, and Department of Chemistry and Instrumentation Center, National Taiwan University, Taipei 10764, Taiwan

Received November 30, 1998

The heterometallic compound $\text{Os}_3(\text{CO})_{10}(\mu\text{-H})(\mu\text{-}\eta^3\text{-C(=CHPh)C}\equiv\text{CW(O)}_2(\text{C}_5\text{Me}_5))$ (**1**), which possesses a bridging hydride and a $\text{C}\equiv\text{CC}=\text{CHPh}$ ligand, is obtained on addition of the diyne complex $(\text{C}_5\text{Me}_5)\text{W(O)}_2(\text{C}\equiv\text{CC}\equiv\text{CPh})$ to $\text{Os}_3(\mu\text{-H})_2(\text{CO})_{10}$. ¹H NMR spectra show the occurrence of three isomers at equilibrium in solution, of which two isomers have been unambiguously characterized by an X-ray diffraction study. During subsequent tests of reactivity we isolated $\text{Os}_3(\text{CO})_{10}(\mu\text{-}\sigma\text{-}\eta^2\text{-C}\equiv\text{CCHCHPh})(\mu\text{-H})$ (**2**) and $(\text{C}_5\text{Me}_5)\text{W}(\mu\text{-O})_2\text{Os}_3(\text{CO})_9(\mu\text{-}\sigma\text{-}\eta^2\text{-C}\equiv\text{CCHCHPh})$ (**3**) through two distinct processes initiated by removal of the $(\text{C}_5\text{Me}_5)\text{W(O)}_2$ fragment and decarbonylation, respectively.

Examination of the bonding of ligated hydrocarbon moieties in metal cluster compounds provides detailed data that may be applicable to the characterization of chemisorbed intermediates on metal surfaces.¹ While the chemistry of cluster complexes with small carbon-rich ligands such as carbide, dicarbide, and alkynyl ligands has been investigated,² it is only recently that the higher analogues of diyne or polyynyl fragments have received similar attention as a result of the emerging synthetic application of materials possessing extended carbon unsaturation.³ This new trend is evident from several reports devoted to the syntheses and characterization of monometallic or dimetallic diyne and polyynyl complexes.⁴

In seeking to extend these studies, the chemistry of diyne/metal complexes with organic substrates or with transition-metal clusters has been investigated.⁵ It is shown that the coupling occurred at the alkyne unit farther from the metal atom, displaying a reactivity

pattern influenced by steric effects. We report here a related coupling reaction of $(\text{C}_5\text{Me}_5)\text{W(O)}_2(\text{C}\equiv\text{CC}\equiv\text{CPh})$ with the dihydride complex $\text{Os}_3(\mu\text{-H})_2(\text{CO})_{10}$, which proceeds smoothly to give the product $\text{Os}_3(\text{CO})_{10}(\mu\text{-H})(\mu\text{-}\eta^3\text{-C(=CHPh)C}\equiv\text{CW(O)}_2(\text{C}_5\text{Me}_5))$ (**1**), in which the organic ligand $\text{C}\equiv\text{CC}=\text{CHPh}$ is formed by facile hydride transfer to the diyne unit, a result consistent with the expected higher activity of $\text{Os}_3(\mu\text{-H})_2(\text{CO})_{10}$.⁶ Moreover, X-ray diffraction analyses of two single crystals of **1** provide not only unambiguous structural identification of the linear $\mu\text{-}\eta^3\text{-C}\equiv\text{CC}=\text{CHPh}$ ligand but also a unique case for hydride dislocation on the triosmium framework by crystal polymorphism.

Experimental Section

General Information and Materials. Infrared spectra were recorded on a Perkin-Elmer 2000 FT-IR spectrometer. ¹H and ¹³C NMR spectra were recorded on a Bruker AM-400

[†] National Tsing Hua University.

[‡] National Taiwan University.

(1) (a) Thimmappa, B. H. S. *Coord. Chem. Rev.* **1995**, *143*, 1. (b) Zaera, F. *Chem. Rev.* **1995**, *95*, 2651.

(2) (a) John, K. D.; Geib, S. J.; Hopkins, M. D. *Organometallics* **1996**, *15*, 4357. (b) Jensen, M. P.; Phillips, D. A.; Sabat, M.; Shriver, D. F. *Organometallics* **1992**, *11*, 1859. (c) Norton, D. M.; Eveland, R. W.; Hutchison, J. C.; Stern, C.; Shriver, D. F. *Organometallics* **1996**, *15*, 3916. (d) Akita, M.; Moro-oka, Y. *Bull. Chem. Soc. Jpn.* **1995**, *68*, 420. (e) Carty, A. J.; Hogarth, G.; Enright, G.; Frapper, G. *Chem. Commun.* **1997**, 1883. (f) Doherty, S.; Corrigan, J. F.; Carty, A. J.; Sappa, E. *Adv. Organomet. Chem.* **1995**, *37*, 39. (g) Jensen, M. P.; Phillips, D. A.; Sabat, M.; Shriver, D. F. *Organometallics* **1992**, *11*, 1859. (h) Adams, C. J.; Bruce, M. I.; Skelton, B. W.; White, A. H. *Chem. Commun.* **1996**, 975.

(3) (a) Blenkinsop, P.; Taylor, N. J.; Carty, A. J. *J. Chem. Soc., Chem. Commun.* **1995**, 327. (b) Bruce, M. I.; Ke, M.; Low, P. J. *Chem. Commun.* **1996**, 2405. (c) Bruce, M. I.; Denisovich, L. I.; Low, P. J.; Peregudova, S. M.; Ustynyuk, N. A. *Mendeleev Commun.* **1996**, 200. (d) Norton, D. M.; Stern, C. L.; Shriver, D. F.; Jensen, M. P.; Shriver, D. F. *Inorg. Chem.* **1994**, *33*, 2701. (e) Lang, H.; Weber, C. *Organometallics* **1995**, *14*, 4415.

(4) (a) Lang, H. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 547. (b) Lagow, R. J.; Kampa, J. J.; Wei, H.-C.; Battle, S. L.; Genge, J. W.; Laude, D. A.; Harper, C. J.; Bau, R.; Stevens, R. C.; Haw, J. F.; Munson, E. *Science* **1995**, *267*, 362. (c) Bartik, T.; Bartik, B.; Brady, M.; Dembinski, R.; Galdysz, J. A. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 414. (d) Chi, Y.; Carty, A. J.; Blenkinsop, P.; Delgado, E.; Enright, G. D.; Wang, W.; Peng, S.-M.; Lee, G.-H. *Organometallics* **1996**, *15*, 5269. (e) Bartik, B.; Dembinski, R.; Bartik, T.; Arif, A. M.; Gladysz, J. A. *New J. Chem.* **1997**, *21*, 739.

(5) (a) Brady, M.; Weng, W.; Gladysz, J. A. *J. Chem. Soc., Chem. Commun.* **1994**, 2655. (b) Bruce, M. I.; Skelton, B. W.; White, A. H.; Zaitseva, N. N. *J. Chem. Soc., Dalton Trans.* **1996**, 3151. (c) Bruce, M. I.; Ke, M.; Low, P. J. *Chem. Commun.* **1996**, 2405. (d) Gevert, O.; Wolf, J.; Werner, H. *Organometallics* **1996**, *15*, 2806. (e) Bruce, M. I.; Ke, M.; Low, P. J.; Skelton, B. W.; White, A. H. *Organometallics* **1998**, *17*, 3539.

(6) (a) Shapley, J. R.; Park, J.-T.; Churchill, M. R.; Ziller, J. W.; Beanan, L. R. *J. Am. Chem. Soc.* **1984**, *106*, 1144. (b) Chi, Y.; Shapley, J. R.; Churchill, M. R.; Li, Y. J. *Inorg. Chem.* **1986**, *25*, 4165. (c) Green, M.; Orpen, A. G.; Schaverien, C. J. *J. Chem. Soc., Dalton Trans.* **1989**, 1333. (d) Koike, M.; Hamilton, D. H.; Wilson, S. R.; Shapley, J. R. *Organometallics* **1996**, *15*, 4930.

(400.13 MHz) or a Bruker AMX-300 (300.6 MHz) instrument. Mass spectra were measured on a JEOL-HX110 instrument operating in the fast atom bombardment mode (FAB). All reactions were performed under a nitrogen atmosphere using solvents dried with an appropriate reagent. Products were separated on commercially available preparative thin-layer chromatographic plates (Kieselgel 60 F₂₅₄, E. Merck). Elemental analyses were performed at the NSC Regional Instrumentation Center at National Cheng Kung University, Tainan, Taiwan.

Reaction of Os₃(μ-H)₂(CO)₁₀ with (C₅Me₅)W(O)₂(C≡CC=CPh). In a 100 mL reaction flask, a CH₂Cl₂ solution (60 mL) of Os₃(μ-H)₂(CO)₁₀ (180 mg, 0.21 mmol) and the diyne complex (C₅Me₅)W(O)₂(C≡CC=CPh) (100 mg, 0.21 mmol) was heated at reflux for 3 h, during which period the solution changed from red-purple to red-orange. After removal of solvent under vacuum, the oily residue was dissolved in a minimum amount of CH₂Cl₂ and this solution subjected to separation by thin-layer chromatography. Development with a mixture of ethyl acetate, CH₂Cl₂, and hexane (3:1:12) produced orange and dark red bands, which were extracted from silica gel with CH₂Cl₂, affording yellow-orange Os₃(CO)₁₀(μ-σ-η²-C≡CCHCHPh)(μ-H) (**2**; 40.7 mg, 0.042 mmol, 20%) and red Os₃(CO)₁₀(μ-H)(μ-η³-C(=CHPh)C=CW(O)₂(C₅Me₅)) (**1**; 110 mg, 0.083 mmol, 40%). Single crystals of **1**, suitable for X-ray diffraction, were obtained from a mixture of CH₂Cl₂ and heptane or from a solution of THF and methanol at -20 °C, respectively.

Spectral data of **1** are as follows. MS (FAB, ¹⁹²Os, ¹⁸⁴W): *m/z* 1334 (M⁺). IR (C₆H₁₂): ν(CO) 2123 (vw), 2104 (w), 2068 (vs), 2054 (vs), 2041 (w), 2034 (w), 2021 (vs), 2004 (m), 1997 (m), 1989 (w), 1986 (w, sh), 1975 (vw), 1964 (vw), 1922 (vw, br) cm⁻¹. ¹H NMR (400 MHz, CD₂Cl₂, 240 K): δ 7.27–7.16 (broad, C₆H₅ & CH), 6.89 (s, CH, 0.54 H), 2.10 (s, C₅Me₅, 0.14 H), 2.03 (s, C₅Me₅, 0.32 H), 1.92 (s, C₅Me₅, 0.54 H), -16.91 (s, μ-H, 0.54 H), -17.24 (s, μ-H, 0.14 H). Anal. Calcd for C₃₀H₂₂O₁₂Os₃W: C, 27.11; H, 1.67. Found: C, 27.50; H, 1.78.

Spectral data of **2** are as follows. MS (FAB, ¹⁹²Os): *m/z* 984 (M⁺). IR (C₆H₁₂): ν(CO) 2109 (w), 2070 (vs), 2061 (s), 2024 (vs), 2005 (s), 1990 (w), 1984 (m) cm⁻¹. ¹H NMR (400 MHz, CD₂Cl₂, 220 K): δ 7.71 (d, *J*_{HH} = 7.2 Hz, C₆H₅, 2 H), 7.35–7.23 (m, C₆H₅, 3 H), 6.67 (d, *J*_{HH} = 12.2 Hz, CH), 5.94 (d, *J*_{HH} = 12.2 Hz, CH), -16.51 (s, μ-H). ¹³C NMR (100.6 MHz, CD₂Cl₂, 223 K): CO, δ 182.6, 182.3, 178.4 (2 C), 175.3 (2 C), 171.4 (2 C), 170.0 (2 C); δ 142.4 (C_δH), 135.2 (*i*-C₆H₅), 129.6 (*p*-C₆H₅), 129.0 (*m*-C₆H₅, 2 C), 128.5 (*o*-C₆H₅, 2 C), 106.8 (C_γH), 94.1 (C_β), 79.4 (C_α).

Thermolysis of 1. A CH₂Cl₂ solution (40 mL) of Os₃(CO)₁₀(μ-H)(μ-η³-C(=CHPh)C=CW(O)₂(C₅Me₅)) (**1**; 120 mg, 0.090 mmol) was heated at reflux for 40 h. After removal of the solvent in vacuo, the residue was dissolved in CH₂Cl₂ and subjected to chromatography. Development with a similar mixture of ethyl acetate, CH₂Cl₂, and hexane produced two bands, which were extracted from silica gel to afford yellow-orange Os₃(CO)₁₀(μ-σ-η²-C≡CCHCHPh)(μ-H) (**2**; 48 mg, 0.049 mmol, 54%) and orange-red (C₅Me₅)W(μ-O)₂Os₃(CO)₉(μ-σ-η²-C≡CCHCHPh) (**3**; 28.5 mg, 0.022 mmol, 25%).

Spectral data of **3** are as follows. MS (FAB, ¹⁹²Os, ¹⁸⁴W): *m/z* 1306 (M⁺). IR (C₆H₁₂): ν(CO) 2088 (w), 2069 (vs), 2026 (vs), 2001 (s), 1989 (w), 1973 (vw, br), 1955 (w), 1937 (w) cm⁻¹. ¹H NMR (400 MHz, CDCl₃, 230 K): δ 7.48 (d, *J*_{HH} = 7.4 Hz, C₆H₅), 7.40–7.24 (m, C₆H₅), 6.47 (d, *J*_{HH} = 11.9 Hz, CH), 2.08 (s, C₅Me₅). ¹³C NMR (100.6 MHz, CDCl₃, 230 K): δ 187.2 (1 CO), 185.4 (2 CO), 177.6 (2 CO), 172.4 (2 CO), 168.9 (2 CO), 138.6 (C_δH), 134.8 (*p*-C₆H₅), 128.7 (*m*-C₆H₅), 128.1 (*o*-C₆H₅), 126.5 (*i*-C₆H₅), 115.4 (C₅Me₅), 110.1 (C_γH), 95.3 (C_β), 65.0 (C_α), 12.3 (C₅Me₅). Anal. Calcd for C₂₉H₂₂O₁₁Os₃W: C, 26.77; H, 1.70. Found: C, 26.94; H, 1.75.

X-ray Crystallography. The X-ray diffraction measurements were made with a Nonius CAD-4 or a Siemens Smart CCD diffractometer at room temperature. For complex **1a**, lattice parameters were determined from 25 randomly selected

Table 1. X-ray Structural Data of Complexes 1a and 1b^a

	1a	1b
formula	C ₃₀ H ₂₂ O ₁₂ Os ₃ W	C ₃₀ H ₂₂ O ₁₂ Os ₃ W
mol wt	1328.95	1328.95
diffractometer	Nonius CAD-4	Siemens Smart CCD
cryst syst	monoclinic	monoclinic
space group	<i>C2/c</i>	<i>Cc</i>
<i>a</i> (Å)	34.79(5)	21.6673(3)
<i>b</i> (Å)	12.973(4)	16.4637(2)
<i>c</i> (Å)	14.917(4)	14.5454(1)
β (°)	98.88(3)	131.467(1)
<i>V</i> (Å ³)	6653(3)	3888.08(8)
<i>Z</i>	8	4
<i>D_c</i> (g/cm ³)	2.654	2.270
<i>F</i> (000)	4800	2400
2θ(max) (deg)	50.0	55.0
<i>hkl</i> ranges	-41 to +40, 0-15, 0-17	-28 to +24, -21 to +21, -13 to +18
crystal size (mm)	0.30 × 0.25 × 0.25	0.45 × 0.17 × 0.16
μ(Mo Kα) (cm ⁻¹)	150.08	127.72
transmission: max, min	0.737, 0.693	0.746, 0.362
no. of data in refinement	3908 with <i>I</i> ≥ 2σ(<i>I</i>)	7055
no. of params	419	416
max μ/σ ratio	0.001	0.02
<i>R_f</i> ; <i>R_w</i> or <i>RF</i> ; <i>R_{wF}</i> ²	0.037; 0.033	0.036; 0.077
GOF	1.23	1.05
<i>D</i> -map, max/min (e/Å ⁻³)	1.14/-1.81	1.57/-0.77

^a Features common to all determinations: λ(Mo Kα) = 0.7107 Å; function minimized Σ(w|F_o - F_c|²); weighting scheme w⁻¹ = σ²(F_o) + |g|F_o²; GOF = [Σw|F_o - F_c|²/(N_o - N_v)]^{1/2} (N_o = number of observations; N_v = number of variables).

high-angle reflections. Three standard reflections were monitored every 3600 s. No significant change of intensities (≤2%) was observed during the course of all data collection. Intensities of diffraction signals were corrected for Lorentz, polarization, and absorption effects (ψ scans). The structure was solved with the NRCC-SDP-VAX package. All non-hydrogen atoms had anisotropic temperature factors; hydrogen atoms of organic substituents were placed at calculated positions with *U_H* = *U_C* + 0.1.

Single-crystal X-ray diffraction data of **1b** were measured on a Siemens SMART CCD diffractometer. Reflections were collected using three different φ setting angles; each setting was scanned by 0.3 ω between frames. Each frame was exposed for 5 s. The detector was located 4.0 cm away from the crystal. Crystal decay was monitored by repeating the initial 50 frames at the end of data collection and analyzing the duplicate reflections, and no decay was observed. An empirical absorption correction using SADABS was applied. The structure was solved by the direct method. Hydrogen atoms were fixed at calculated positions and refined using a riding model. Anisotropic displacement parameters were used for all non-H atoms, while the given isotropic displacement parameters were used for H atoms (1.2 and 1.5 times the equivalent isotropic displacement parameter of the atom to which they are respectively attached for methylene H atoms and methyl H atoms). Data collection was carried out by using the SMART program. Cell refinement and data reduction were performed by using the SAINT program. Structure analysis was carried out by using the SHELXTL/PC program.

Crystallographic refinement parameters of **1a** and **1b** are summarized in Table 1; selected bond distances and angles are presented in Tables 2 and 3, respectively.

Results and Discussion

Formation and Characterization of 1. The reaction of the hydridoosmium complex Os₃(μ-H)₂(CO)₁₀ with

Table 2. Selected Bond Distances (Å) and Angles (deg) of 1a (Esd's in Parentheses)



Os(1)–Os(2)	2.959(1)	Os(1)–Os(3)	2.915(1)
Os(2)–Os(3)	2.877(1)	Os(1)–C(11)	2.42(1)
Os(1)–C(12)	2.53(1)	Os(2)–C(13)	2.15(1)
W–C(11)	2.10(1)	W–O(11)	1.71(1)
W–O(12)	1.71(1)	C(11)–C(12)	1.23(2)
C(12)–C(13)	1.43(2)	C(13)–C(14)	1.32(2)
Os(1)–H	1.81(1)	Os(2)–H	1.72(1)
\angle Os(1)–Os(2)–Os(3)	59.91(3)	\angle Os(1)–Os(3)–Os(2)	61.45(3)
\angle Os(2)–Os(1)–Os(3)	58.64(3)	\angle W–C(11)–C(12)	152(1)
\angle C(11)–C(12)–C(13)	173(1)	\angle C(12)–C(13)–C(14)	123(1)
\angle C(11)–W–O(11)	103.9(5)	\angle C(11)–W–O(12)	100.6(5)
\angle O(11)–W–O(12)	108.7(5)		

Table 3. Selected Bond Distances (Å) and Angles (deg) of 1b (Esd's in Parentheses)

Os(1)–Os(2)	2.8451(6)	Os(1)–Os(3)	3.0741(6)
Os(2)–Os(3)	2.8996(7)	Os(1)–C(11)	2.30(1)
Os(1)–C(12)	2.46(1)	Os(2)–C(13)	2.13(1)
W–C(11)	2.15(1)	W–O(11)	1.71(1)
W–O(12)	1.72(1)	C(11)–C(12)	1.24(2)
C(12)–C(13)	1.35(2)	C(13)–C(14)	1.36(2)
\angle Os(1)–Os(2)–Os(3)	64.70(2)	\angle Os(1)–Os(3)–Os(2)	56.80(2)
\angle Os(2)–Os(1)–Os(3)	58.51(2)	\angle W–C(11)–C(12)	147(1)
\angle C(11)–C(12)–C(13)	177(1)	\angle C(12)–C(13)–C(14)	124(1)
\angle C(11)–W–O(11)	101.7(5)	\angle C(11)–W–O(12)	102.8(4)
\angle O(11)–W–O(12)	106.4(5)		

the tungsten acetylide complex $(C_5Me_5)W(O)_2(C\equiv CC\equiv CPh)$ in CH_2Cl_2 at reflux gives the heterometallic compound $Os_3(CO)_{10}(\mu-H)(\mu-\eta^3-C(=CHPh)C\equiv CW(O)_2-(C_5Me_5))$ (**1**) as a red solid. This compound is relatively unstable in solution upon exposure to air; therefore, recrystallization is carried out in a mixture of CH_2Cl_2 and heptane at $-20^\circ C$. FAB MS analysis of **1** shows a parent ion M^+ at m/z 1334 that is consistent with formation of a molecular formula corresponding to a 1:1 combination of starting materials. The IR spectrum in cyclohexane solution exhibits nearly 20 overlapping CO stretching bands in the range $2123\text{--}1922\text{ cm}^{-1}$, indicative of at least two structural isomers. This behavior is confirmed by subsequent 1H NMR analysis. In addition to a broad and complicated signal at δ 7.27–7.16 assigned to aromatic and olefinic hydrogens, we observed three distinctive hydride signals at δ –16.91, –17.24, and –17.73 with the relative intensity ratio 3.9:2.3:1, demonstrating the coexistence of three isomers (Scheme 1). To establish their molecular structures and the interrelationships, we carried out the X-ray diffraction study of a single crystal obtained from CH_2Cl_2 /heptane solution.

The molecular structure of this heterometallic compound labeled **1a** is depicted in Figure 1; selected bond lengths and angles appear in Table 2. The structure contains a triangular osmium framework with 10 terminal CO ligands. The arrangement of CO ligands on this osmium triangle is similar to those of the clusters $Os_3(\mu-X)(\mu-H)(CO)_{10}$,⁷ in which one Os–Os edge is bridged by ligand X (X = three-electron donor) and a hydride ligand located below the osmium triangle. One hydride ligand bridging the Os(1)–Os(2) edge is positively defined by difference Fourier synthesis. The second hydride migrates to the terminal carbon atom

**Figure 1.** Molecular structure and atomic labeling scheme of the complex $Os_3(CO)_{10}(\mu-H)(\mu-\eta^3-C(=CHPh)C\equiv CW(O)_2-(C_5Me_5))$ (**1a**) with thermal ellipsoids shown at the 30% probability level.

C(14) of the incoming acetylide complex $(C_5Me_5)W(O)_2(C\equiv CC\equiv CPh)$, giving a novel $C\equiv CC=CHPh$ fragment which is confirmed by observation of a C–C double bond with distance 1.32(2) Å between C(13) and C(14) atoms. The resulting C_4 hydrocarbon fragment is coordinated to the triosmium framework in two axial sites; one is filled by carbon atom C(13), forming a regular σ -bond with distance 2.15(1) Å, whereas the second has a π -interaction from alkyne fragment C(11)–C(12) with $Os(1)\text{--}C(11) = 2.42(1)$ Å and $Os(1)\text{--}C(12) = 2.53(1)$ Å. The resulting C(13)–C(12)–C(11) unit is linear with bond angle $173(1)^\circ$ and is nearly parallel to the Os(1)–Os(1) edge. This novel $\mu-\eta^3$ -bonding mode is in contrast to that of a related hydrocarbonyl ligand in the allyl

(7) (a) Deeming, A. J. *Adv. Organomet. Chem.* **1986**, *26*, 1. (b) Smith, A. K. In *Comprehensive Organometallic Chemistry II*; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: Oxford, U.K., 1982; Vol. 7, Chapter 13.




Figure 2. Molecular structure and atomic labeling scheme of the complex $\text{Os}_3(\text{CO})_{10}(\mu\text{-H})(\mu\text{-}\eta^3\text{-C(=CHPh)C}\equiv\text{CW(O)}_2\text{-}(\text{C}_5\text{Me}_5))$ (**1b**) with thermal ellipsoids shown at the 30% probability level.


complex $\text{Os}_3(\text{CO})_{10}(\mu\text{-AuPEt}_3)(\mu\text{-}\eta^3\text{-C}_3\text{H}_5)$,⁸ in which the AuPEt_3 ligand is isolobal with a hydride, whereas the allyl ligand, adapting a $\mu_2\text{-}\eta^3$ -mode and serving as a three-electron donor, is found to coordinate to equatorial sites on the triosmium triangle, not the axial sites. Another notable feature is the pendant $(\text{C}_5\text{Me}_5)\text{W(O)}_2$ fragment that resides above equatorial CO ligands on the atom Os(2). The lengths of bonds between the W atom and terminal oxo ligands (W-O(11) , $\text{W-O(12)} = 1.71(1)$ Å) are similar to the average W=O distance observed in the anionic complex $[(\text{C}_5\text{Me}_5)\text{W(O)}_3][\text{PPN}]$ ⁹ and in the parent dioxo complex $(\text{C}_5\text{Me}_5)\text{W(O)}_2(\text{C}\equiv\text{CPh})$.¹⁰

The structure of **1a** evidently represents one isomeric structure detected in ¹H NMR spectra. The second structure (**1b**) is established with X-ray diffraction study of another single crystal obtained from a mixed solution of THF and methanol at -20 °C. Preliminary X-ray analysis indicates that the cell parameters of **1b** are not the same as those of **1a**. For example, the unit cell volume of **1b** is substantially greater, indicating a less efficient packing for molecules in the solid state. This observation prompted us to undertake the next single-crystal X-ray analysis.

As indicated in Figure 2, the structure of **1b** differs little from that of **1a**, consisting of three fundamental building blocks: a triangular $\text{Os}_3(\text{CO})_{10}$ unit, a bridging

hydride, and a coordinated $(\text{C}_5\text{Me}_5)\text{W(O)}_2(\text{C}\equiv\text{CC}=\text{CHPh})$ unit attached to two osmium atoms at adjacent axial sites. However, the $(\text{C}_5\text{Me}_5)\text{W(O)}_2$ fragment here undergoes a 180° rotation about the W-C bond, placing terminal oxo ligands in positions away from the unique $\text{Os}(\text{CO})_4$ fragment. Moreover, a hydride is found to bridge the Os(1)-Os(3) edge ($3.0741(6)$ Å), not the Os(1)-Os(2) edge as observed earlier, which is verified by a slight elongation of the Os-Os vector with respect to other Os-Os edges: $\text{Os(1)-Os(2)} = 2.845(6)$ Å and $\text{Os(2)-Os(3)} = 2.8996(7)$ Å. Thus, the structures of **1a** and **1b** are related by hydride migration between two adjacent Os-Os edges. By further extending this delineation, we assume that the last isomer (**1c**) would possess a structure with the hydride ligand associated with the third Os-Os edge (Scheme 1). This isomerization is consistent with a well-established tendency of hydride to migrate on a polymetallic cluster framework.¹¹

Thermolysis of 1. Decarbonylation of **1** is achieved on extensive thermolysis in CH_2Cl_2 at reflux (40 h). Separation of the reaction mixture by thin-layer chromatography yields two major bands. The first fast-eluting band is due to the yellow triosmium compound $\text{Os}_3(\text{CO})_{10}(\mu\text{-}\sigma\text{-}\eta^2\text{-C}\equiv\text{CCHCHPh})(\mu\text{-H})$ (**2**), but this com-



plex is relatively unstable, forming some intractable materials on dissolution in air at room temperature. As such, its recrystallization must be carried out at -20 °C and no attempt was made to obtain microanalysis data because of the poor thermal stability. In contrast, the second, red-orange band is more stable and consists of the complex $(\text{C}_5\text{Me}_5)\text{W(O)}_2\text{Os}_3(\text{CO})_9(\mu\text{-}\sigma\text{-}\eta^2\text{-C}\equiv\text{CCHCHPh})$ (**3**). Its formation is a result of competitive loss of CO and coordination of the $(\text{C}_5\text{Me}_5)\text{W(O)}_2$ unit to the osmium framework, a sequence which is typical for the related oxo carbonyl cluster compounds.¹²

Characterization of 2 and 3. For complex **2**, the FAB mass spectrum shows parent molecular ions assigned to $\text{C}_{20}\text{H}_{10}\text{H}_8\text{Os}_3$; the IR spectrum in the $\nu(\text{CO})$ region closely resembles that of the compound $\text{Os}_3(\text{CO})_{10}(\mu\text{-H})(\mu\text{-}\sigma\text{-}\eta^2\text{-C}\equiv\text{CPh})$.¹³ Furthermore, the ¹³C NMR spectrum was then recorded at 220 K, giving eight ¹³C NMR signals in the region δ 142.4–79.4, attributed to a $\mu\text{-C}\equiv\text{CCH}=\text{CHPh}$ fragment. Six sharp ¹³C NMR signals at δ 182.6, 182, 178.4, 175.3, 171.4, and 170.0 with intensity ratio 1:1:2:2:2:2 are also observed, show-

(8) Housecroft, C. E.; Johnson, B. F. G.; Lewis, J.; Lunniss, J. A.; Owen, S. M.; Raithby, P. R. *J. Organomet. Chem.* **1991**, *409*, 271.

(9) (a) Rau, M. S.; Kertz, C. M.; Mercado, L. A.; Geoffroy, G. L.; Rheingold, A. L. *J. Am. Chem. Soc.* **1991**, *113*, 7420. (b) Rau, M. S.; Kertz, C. M.; Geoffroy, G. L. *Organometallics* **1993**, *12*, 3447.

(10) Shiu, C.-W.; Su, C.-J.; Pin, C.-W.; Chi, Y.; Peng, S.-M.; Lee, G.-H. *J. Organomet. Chem.* **1997**, *545–546*, 151.

(11) (a) Band, E.; Muetterties, E. L. *Chem. Rev.* **1978**, *78*, 639. (b) Humphries, A. P.; Kaesz, H. D. *Prog. Inorg. Chem.* **1979**, *25*, 146. (c) Farrugia, L. J. *Adv. Organomet. Chem.* **1990**, *31*, 301.

(12) (a) Blenkinsop, P.; Carty, A. J.; Peng, S.-M.; Lee, G.-H.; Su, C.-J.; Shiu, C.-W.; Chi, Y. *Organometallics* **1997**, *16*, 519. (b) Chao, W.-J.; Chi, Y.; Chung, C.; Carty, A. J.; Delgado, E.; Peng, S.-M.; Lee, G.-H. *J. Organomet. Chem.* **1998**, *565*, 3.


(13) (a) Deeming, A. J.; Hasso, S.; Underhill, M. *J. Chem. Soc., Dalton Trans.* **1975**, 1614. (b) Koridze, A. A.; Kizas, O. A.; Petrovskii, P. V.; Kolobova, N. E.; Struchkov, Yu. T.; Yanovsky, A. I. *J. Organomet. Chem.* **1988**, *338*, 81.

ing the acetylide ligand undergoing a facile $\sigma \rightarrow \pi$, $\pi \rightarrow \sigma$ rearrangement, which produces a time-averaged plane of mirror symmetry bisecting the entire molecule. The CO chemical shifts observed and the respective dynamic behavior are all consistent with those of the complex $\text{Os}_3(\text{CO})_{10}(\mu\text{-H})(\mu\text{-}\sigma\text{-}\eta^2\text{-C}\equiv\text{CPh})$. In addition, the ^1H NMR spectrum exhibits a signal at $\delta -16.51$, characteristic of hydride bridging a Os–Os bond, a doublet at $\delta 7.71$, and a multiplet centered at $\delta 7.29$ due to the phenyl group. Notably, a pair of doublets at $\delta 6.67$ and 5.94 with the coupling constant $^3J_{\text{HH}} = 12.2$ Hz are observed, which pertain to olefinic protons that are *cis*-disposed.

On the other hand, the structure of **3** is assigned with reasonable certainty from FAB MS analysis and from the analysis of the IR $\nu(\text{CO})$ spectrum in the region $2088\text{--}1937\text{ cm}^{-1}$. Agreement between this $\nu(\text{CO})$ pattern with those observed for the structurally characterized complexes $\text{CpW}(\mu\text{-O})_2\text{Os}_3(\text{CO})_9(\mu\text{-H})^{14}$ and $(\text{C}_5\text{Me}_5)\text{W}(\mu\text{-O})_2\text{Os}_3(\text{CO})_9(\mu\text{-}\sigma\text{-}\eta^2\text{-C}\equiv\text{CPh})^{15}$ is excellent. Thus, the cluster core geometry can be definitely assigned to have the symmetrical butterfly skeletal arrangement of a $(\text{C}_5\text{Me}_5)\text{W}(\mu\text{-O})_2\text{Os}_3(\text{CO})_9$ unit. In agreement with this assignment, the ^{13}C NMR spectrum shows a time-averaged, five-line pattern for Os–CO signals at $\delta 187.2$, 185.4 , 177.6 , 172.4 , and 168.9 with intensity ratio 1:2:2:2:2. The ^1H NMR spectrum exhibits the expected signals for the phenyl group and two doublets at $\delta 6.47$ and 6.14 assigned to the $\text{CH}=\text{CHPh}$ unit. The coupling constant $^3J_{\text{HH}} = 11.9$ Hz is essentially identical with that observed in **2**, demonstrating the same *cis*-disposition of the $\text{CH}=\text{CHPh}$ unit.

Summary. Complex **1** forms three isomers in solution and maintains a constant ratio of isomers in solvents such as CD_2Cl_2 and $d_8\text{-THF}$. The structures of **1a** and **1b** represent a unique example of tetrametallic clusters identified by X-ray diffraction to have a hydride located on two distinct positions of the cluster framework. The reason for variation of the solid-state structure is unclear but may be associated with the polarity of solvents used for recrystallization. It is possible that the interaction of solvent molecules with oxo ligands perturbs packing of the $(\text{C}_5\text{Me}_5)\text{W}(\text{O})_2$ pendant within the molecule, inducing the hydride to move from one

Scheme 2



Os–Os edge to another during crystallization. A crystal polymorphism involving the hydride ligand has been documented for the cluster $\text{Ru}_6(\mu\text{-H})(\mu_6\text{-B})(\text{CO})_{17}$,¹⁶ in which the isomeric pairs differ in the rotameric conformation of $\text{Ru}(\text{CO})_3$ units and in the location of hydride. Other isomerism that is relevant to the present study is also observed in the compounds $\text{Ir}_6(\text{CO})_{16}$ and $\text{Ru}_6(\mu_6\text{-C})(\text{CO})_{17}$.¹⁷ As they possess no bridging hydride ligand, the isomerism arises entirely from the disposition of CO ligands with respect to the central cluster framework.

Furthermore, a mechanism leading to formation of **2** and **3** can be deduced according to their structures. Complex **3** is clearly produced in a sequence involving decarbonylation and migration of hydride to the $\mu\text{-}\eta^3\text{-C}\equiv\text{CC}=\text{CHPh}$ ligand. Formation of this *cis*- $\text{C}\equiv\text{CC}=\text{CHPh}$ ligand favors a pathway involving direct transfer of hydride to the $\text{C}\equiv\text{CC}=\text{CHPh}$ ligand of **1**; otherwise, production of a mixture of *trans* and *cis* isomers is envisaged. In contrast, the formation of **2** seems to involve removal of the $(\text{C}_5\text{Me}_5)\text{W}(\text{O})_2$ unit and concomitant addition of a hydrogen atom. This novel process is reminiscent of conversion from the heterometallic oxo carbonyl complex $\text{Ru}_3(\text{CO})_8(\mu_3\text{-NPh})(\mu\text{-}\eta^4\text{-CH}_2\text{CMeC}\equiv\text{CW}(\text{O})_2(\text{C}_5\text{Me}_5))$ to the triruthenium complex $\text{Ru}_3(\text{CO})_9\text{[CH}_2\text{CMeCC}(\text{CONHPh})\text{CO]}$ (Scheme 2).¹⁸ To understand how this reaction occurs, imagine that an H_2O molecule (or an equivalent) is introduced into the reaction system by chance or during workup. In effect, it reacts with the $(\text{C}_5\text{Me}_5)\text{W}(\text{O})_2$ fragment, giving off the unstable $[(\text{C}_5\text{Me}_5)\text{W}(\text{O})_2\text{OH}]$ and a proton. The coproduct $[(\text{C}_5\text{Me}_5)\text{W}(\text{O})_2\text{OH}]$ adsorbs on silica gel, and the proton remains on the cluster framework to afford the isolated homometallic complex **2**. This postulation has been substantiated by *in situ* ^1H NMR studies, which indicate almost no formation of **2** prior to chromatographic separation.

Acknowledgment. We thank the National Sciences Council of Taiwan (NSC Grant No. 87-2113-M007-047) for support of this work.

Supporting Information Available: Tables of non-essential crystal data, bond lengths, atomic coordinates, and corresponding anisotropic thermal parameters for structures **1a** and **1b**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM980957Z

(14) Chi, Y.; Hwang, L.-S.; Lee, G.-H.; Peng, S.-M. *J. Chem. Soc., Chem. Commun.* **1988**, 1456.

(15) (a) Shiu, C.-W.; Chi, Y.; Carty, A. J.; Peng, S.-M.; Lee, G.-H. *Organometallics* **1997**, *16*, 5368. (b) Shiu, C.-W.; Chi, Y.; Chung, C.; Peng, S.-M.; Lee, G.-H. *Organometallics* **1998**, *17*, 2970.

(16) (a) Draper, S. M.; Housecroft, C. E.; Keep, A. K.; Matthews, D. M.; Song, X.; Rheingold, A. L. *J. Organomet. Chem.* **1992**, *423*, 241. (b) Hong, F.; Coffy, T. L.; McCarthy, D. A.; Shore, S. G. *Inorg. Chem.* **1989**, *28*, 3284.

(17) (a) Garlaschelli, L.; Martinengo, S.; Bellon, P. L.; Demartin, F.; Manassero, M.; Chiang, M. Y.; Wei, C. Y.; Bau, R. *J. Am. Chem. Soc.* **1984**, *106*, 6664. (b) Braga, D.; Grepioni, F.; Dyson, P. J.; Johnson, B. F. G.; Frediani, P.; Bianchi, M.; Piacenti, F. *J. Chem. Soc., Dalton Trans.* **1992**, 2565.

(18) Pin, C.-W.; Chi, Y.; Chung, C.; Carty, A. J.; Peng, S.-M.; Lee, G.-H. *Organometallics* **1998**, *17*, 4161.