

Heterobimetallics of Nickel–Iron Dinitrosyl: Electronic Control by Chelate and Diatomic Ligands

Wen-Feng Liaw,^{*,†} Chao-Yi Chiang,[†] Gene-Hsiang Lee,[‡] Shie-Ming Peng,[‡]
Chia-Huei Lai,[§] and Marcetta Y. Darensbourg^{*,§}

Department of Chemistry, National Changhua University of Education, Changhua 50058, Taiwan,
Department of Chemistry & Instrumentation Center, National Taiwan University, Taipei 10764, Taiwan,
and Department of Chemistry, Texas A&M University, College Station, Texas 77843

Received June 2, 1999

Reaction of [PPN][Fe(NO)₂(SePh)₂] (**1**) with dimeric [Ni(μ -SCH₂CH₂SCH₂CH₂S)]₂ in the presence of additional NO₂⁻ produced the neutral heterobimetallic [(ON)Ni{(μ -SCH₂CH₂)₂S}Fe(NO)₂] complex (**2**). The X-ray crystal structures of **1** and **2** show distorted tetrahedral iron dinitrosyl groups, assigned according to the Feltham–Enemark notation as {Fe(NO)₂}⁹. The Fe–NO bonds are off linearity by an average of $\approx 10^\circ$ for compounds **1** and **2**, while a more linear Ni–NO coordination with a Ni–NO distance of 1.644(2) Å was found in **2**. The ν (NO) value of complex **2** is consistent with an assignment for {Ni(NO)}⁹ of Ni⁰(NO)⁺ as is known for analogous phosphine derivatives, P₃Ni⁰(NO)⁺. EPR signals of g values = 2.02–2.03 confirmed the existence of the odd electron in the chalcogenated {Fe(NO)₂}⁹ compounds. Two {Fe(NO)₂}¹⁰ complexes coordinated by the nickel(II) dithiolate, (bismercaptoethanediazacyclooctane)nickel(II), (**Ni-1**), (**Ni-1**)Fe(CO)(NO)₂ and (**Ni-1**)Fe(NO)₂, were prepared for comparison to the Ni⁰(NO)⁺ derivative and other monomeric and homodimetallic derivatives of the Fe(NO)₂ fragment. While the oxidation level of Fe(NO)₂ is the primary determinant of ν (NO) values, they are also highly sensitive to ancillary ligands and, thereby, the distal metal influence through the bridging thiolate donor.

Introduction

The selective synthesis of heterobimetallic complexes presents challenges, particularly in the area of thiolate-bridged M(μ -SR)₂M' systems where ideal precursors are apt to exist as stable polynuclear species and product speciation into homopoly-nuclear compounds is also prevalent. Interesting targets of this type are in biomimicry of the (CysS)₂Ni(μ -SCys)₂Fe(CN)₂(CO) active site of [NiFe]H₂ase, intriguing for the heterobimetallic site and also for the initial observation of nature's use of diatomic ligands in catalytically active site construction.¹ Since then, the Fe-only H₂ase has likewise been established to contain diatomic ligands in its binuclear active site.^{2,3} While the origin and role of the diatomic ligands are as much a question as is the requirement for two metals in the active site, two *different* metals in the case of [NiFe]H₂ase, advances in understanding are expected to result from the advantages offered by diatomic

ligands as spectroscopic probes if well-characterized model complexes are made available.

Recently, Osterloh and co-workers reported that a Ni(μ -SR)₂Fe(NO)₂ heterobimetallic was produced upon displacement of two carbonyls of Fe(CO)₂(NO)₂ by a square planar N₂S₂ complex of Ni(II).⁴ The resultant iron dinitrosyl unit maintained quite low ν (NO) values of <1700 cm⁻¹, as in its precursor Fe(CO)₂(NO)₂, where the iron nitrosyl unit has the formal electron assignment, {Fe(NO)₂}¹⁰. This formalism invokes the Enemark–Feltham notation which stresses the well-known covalency and delocalization in the electronically amorphous Fe(NO)₂ unit, without committing to a formal oxidation state on Fe, that is, assignments to Fe^{II}(NO⁻)₂, Fe⁰(NO)₂, Fe^{-II}(NO⁺)₂, or some mixture thereof.⁵ By application of our stepwise ligand exchange route, and with the use of a flexible S₃ precursor of Ni(II) in the presence of nitrite anion, we have prepared a second Ni(μ -SR)₂Fe(NO)₂ unit, complex **2** below, and report it herein as an example of formal oxidation state assignments of Ni⁰-{Fe(NO)₂}⁹.

Examples of nitric oxide coordination to iron and the spectroscopic signals of dinitrosyl iron complexes (DNIC) are of much interest, particularly in light of their role(s) in sulfur-rich protein uptake and degradation.⁶ Furthermore, DNICs have been suggested as possible forms for stabilization and transport of NO in biological systems.⁷ Thus, the precursor to the heterobimetallic complex **2**, (PhSe)₂Fe(NO)₂⁻, (complex **1**), was also isolated and characterized by X-ray crystallography and infrared spectroscopy. To address the heterometal influence on

* To whom correspondence should be addressed.

[†] National Changhua University of Education.

[‡] National Taiwan University.

[§] Texas A&M University.

- (1) (a) Volbeda, A.; Charon, M.-H.; Piras, C.; Hatchikian, E. C.; Frey, M.; Fontecilla-Camps, J. C. *Nature* **1995**, *373*, 580. (b) Volbeda, A.; Garcin, E.; Piras, C.; de Lacey, A. L.; Fernandez, V. M.; Hatchikian, E. C.; Frey, M.; Fontecilla-Camps, J. C. *J. Am. Chem. Soc.* **1996**, *118*, 12989. (c) de Lacey, A. L.; Hatchikian, E. C.; Volbeda, A.; Frey, M.; Fontecilla-Camps, J. C.; Fernandez, V. M. *J. Am. Chem. Soc.* **1997**, *119*, 7181. (d) Garcin, E.; Vernede, X.; Hatchikian, E. C.; Volbeda, A.; Frey, M.; Fontecilla-Camps, J. C. *Structure* **1999**, *7*, 557.
- (2) (a) Pierik, A. J.; Hulstein, M.; Hagen, W. R.; Albracht, S. P. J. *Eur. J. Biochem.* **1998**, *258*, 572. (b) Van Dam, P. J.; Reijerse, E. J.; Hagen, W. R. *Eur. J. Biochem.* **1997**, *248*, 355. (c) Van Der Spek, T. M.; Arendsen, A. F.; Happe, R. P.; Yun, S.; Bagley, K. A.; Stufkens, D. J.; Hagen, W. R.; Albracht, S. P. J. *Eur. J. Biochem.* **1996**, *237*, 629.
- (3) (a) Nicolet, Y.; Piras, C.; Legrand, P.; Hatchikian, E. C.; Fontecilla-Camp, J. C. *Structure* **1999**, *7*, 13. (b) Peters, J. W.; Lanzilotta, W. N.; Lemon, B. J.; Seefeldt, L. C. *Science* **1998**, *282*, 1853.

(4) Osterloh, F.; Saak, W.; Haase, D.; Pohl, S. *Chem. Commun.* **1997**, 979.

(5) Enemark, J. H.; Feltham, R. D. *Coord. Chem. Rev.* **1974**, *13*, 339.

(6) Foster, M. W.; Cowan, J. A. *J. Am. Chem. Soc.* **1997**, *121*, 44093.

(7) Vanin, A. F. *Biochem. (Moscow)* **1998**, *63*, 782.

Table 1. Summary of Crystallographic Data for [PPN][Fe(NO)₂(SePh)₂] (1) and [(ON)Ni(μ-S(CH₂)₂S(CH₂)₂S)Fe(NO)₂] (2)

	1	2
empirical formula	C ₄₈ H ₄₀ O ₂ N ₃ P ₂ FeSe ₂	C ₄ H ₈ O ₃ N ₃ S ₃ NiFe
fw (g/mol)	966.57	356.86
temp, K	298	298
wavelength, Å	0.7107	0.7107
cryst syst	monoclinic	triclinic
space group	C2/c	P1
unit cell dimensions	<i>a</i> = 16.769(4) Å <i>b</i> = 14.972(4) Å; <i>c</i> = 18.019(5) Å	<i>a</i> = 7.0616(12) Å; <i>b</i> = 7.4228(11) Å; <i>c</i> = 11.4142(12) Å
vol, Å ³	4450.2(20)	589.71(15)
Z	4	2
ρ _{calc} , Mg/m ³	1.443	2.010
<i>R</i> (<i>F</i>) [<i>I</i> > 2σ(<i>I</i>)] ^a %	3.7	2.6
w <i>R</i> (<i>F</i> ²) all data ^a %	3.2	3.8

^a Residuals: $R(F) = \sum |F_o - F_c| / \sum F_o$; $wR(F^2) = \{ \sum w(|F_o|^2 - |F_c|^2)|^2 / \sum w(F_o^2) \}^{1/2}$.

adjacent metal diatomics, analogues of the Osterloh et al. complex⁴ based on (bismercaptoethanediazacyclooctane)nickel(II), (bme-daco)Ni(II), **Ni-1**, as the metallathiolate ligand to {Fe(NO)₂(CO)} and {Fe(NO)₂} were prepared and their spectroscopic signals compared with a host of monomeric and homodimetallic derivatives of the Fe(NO)₂ fragment. The ability to reference the Fe(NO)₂ moiety within a consistent framework permits conclusions regarding the electronic character of the metal-modified ancillary ligands.

Experimental Section

General Methods and Materials. Solvents were reagent grade and were purged by nitrogen prior to use. All syntheses and product isolations were carried out under N₂ using standard Schlenk and glovebox techniques.

Physical Measurements. Infrared spectra were recorded on a Bio-Rad FTS-185 instrument using a 0.1-mm-spaced KBr sealed cell. Elemental analyses were carried out by a CHN analyzer (Heraeus). EPR spectra were measured on a Bruker ESP 300 spectrometer equipped with an Oxford ER910A cryostat at 100 K. The spin concentration for each EPR sample was determined by the comparison of the double integral of the entire EPR spectrum of the sample with that of a standard, that is, 1 mM Cu(II) in 10 mM EDTA.⁸ Electrochemical measurements were performed by a BAS-100A electroanalyzer utilizing glassy carbon working, Ag/AgNO₃ reference, and platinum auxiliary electrodes. Cyclic voltammograms were obtained from 2.5 mM analyte concentration in CH₂Cl₂ using 0.1 M [*n*-Bu₄N][PF₆] as a supporting electrolyte. Potentials were scaled to NHE using ferrocene as an internal standard.⁹

The X-ray crystal structures were solved at the Department of Chemistry and Instrumentation Center at National Taiwan University, Taipei, Taiwan. X-ray crystallographic data were obtained on a Nonius CAD 4 diffractometer with graphite-monochromated Mo Kα radiation employing the θ/2θ scan mode. A φ scan absorption correction was made. Structural determinations were made using the NRCC-SDP-VAX package of programs. Cell parameter and data collection summaries for [PPN][Fe(NO)₂(SePh)₂] (1) and [(ON)Ni(μ-S(CH₂)₂S(CH₂)₂S)Fe(NO)₂] (2) are given in Table 1.

Syntheses. Complexes [Ni(μ-S(CH₂)₂S(CH₂)₂S)]₂¹⁰ [PPN][[Fe(CO)₃-(SePh)₃],¹¹ Fe(NO)₂(CO)₂,¹² and (bme-daco)Ni(II)¹³ were synthesized by published procedures. (PPN = bis triphenylphosphineiminium cation).

Preparation of [PPN][Fe(NO)₂(SePh)₂] (1). The compound [PPN]-[[Fe(CO)₃(SePh)₃] (0.2 mmol, 0.115 g) was loaded into a 100-mL Schlenk flask with 0.028 g (0.4 mmol) of NaNO₂, and 30 mL of THF was added. The reaction mixture was stirred at 50 °C for 24 h, after which 15 mL of hexane was added. The resulting mixture was filtered to separate the yellow precipitate, NaSePh. Upon solvent removal in vacuo from the filtrate, [PPN][Fe(NO)₂(SePh)₂] was obtained as a dark red brown solid (0.075 g, 39%). Recrystallization from concentrated THF solution with ether diffusion gave dark red crystals used in the X-ray diffraction study. IR (ν_{NO}): 1735 s, 1694 vs cm⁻¹ (THF); 1741 s, 1697 vs cm⁻¹ (CH₂Cl₂). UV–vis absorption spectrum (THF) [λ_{max}, nm (ε, M⁻¹ cm⁻¹): 452(1544), 360(3943), 342(5095)]. Anal. Calcd for C₄₈H₄₀O₂N₃P₂FeSe₂: N, 1.28; C, 60.4; H, 4.15. Found: N, 1.34; C, 60.2; H, 4.13.

Preparation of [(ON)Ni(μ-S(CH₂)₂S(CH₂)₂S)Fe(NO)₂] (2). Into a 100-mL Schlenk flask were loaded the starting materials [PPN][Fe(NO)₂(SePh)₂] (0.40 mmol, 0.388 g), NaNO₂ (0.40 mmol, 0.028 g), and [Ni(μ-S(CH₂)₂S(CH₂)₂S)]₂ (0.20 mmol, 0.086 g). A 30-mL portion of CH₂Cl₂ was added to give a dark purple solution. The solution mixture was stirred at 50 °C overnight, after which 15 mL of hexane was added to precipitate the side products [PPN][Fe(NO)Cl₃] and NaSePh. The resulting mixture was filtered and solvent was removed from the filtrate under vacuum. One milliliter of CH₂Cl₂ was added to redissolve the products and hexane (12 mL) was added to obtain a dark green solid isolated by filtration (0.061 g, 43%). Recrystallization from ether gave green brown crystals which were soluble in hexane, THF, and methylene dichloride. IR (ν_{NO}): 1805 m, 1767 s, 1725 s cm⁻¹ (CH₂Cl₂); 1798 m, 1763 s, 1723 s cm⁻¹ (THF). Absorption spectrum (CH₂Cl₂) [λ_{max}, nm (ε, M⁻¹ cm⁻¹): 491(519), 385(919)]. Anal. Calcd for C₄H₈O₃N₃S₃NiFe: N, 11.78; C, 13.46; H, 2.26. Found: N, 11.68; C, 13.54; H, 2.21.

Preparation of (Ni-1)Fe(CO)(NO)₂ and (Ni-1)Fe(CO)₂(NO)₂. To a stirred solution of (bme-daco)Ni(II), **Ni-1** (0.044 g, 0.15 mmol), in acetonitrile (30 mL) was added Fe(CO)₂(NO)₂ (16 μL, 0.16 mmol). The reaction was monitored immediately by IR to confirm the loss of bands for the starting material with formation of (Ni-1)Fe(CO)(NO)₂, THF solution spectra (cm⁻¹): (ν_{CO}) 2006 br; (ν_{NO}): 1733 s, 1691 vs. After 1.5 h of stirring, the resulting red-brown solution had two ν_{NO} bands at 1677 (ms) and 1630 (s) cm⁻¹, as expected for (Ni-1)Fe(NO)₂, based on the spectroscopy of the Osterloh et al. complex.⁴ Both compounds are thermally unstable and decomposed under vacuum; they dissolve in polar solvents such as THF and acetonitrile.

Results and Discussion

Syntheses and Molecular Structures. Presented in Scheme 1 is the reaction which led to dark red, crystalline [PPN⁺][Fe(NO)₂(SePh)₂⁻]. The molecular structure of the anion, Figure 1, is that of a distorted tetrahedron, elongated along the Se–Fe–Se and N–Fe–N edges. The nitrosyls are slightly bent (∠Fe–N–O = 169°) and flared toward each other.⁶ The Fe–SePh average distance of 2.395(1) Å is significantly shorter than that of the six-coordinate fac-Fe(CO)₃(SePh)₃⁻ (Fe–Se average = 2.459(2) Å)^{11a} and tetrahedral [Fe(SePh)₄]²⁻ (Fe–Se average = 2.460(12) Å).¹⁴ This is rationalized by the higher coordination

(8) Orme-Johnson, N. R.; Orme-Johnson, W. H. *Methods Enzymol.* **1978**, 52, 252.
(9) Gagne, R. R.; Koval, C. A.; Lisensky, G. C. *Inorg. Chem.* **1980**, 19, 2854.

(10) (a) Harley-Mason, J. J. *Chem. Soc.* **1952**, 146. (b) Baker, D. J.; Goodall, D. C.; Moss, D. S. *Chem. Commun.* **1969**, 325.
(11) (a) Liaw, W.-F.; Lai, C.-H.; Chiang, M.-H.; Lee, C.-K.; Lee, G.-H.; Peng, S.-M. *J. Chem. Soc., Dalton Trans.* **1993**, 2421. (b) Liaw, W.-F.; Chiang, M.-H.; Liu, C.-J.; Harn, P.-J.; Liu, L.-K. *Inorg. Chem.* **1993**, 32, 1536. (c) Liaw, W.-F.; Chen, C.-H.; Lee, C.-M.; Lin, G.-Y.; Ching, C.-Y.; Lee, G.-H.; Peng, S.-M. *J. Chem. Soc., Dalton Trans.* **1998**, 353.
(12) Hieber, W.; Beutner, H. Z. *Anorg. Allg. Chem.* **1963**, 320, 101.
(13) Mills, D. K.; Reibenspies, J. H.; Darensbourg, M. Y. *Inorg. Chem.* **1990**, 29, 4364.
(14) McConnachie, J. M.; Ibers, J. A. *Inorg. Chem.* **1991**, 30, 1770.





Figure 1. Labeling scheme for $[\text{Fe}(\text{NO})_2(\text{SePh})_2]^-$ (**1**) with thermal ellipsoids drawn at the 50% probability level. Fe—Se, 2.395(1); Fe—N(1), 1.669(4); N(1)—O(1), 1.162(5). Se—Fe—Se(a), 114.08(5); Se—Fe—N(1), 107.88(13); Se—Fe—N(1a), 105.06(13); N(1)—Fe—N(1a), 117.20(19); Fe—N(1)—O(1), 169.2(4).

Scheme 1



number of the former and higher anionic charge of the latter. Structural and spectroscopic features are the same for **1** and $[\text{Fe}(\text{NO})_2(\text{SPh})_2]^-$.¹⁵

Subsequent reaction of the $[\text{Fe}(\text{NO})_2(\text{SePh})_2]^-$ anion with dimeric $[\text{Ni}(\mu\text{-SCH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{S})_2]$ in the presence of additional NO_2^- produced the neutral heterobimetallic $[(\text{ON})\text{Ni}\{\mu\text{-SCH}_2\text{CH}_2\text{SCH}_2\text{S}\}\text{Fe}(\text{NO})_2]$ complex **2**, as a thermally stable, dark green-brown solid isolated in $\approx 70\%$ yield. Complex **2** is soluble in hexane and THF and forms slightly air-sensitive solutions in CH_2Cl_2 . Crystals suitable for X-ray crystallography were grown from concentrated hexane solution and were found to be in the triclinic $\bar{P}1$ space group.

Figure 2 displays a thermal ellipsoid plot of the neutral Ni—Fe bimetallic complex **2** and selected distances and angles are given in the caption. The constraints of the bismercaptoethylenesulfide ligand generates $\approx 90^\circ$ $\text{S}_{\text{thioether}}\text{—Ni—S}_{\text{thiolate}}$ angles,¹⁶ enforcing a severe distortion from a tetrahedral at the four-coordinate nickel site. Nevertheless, the NiS_2Fe core is minimally hinged (the angle between NiS_2 and FeS_2 planes = 162.8°).

All $\text{M—S}_{\text{br thiolate}}$ distances of complex **2** are substantially the same, 2.29 Å, while the $\text{Ni—S}_{\text{thioether}}$ distance is 0.02-Å shorter. The average Fe—S—Ni angle of 75.2° and the Fe—Ni distance of 2.8001(6) Å are similar to those of complex **4**, $76.11(4)^\circ$ and 2.797(1) Å, respectively. (A stick drawing of complex **4** is found in Table 2.) The large $\angle \text{S}_{\text{br}}\text{—Ni—S}_{\text{br}}$ of $103.94(3)^\circ$ and $\angle \text{S—Fe—S}$ of $103.73(3)^\circ$ found in complex **2** however dramati-




Figure 2. Labeling scheme for $[(\text{ON})\text{Ni}(\mu\text{-S}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{S})\text{Fe}(\text{NO})_2]$ (**2**) with thermal ellipsoids drawn at the 50% probability level. Fe—S(1), 2.2907(9); Fe—S(3), 2.3039(9); Fe—N(1), 1.670(2); Fe—N(2), 1.674(3); N(1)—O(1), 1.163(3); N(2)—O(2), 1.159(3); Ni—S(1), 2.2947(9); Ni—S(2), 2.2701(9); Ni—S(3), 2.2931(9); Ni—N(3), 1.644(2); N(3)—O(3), 1.119(3); Ni···Fe, 2.8001(6). S(1)—Fe—S(3), 103.73(3); S(1)—Fe—N(1), 114.88(9); S(1)—Fe—N(2), 106.02(9); N(1)—Fe—N(2), 115.06(12); S(1)—Ni—S(2), 91.67(3); S(1)—Ni—S(3), 103.94(3); S(1)—Ni—N(3), 120.53(10); S(2)—Ni—S(3), 92.31(4); S(2)—Ni—N(3), 126.69(9); S(3)—Ni—N(3), 115.83(10); Fe—S(1)—Ni, 75.28(3); Fe—S(3)—Ni, 75.05(3); Fe—N(1)—O(1), 166.21(23); Fe—N(2)—O(2), 167.3(3); Ni—N(3)—O(3), 175.1(3).

cally differ from the analogous features in complex **4**, $81.46(6)^\circ$ and $74.10(6)^\circ$, respectively. Further appropriate comparisons lie in the Fe—N and N—O parameters; the Fe—N bond distances (average 1.67 Å) are not disturbed by the change of donor sets in complexes **1** versus **2** which are very slightly longer than the 1.65-Å average seen for **4**. The Fe—NO bonds are off linearity by an average of $\approx 10^\circ$ and, as previously noted for complex **1**, they bend toward each other.

The Ni—NO distance (1.644(2) Å) in complex **2** is comparable with the terminal Ni—NO of 1.653(4) Å in $[\text{Ni}(\text{NO})(\text{PTA})_3]^+$ (PTA = 1,3,5-triaza-7-phosphaadamantane);¹⁷ the nitrosyl ligand is coordinated in a nearly linear mode with the $\angle \text{Ni—N(3)—O(3)} = 175.1(3)^\circ$. Together with similar $\nu(\text{NO})$ values found for complexes **2** and $[\text{Ni}(\text{NO})(\text{PTA})_3]^+$, vide infra, we conclude the site is best described as NO^+ coordinated to nickel(0).

Spectroscopic and Electrochemical Measurements. Complexes **1**, **2**, and **3**, the $\{\text{Fe}(\text{NO})_2\}^0$ species, in Table 2, have an odd number of electrons and were found to have similar EPR properties: a single isotropic signal with no hyperfine coupling to the NO ligands. The g values of 2.021, 2.021, and 2.028, for **1**, **2**, and **3**, respectively, are consistent with halogenonitrosyl iron complexes, $[\text{X}_2\text{Fe}(\text{NO})_2]^-$ (X = Cl, Br, I) which also have tetrahedral geometry and an electron assignment of $\{\text{Fe}(\text{NO})_2\}^0$ with $S = 1/2$.¹⁸ Weaker EPR signals, assumed to be due to impurities, were observed for solutions of complexes **4**, **5**, and **6** (spin quantitations of 0.07–0.10 spin per molecule as compared to ≈ 1.0 for **1**, **2**, and **3**).

On treatment of the complex **2** with 1 equiv of Ce^{IV} (as cerium ammonium nitrate), the EPR signal at $g = 2.021$ was lost. Addition of the weak reductant, Cp_2Co , produced no change in the EPR spectrum. However, addition of the LiHBET_3 reagent to the EPR sample tube produced an axial signal of $g = 2.602$ and 2.018. This product has not been identified.












(15) Strasdeit, H.; Krebs, B.; Henkel G. *Z. Naturforsch.* **1986**, *41b*, 1357.

(16) Kaasjager, V. E.; Henderson, R. K.; Bouwman, E.; Lutz, M.; Spek, A. L.; Reedijk, J. *Angew. Chem., Int. Ed.* **1998**, *37*, 1668.

(17) PTA = phosphotriazaadamantane, a neutral P-donor ligand. Darensbourg, D. J.; Decuir, T. J.; Stafford, N. W.; Robertson, J. B.; Draper, J. D.; Reibenspies, J. H.; Kathó, A.; Joó, F. *Inorg. Chem.* **1997**, *36*, 4218.

(18) Connelly, N. G.; Gardner, C. J. *Chem. Soc., Dalton Trans.* **1976**, 1525.

Table 2. Selected Structure and Infrared Data for Dichalcogenide Fe(NO)₂ Derivatives {Fe(NO)₂}^{9,10} Based on Enemark–Feltham Notation⁶

Compound	Compound	$\nu(\text{NO}), \text{cm}^{-1}$ [$\nu(\text{CO}), \text{cm}^{-1}$]	$\angle\text{N-Fe-N, deg}$ [$\angle\text{L-Fe-L}$]	$\angle\text{Fe-N-O, deg}$
1	 PhSeO ₂ (Fe(NO) ₂) ⁹ PhSe	1697, 1741 ^a	117.20(19) [114.08(5)]	169.2(4)
1a	 PhS ₂ (Fe(NO) ₂) ⁹ PhS	1709, 1744 ^b	117.36(9) [109.73(2)]	168.5(2) 169.4(2)
2	 (2.8001(6) Å)	1725, 1767 ^a	115.06(12) [103.73(3)]	166.21(23) 167.3(3)
3 ²⁰		1695, 1740 ^c	118.09(12) [109.15(3)]	169.8(3) 169.8(3)
4 ⁴	 (2.797(1) Å)	1624, 1663 ^c 1630, 1675 ^d	118.2(3) [74.10(6)]	167.0(5) 174.6(5)
5		1691, 1733 ^e [2006]		
6		1630, 1677 ^d		
7 ²¹		1756, 1810 ^f [2034, 2087]	114(6) [110(3)]	ca. 180
8 ²²		1722, 1764 ^g [2009]	114.4(4) ^h [103.9(4)]	177.9(6) ^h
9 ²²		1678, 1724 ^g	123.8(4) [111.9(1)]	178.2(7)
10 ²⁷	 (2.6166 (4) Å)	1710, 1760 ^c	120.8(1) [101.1(3)]	173.1(3) 171.4(3)

^a Dichloromethane. ^b CsI. ^c KBr. ^d Acetonitrile. ^e THF. ^f Cyclohexane. ^g Tetrachloroethylene. ^h Shown are average values.

As noted in Table 2, the CH₂Cl₂ solution spectrum of the neutral heterobimetallic compound **2** shows three infrared bands in the ν_{NO} region. The highest frequency band at 1805 cm⁻¹ is identical to the ν_{NO} band of the tetracoordinate, pseudotetrahedral Ni(0) nitrosyl complex [Ni(NO)(PTA)₃]⁺ ($\nu_{\text{NO}} = 1805$ cm⁻¹ in CH₂Cl₂); thus, we can project an electron donor ability of the (μ -((SCH₂CH₂)₂S)Fe(NO)₂) unit to [NiNO]⁺ to be the same as that of the P-donor ligands, assuming an oxidation state of zero for nickel and the presence of NO as NO⁺.¹⁷ That is, the three S donors comprised of a {Fe(NO)₂}⁹-modified dithiolate and the one thioether is equivalent to three P donors of the phosphotriazaadamantane ligand, PTA.

The lower energy NO bands of **2** are shifted by $\approx +30$ cm⁻¹ from those of the anionic monomeric precursor, Fe(NO)₂(SePh)₂⁻. Known to serve as a reporter of electron density at iron similarly to CO,¹⁹ the shifts in NO frequencies reflect a variation in negative charge on the Fe(NO)₂ unit as follows: **4** < **3** ~ **1** <

(19) Horrocks, W. D.; Taylor, R. C. *Inorg. Chem.* **1963**, *2*, 723.




Figure 3. Cyclic voltammogram of a 2.5-mV solution of complex **2** in 0.1 M TBAHFP/CH₂Cl₂ with a glassy carbon working electrode at a scan rate of 200 mV/s.

2.^{4,20} Thus, the charge separation indicated by the formal oxidation states of the heterobimetallic complexes **2** (Ni⁰{Fe(NO)₂}⁹) and **4** (Ni^{II}{Fe(NO)₂}¹⁰) is reflected in the ν_{NO} values. The similarity of ν_{NO} values in the Fe(NO)₂ units of complexes **1** and **3**, despite the negative charge on the former, is rationalized by the firm localization of H⁺ on the two nitrogen bases, as indicated in the structure drawn for **3**. The higher ν_{NO} values of **2** versus **3** would then imply that the Ni⁰(NO⁺) unit substantially neutralizes the cis thiolate negative charges, and the [(ON)Ni⁰S₃]⁻ unit serves as a bidentate donor ligand of -1 charge to the {Fe(NO)₂}⁹ unit.

Figure 3 displays the cyclic voltammogram (CV) of complex **2** with its two quasi-reversible redox events centered at +690 and -970 mV. Because the CV of [PPN][PhSe)₂{Fe(NO)₂}⁹] has a reduction feature at -1160 mV, the -970-mV event of complex **2** is assumed to be due to the {Fe(NO)₂}⁹{Fe(NO)₂}¹⁰ couple. With this assignment, we conclude that the anionic (ON)Ni(0)-dithiolate unit donates slightly less electron density to the {Fe(NO)₂}⁹ unit as compared to the two phenylselenide donors and stabilizes this event by 190 mV. This result is consistent with the $\nu(\text{NO})$ values in the IR spectra as discussed above. The quasi-reversible anodic event at +687 mV could be assigned to the {Fe(NO)₂}⁹{Fe(NO)₂}⁸ couple, consistent with the observed loss of the EPR signal of chemically oxidized **2**. Alternatively, assignment to a nickel-based oxidation, the Ni^{0/1} couple, would have to invoke coupling of the {Fe(NO)₂}⁹ and the Ni^I (*d⁹*) centers to account for the loss of paramagnetism and the EPR signal.

Further Comments

The assignment of formal oxidation levels to the Fe(NO)₂ units compared in this study was done in an effort to describe the electron-donating abilities of the ancillary ligands within a similar framework. With this approach, we are able to reference the nickel(0)dithiolate, compound **2**, and nickel(II)dithiolate ligands, compounds **4**, **5**, and **6**, with respect to mononuclear complexes with rather unambiguous additional ligands to Fe(NO)₂.^{21,22} The following conclusions are made:

- (20) Baltusis, L. M.; Karlin, K. D.; Rabinowitz, H. N.; Dewan, J. C.; Lippard, S. J. *Inorg. Chem.* **1980**, *19*, 2627.
- (21) (a) Brockway, L. O.; Anderson, J. S. *Trans. Faraday Soc.* **1937**, *38*, 1233. (b) Hedberg, L.; Hedberg, K.; Satija, S. K. Swanson, B. I. *Inorg. Chem.* **1985**, *24*, 2766.
- (22) Albano, V. G.; Araneo, A.; Bellon, P. L.; Ciani, G.; Manassero, M. J. *Organomet. Chem.* **1974**, *67*, 413.

(1) Two oxidation levels, $\{\text{Fe}(\text{NO})_2\}^9$ and $\{\text{Fe}(\text{NO})_2\}^{10}$ are observed in the heterobimetallic molecular units examined in this work. The paradigms for the former are $(\text{PhS})_2\text{Fe}(\text{NO})_2^-$ and $(\text{PhSe})_2\text{Fe}(\text{NO})_2^-$ while that of the latter is $(\text{CO})_2\text{Fe}(\text{NO})_2$. The π -donor ability of thiolate (or selenoate) stabilizes the former $\{\text{Fe}(\text{NO})_2\}^9$ or oxidized form, while the π -accepting ability of CO stabilizes the latter, reduced form.


(2) The oxidation level of $\text{Fe}(\text{NO})_2$ is the primary determinant of the $\nu(\text{NO})$ position. Within a series of the same oxidation level, the $\nu(\text{NO})$ reporter shows a sensitivity to ancillary ligands to $\{\text{Fe}(\text{NO})_2\}$ similarly to $\nu(\text{CO})$ as described in classic studies such as that of Horrocks and Taylor in a homologous series of $\text{Co}(\text{NO})(\text{CO})(\text{PR}_3)$ complexes.¹⁹ In an earlier study, we referenced the nickel(II) dithiolate, (bme-daco)Ni as a S-donor ligand to $\text{Fe}^0(\text{CO})_4$ relative to $(\text{Ph}_3\text{P})\text{Fe}(\text{CO})_4$ using the noncontroversial CO ligand as a spectroscopic probe.²³ There, we found the monodentate metalthiolate ligand to be a better donor, shifting the $\nu(\text{CO})$ values negatively on average by $\approx 30 \text{ cm}^{-1}$ from $(\text{Ph}_3\text{P})\text{Fe}(\text{CO})_4$.

(3) The formation and the $\nu(\text{NO})$ values of the NiFe bimetallic compound **2** is consistent with a dichalcogenide derivative of $\{\text{Fe}(\text{NO})_2\}^9$ that is partially charge-neutralized by interaction of a single cationic charge, that is, $[\text{Ni}(\text{NO})]^+$. This gives rise to the highest $\nu(\text{NO})$ frequency of the $\{\text{Fe}(\text{NO})_2\}^9$ series. Alternatively, the heterometallic may be described as a $[\text{S}'\text{S}'\text{Ni}(\text{NO}^+)]^-$ moiety bound by an interacting $\{\text{Fe}(\text{NO})_2\}^9$ unit which effectively softens the thiolate donors of $\text{S}(\text{CH}_2\text{CH}_2\text{S})_2$.

(4) The $\nu(\text{NO})$ positions of compounds **4**, **5**, and **6**, those containing the $\{\text{Fe}(\text{NO})_2\}^{10}$ moiety, Table 2, are consistent with those of compounds **7**, **8**, and **9** given the established donor ability of NiSR versus PPh_3 discussed above. That is, the nickel thiolate is a better donor than phosphine, producing the lowest observed $\nu(\text{NO})$ in the entire series for compounds **4** and **6**.

Thus, for the purpose of interpreting the IR spectra, the $\text{Ni}^{\text{II}}(\mu\text{-SR})_2\text{Fe}(\text{NO})_2$ and the $\text{Ni}^0(\mu\text{-SR})_2\text{Fe}(\text{NO})_2$ complexes may be viewed as mononuclear $\text{Fe}(\text{NO})_2$ centers with ligands (i.e., thiolates) modified by distal metal interactions.²⁴ The oxidation level of the $\text{Fe}(\text{NO})_2$ unit may be rationalized according to the ability of the chalcogenide (or metal-modified chalcogenide) ligand to stabilize the oxidized, $\{\text{Fe}(\text{NO})_2\}^9$ or reduced, $\{\text{Fe}(\text{NO})_2\}^{10}$ forms. A 2+ charge on the distal metal, as in the Ni^{II} -bound dichalcogenide, creates neutral sulfur donor sites and stabilizes the electron-rich or reduced form, as does CO or PR_3 . The $[\text{Ni}^0\text{NO}^+]$ unit effectively generates a monoanionic dithiolate ligand donor set, which stabilizes the oxidized form. Interestingly, both have the same Ni–Fe distances of 2.8 Å, but distinctly different hinge angles of the $\text{Ni}(\mu\text{-SR})_2\text{Fe}$ unit, 104.6° for complex **4** and 162.8° for compound **2**. The maintenance of the same Ni–Fe distance is possible because the $\angle\text{S}-\text{Fe}-\text{S}$ and $\angle\text{S}-\text{Ni}-\text{S}$ in the core NiS_2Fe unit open by $\approx 30^\circ$ on going from the butterfly structure of **4** to the roughly planar structure of **2**, with a concomitant shift of the $\text{Fe}(\text{NO})_2$ unit relative to the NiS_2 plane.

Such geometric differences as determined by the nickel oxidation state in **4** versus that in **2** demand ligands capable of



matching the geometrical requirement of the $\text{Ni}(\mu\text{-SR})_2\text{Fe}$ unit as well as maintaining the preferred coordination number of the nickel. One would expect the protein's cysteinyl sulfurs which bridge the Ni–Fe active site of $[\text{NiFe}]_2\text{H}_2\text{ases}$ to be capable of such flexibility. *In fact, as observed in the protein crystal structures, the irregular orientation of the two terminal and two bridging CysS about nickel (pseudo-tetrahedral, or distorted, open-site square pyramid) could indicate a balance of geometrical requirements for different redox levels of nickel in different states of the enzyme.* In small molecule model chemistry, two different ligands are required to mimic the effect. [It should be noted that the usefulness of the $(\mu\text{-SCH}_2\text{CH}_2)_2\text{S}$ ligand, was previously recognized in the preparation of a binuclear structural model for Fe only hydrogenase.^{16]}

Electron coupling through the 2.8-Å Ni–Fe distances found in the heterometallics with the $\text{Fe}(\text{NO})_2$ unit is expected, but not yet established. The Ni–Fe distances in $\text{Ni}^{\text{II}}\text{-Fe}^{\text{II}}$ heterometallics that do not contain the $\text{Fe}(\text{NO})_2$ unit are $\approx 3.0 \text{ \AA}$ or larger.^{22,25,26} Interestingly, the natural heterobimetallic found in the $[\text{NiFe}]_2\text{H}_2\text{ase}$ enzyme active site has a Ni–Fe distance of 2.9 Å.¹ The much shorter distance of 2.6 Å in compound **10**,²⁷ listed in Table 2 as $\text{Ni}^{\text{II}}\{\text{Fe}(\text{NO})_2\}^{10}$, probably implies a Ni–Fe bond character and an assignment of $\text{Ni}^{\text{I}}\{\text{Fe}(\text{NO})_2\}^9$ would be equally, if not more, realistic.

Finally, it is noted that the usual exchange of two NO donor ligands for three two-electron donors such as CO and CN^- produces a fragment, $\text{Fe}(\text{NO})_2$, that could be interchangeable with isolobal $\text{Fe}(\text{CO})(\text{CN})_2$, recently established as important in the hydrogenases. Because all three diatomic ligands have infrared stretches in the same region, isotopic labeling should be a requirement for confidence in spectroscopic assignments and vibrational studies.

Acknowledgment. The authors acknowledge the financial support of the National Science Council (Taiwan to W.F.L. and S.M.P.) and the National Science Foundation (U.S.A., CHE-9812355 to M.Y.D.). We are grateful for the assistance of Huay-Keng Loke in the EPR measurements, to Jason Smee for information retrieval and computer display, and to Dr. Frank Osterloh for his helpful critique of the manuscript.

Supporting Information Available: Tables of crystal data and experimental conditions for the X-ray studies, atomic coordinates and B_{eq} values, complete listings of bond lengths and bond angles, and anisotropic temperature factors for $[\text{PPN}][\text{Fe}(\text{NO})_2(\text{SePh})_2]$ and $[(\text{ON})\text{-Ni}(\mu\text{-S}(\text{CH}_2)_2\text{SCH}_2)_2\text{S}]\text{Fe}(\text{NO})_2$. This material is available free of charge via the Internet at <http://pubs.acs.org>.

IC990631Y

(23) Lai, C.-H.; Reibenspies, J. H.; Darensbourg, M. Y. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 2390.

(24) Musie, G.; Farmer, P. J.; Tuntulani, T.; Reibenspies, J. H.; Darensbourg, M. Y. *Inorg. Chem.* **1996**, *35*, 2176.

(25) Mills, D. K.; Hsiao, Y. M.; Farmer, P. J.; Atnip, E. V.; Reibenspies, J. H.; Darensbourg, M. Y. *J. Am. Chem. Soc.* **1991**, *113*, 1421.

(26) Colpas, G. J.; Day, R. O.; Maroney, M. J. *Inorg. Chem.* **1992**, *31*, 5053.

(27) Chau, C. N.; Wojcicki, A.; Calligaris, M.; Nardin, G. *Inorg. Chem. Acta* **1990**, *168*, 105.