

Synthesis, spectra and crystal structure of (*E*)-(CO)₂(NO)Cr[(η^5 -C₅H₄)-CH=CH(η^5 -C₅H₄)]Cr(CO)₂(NO)

Yu-Pin Wang^{a,*}, Xen-Hum Lui^a, Bi-Son Lin^a, Wei-Der Tang^a, Tso-Shen Lin^a,
Jen-Hai Liaw^a, Yu Wang^b, Yi-Hung Liu^b

^a Department of Chemistry, Tunghai University, Taichung, Taiwan, ROC

^b Department of Chemistry, National Taiwan University, Taipei, Taiwan, ROC

Received 14 April 1998; received in revised form 19 September 1998

Abstract

Compounds 1,2-bis[(η^5 -cyclopentadienyl)dicarbonylnitrosylchromium]ethene (**7**) (hereafter called 1,2-dicynichrodenylethene) and 1,2-diferrocenylethene (**10**) were prepared from formylcynichrodene (**3**) and formylferrocene (**9**), respectively, via the McMurry's low-valent titanium coupling method. Compounds (η^5 -vinylcyclopentadienyl)dicarbonylnitrosylchromium (**6**) and vinylferrocene (**11**) were obtained by the dehydration of the corresponding alcohols. The structure of **7** was solved by an X-ray diffraction study: space group, $P2_1/c$; monoclinic; $a = 6.379(5)$, $b = 11.295(3)$ and $c = 11.9352(24)$; $Z = 2$. It turns out that compound **7** adopts a *transoid* conformation at the ethenylene bridge and the two cyclopentadienyl rings are coplanar. The nitrosyl group in each cynichrodenyl moiety of **7** is located at the side towards the corresponding ethenylene carbon atom with a twist angle of 46.5° . The chemical shifts of H(2)–H(5) protons and C(2)–C(5) carbon atoms of a series of vinyl derivatives of compounds bearing cyclopentadienyl rings have been assigned using two-dimensional HetCOR-NMR spectroscopy. For the derivatives of cynichrodene (**1**) and ferrocene, it was found that the shielding of C(2,5) and C(3,4) carbon atoms is parallel to the shielding of the *ortho*- and *para*-carbon atoms of benzene derivatives. The electron density distribution in the cyclopentadienyl ring is discussed on the basis of ¹³C-NMR data. Surprisingly, the vinyl group donates electron density to the adjacent cynichrodene moieties rather than withdraws from them. © 1999 Elsevier Science S.A. All rights reserved.

Keywords: Chromium; Synthesis; Spectra; Crystal structure

1. Introduction

Earlier [1,2], we reported the unequivocal assignments of C(2,5) and C(3,4) on the Cp ring of cynichrodene (**1**) derivatives bearing electron-donating or electron-withdrawing substituent in ¹³C-NMR spectra. For derivatives with electron-donating substituents, an analogy was observed between the shielding of C(2,5) and C(3,4) carbon atoms of cynichrodene **1** derivatives and ferrocene derivatives and that of the *ortho*-

and *para*-carbon atoms of benzene derivatives. For derivatives bearing electron-withdrawing substituents, the opposite correlation on the assignments was observed between cynichrodene derivatives and the derivatives of ferrocene and benzene. The high-field and low-field chemical shifts are assigned to C(3,4) and C(2,5), respectively, in the case of cynichrodene derivatives bearing electron-withdrawing groups, while the opposite assignment was made for the ferrocene derivatives. This amazing finding prompted us to study **6–7** and **10–11**, compounds containing vinyl moiety, a group capable of being either electron-donating or electron-withdrawing.

Several methods have been employed to couple the

* Corresponding author. Fax: +886-4-359-96851.

carbonyl groups to yield orefins [3,4]. However, the applications to organometallic compounds are limited [5]. Herein, we report the preparations of **7** and **10** via the McMurry's low valent titanium method and the crystal structure of (*E*)-(CO)₂(NO)Cr(Cp-CH=CHCp)Cr(NO)(CO)₂ (**7**). Spectral comparison between benzene, ferrocene and cynichrodene derivatives bearing corresponding substituents is also included.

2. Experimental details

All the syntheses were carried out under nitrogen by the use of Schlenk techniques. Traces of oxygen in the nitrogen were removed with BASF catalyst and the deoxygenated nitrogen was dried over molecular sieves (3 Å) and P₂O₅. Hexane, pentane, benzene, and dichloromethane were dried over calcium hydride and freshly distilled under nitrogen. Diethyl ether was dried over sodium and re-distilled under nitrogen from sodium-benzophenone ketyl. All the other solvents were used as commercially obtained.

Column chromatography was carried out under nitrogen with Merck Kiesel-gel 60. The silica gel was heated with a heat gun during mixing in a rotary evaporator attached to a vacuum pump for 2 h to remove water and oxygen. The silica gel was then stored under nitrogen until use. Compounds **3** and **9** were prepared according to the literature procedures [6,7].

¹H- and ¹³C-NMR were acquired on a Varian Unity-300 spectrometer. Chemical shifts were referenced to tetramethylsilane. IR spectra were recorded with a Perkin-Elmer Fourier transform IR 1725X spectrophotometer. Microanalyses were carried out by the Microanalytical Laboratory of the National Chung Hsing University.

2.1. Preparation of (η^5 -vinylecyclopentadienyl)dicarbonylnitrosylchromium (**6**)

This compound was prepared according to Mintz and Rausch [8] with minor modifications. [η^5 -(1-Hydroxyethyl)cyclopentadienyl]dicarbonylnitrosyl-chromium (**8**) (0.54 g, 2.2 mmol), 5 mg of hydroquinone, and 0.05 g of *p*-toluenesulfonic acid were dissolved in 50 ml of benzene. The mixture was refluxed for 1.25 h, then 100 ml of dichloromethane was added. The solution was washed with distilled water and then dried with anhydrous magnesium sulfate. After the solution was filtered and concentrated, 0.42g of a residue of **6** (84%) was obtained. An analytical sample was prepared from centrifugal TLC (hexane:benzene (1:1)).

Proton-NMR (CDCl₃): δ (relative intensity, multi-

plicity, assignment): 5.01 (2H, t, Cp H(3,4)); 5.15 (1H, dd, H_B, $J_{AB} = 3.3$, $J_{BX} = 11$); 5.20 (2H, t, Cp H(2,5)); 5.44 (1H, dd, H_A, $J_{AX} = 17.7$); 6.22 (1H, dd, H_X). ¹³C-NMR (CDCl₃): δ (assignment): 88.13 (Cp, C(2,5)); 89.74 (Cp, C(3,4)), 108.31 (Cp, C(1)); 115.13 (=CH₂); 128.70 (-CH=); 236.97 (Cr-C≡O). IR(KBr): ν (cm⁻¹) (intensity): 2020 (vs), 1946 (vs), 1696 (vs), 628 (m).

2.2. Preparation of (*E*)-1,2-dicynichrodeneylethene (**7**)

Titanium tetrachloride (3.30 ml, 30 mmol) was added into a Schlenk tube containing lithium alu-

Table 1
Summary of crystal data and intensity collection of **7**^a

Empirical formula	C ₁₆ H ₁₀ O ₆ N ₂ Cr ₂
Formula weight	430.25
Diffractometer used	Nonius
Space group	<i>P</i> 2 ₁ / <i>c</i> ; monoclinic
<i>a</i> (Å)	6.379(5)
<i>b</i> (Å)	11.295(3)
<i>c</i> (Å)	11.9352(24)
β (°)	99.17(4)
Volume (Å ³)	849.0(7)
<i>Z</i>	2
Density (calculated) (g cm ⁻³)	1.683
λ (Å)	0.7093
<i>F</i> (000)	432
Number of reflections for indexing	25 ($14.44^\circ \leq 2\theta \leq 26.00^\circ$)
Scan type	0–2 θ
Scan width (°)	2(0.70 + 0.35 tan(θ))
Scan speed (° min ⁻¹)	2.06–8.24
2 θ maximum (°)	50
<i>h</i> , <i>k</i> , <i>l</i> ranges	(–7, 7), (0, 13), (0, 14)
μ (cm ⁻¹)	12.833
Crystal size (mm)	0.05 × 0.20 × 0.50
Transmission	0.865, 1.000
Temperature (K)	298
Number of measured reflections	1492
Number of observed reflections ($I > 2.0\sigma(I)$)	1071
Number of unique reflections	1492
<i>R</i> _p , <i>R</i> _w	0.035, 0.027
Goodness-of-fit	1.86
Refinement program	NRCVAX
Number of atoms	18
Number of parameters refined	138 (1071 out of 1492 reflections)
Minimize function	Sum($w F_o - F_c ^2$)
Weights scheme	(1/ $\sigma(F_o)^2$)
(Δ/σ) max	0.0085
(<i>D</i> -map) max, min (e Å ⁻³)	–0.260, 0.380

^a $R_f = \text{Sum}(F_o - F_c) / \text{Sum}(F_o)$; $R_w = \text{Sqrt}[\text{Sum}(w(F_o - F_c)^2) / \text{Sum}(wF_o^2)]$; goodness-of-fit = $\text{Sqrt}[\text{Sum}(w(F_o - F_c)^2) / (\text{no. of reflections} - \text{no. of parameters})]$.

Table 2
The contracted 2D HetCOR spectra of 2–7

Compound	R	^1H , Cp(Cr) ^a	2D HetCOR ^b	^{13}C , Cp(Cr) ^a
2	COOH			
3	CHO			
4	C(O)CH ₃			
5	NH ₂			
6	CH=CH ₂			
7	CH=CH-(C ₆ H ₅)Cr(CO) ₂ (NO)			

^a ○, (2,5); *, (3,4); the magnetic field increases towards the right.

^b The magnetic fields of ^1H - and ^{13}C -NMR spectra increase towards the right and upper side respectively.

minium hydride (0.57 g, 15 mmol) at 0°C. The temperature was slowly raised to 90°C and held for 12 h with stirring. After cooling down to 0°C, 100 ml of tetrahydrofuran and formylcynichrodene (**3**) (0.92 g, 4.0 mmol) were added into the residue. The reaction mixture was heated to 55°C and held for 16 h before cooling to 0°C, then 50 ml of a saturated potassium carbonate aqueous solution was added in slowly. Thereupon the reaction mixture was extracted with three 50 ml portions of dichloromethane. The combined extracts were washed three times with distilled water and dried with anhydrous magnesium sulfate. The solution was filtered and concentrated to a residue. The residue was dissolved in 30 ml of dichloromethane. Three grams of silica gel were added to the solution, and the solvent was then removed under vacuum. The residue was added to a dry-packed column (2.0 × 30 cm) of silica gel. Elution of the column with hexane:benzene (3:1) gave two bands. The first band was unidentified. After removal of solvent, the second (red) band gave (*E*)-1,2-dicynichrodenylethene (**7**) (0.31 g, 36%). An analytical sample (m.p. 186.5°C) was prepared by recrystallization using the solvent evaporation method from pentane:dichloromethane (5:1) at 0°C.

Anal. Found: C, 44.53; H, 2.46; N, 6.66. C₁₆H₁₀Cr₂N₂O₆. Calc.: C, 44.67; H, 2.34; N, 6.51%. Proton-NMR (CDCl₃): δ (relative intensity, multiplicity, assignment): 5.05 (2H, t, Cp H(3,4)); 5.25 (2H, t, Cp H(2,5)); 6.31 (2H, s, -CH=). ^{13}C -NMR (CDCl₃): δ (assignment): 88.57 (Cp, C(2,5)); 89.93 (Cp, C(3,4)), 106.72 (Cp, C(1)); 121.59 (-CH=); 236.70 (Cr-C=O).

IR(KBr): ν (cm⁻¹) (intensity): 2010 (vs), 1932 (vs), 1682 (vs), 644 (m). Mass spectrum: *m/z*: 430 (M⁺).

2.3. Preparation of (*E*)-1,2-diferrocenylethene (**10**)

To a solution of titanium trichloride (2.80 g, 18 mmol) in tetrahydrofuran, lithium aluminium hydride (0.34 g, 9.0 mmol) was added in an ice salt bath. After stirring for 30 min, formylferrocene (**9**) (1.00 g, 4.6 mmol) in 20 ml of tetrahydrofuran was added into the deep black solution. After stirring for 3 h at room temperature, 20 ml of hydrochloric acid (4 N) was added. The reaction mixture was then extracted with three 50 ml portions of dichloromethane. The combined extracts were washed three times with distilled water and dried with anhydrous magnesium sulfate. The solution was filtered. Ten grams of silica gel were added and the solvent was then removed under vacuum. The residue was added to a dry-packed column (2.0 × 10 cm) of silica. Elution of the column with hexane/benzene (1:1) gave two bands. The first band was unidentified. After removal of solvent under vacuum, the second (orange) band gave (*E*) (**10**) and (*Z*)-1,2-diferrocenylethene with a ratio of 3.5:1 (0.47 g, 25%). An analytical sample of **10** was prepared by recrystallization using the solvent evaporation method from hexane:dichloromethane (5:1) at 0°C.

Proton-NMR (CDCl₃): δ (relative intensity, multiplicity, assignment): 4.12 (10H, s, Cp²(Fe)); 4.24 (4H, t, Cp¹(Fe) H(3,4)); 4.38 (4H, t, Cp¹(Fe) H(2,5)); 6.41 (2H,




Fig. 1. 2D $^1\text{H}\{^{13}\text{C}\}$ -HetCOR spectrum of **7** in CDCl_3 .

s, $-\text{CH}=\text{}$). ^{13}C -NMR (CDCl_3): δ (assignment): 69.16 ($\text{Cp}^2(\text{Fe})$); 66.21 ($\text{Cp}^1(\text{Fe})$, C(2,5)); 68.54 ($\text{Cp}^1(\text{Fe})$, C(3,4)); 84.33 ($\text{Cp}^1(\text{Fe})$, C(1)); 123.63 ($-\text{CH}=\text{}$). IR(KBr): ν (cm^{-1}) (intensity): 1103 (w), 809 (m), 678 (s).

2.4. Preparation of vinylferrocene (**11**)

The preparation of vinylferrocene compound **11** was previously reported by Rausch and Seigel [9]. However, a more reliable method is described as following. (1-Hydroxyethyl)ferrocene (0.52 g, 2.24 mmol), anhydrous cupric sulfate (3 g, 18.73 mmol), and 5 mg of hydroquinone were dissolved in 100 ml of toluene. The

mixture was refluxed for 45 min and then cooled to room temperature. After the filtration the solvent was removed under vacuum. The residue was extracted with ether and dried with anhydrous magnesium sulfate. The solution was filtered, concentrated to 50 ml under vacuum. The residue was added to a dry-packed column (1.8 \times 9 cm) of silica gel. Elution of the column with hexane gave an orange band which upon removal of the solvent gave vinylferrocene (**11**) (0.40 g, 84%).

Proton-NMR (CDCl_3): δ (relative intensity, multiplicity, assignment): 4.11 (Cp^2); 4.21 (2H, t, Cp^1 H(3,4)); 4.36 (2H, t, Cp^1 H(2,5)); 5.03 (1H, dd, H_B , $J_{\text{AB}} = 3.3$, $J_{\text{BX}} = 10.8$); 5.34 (1H, dd, H_A , $J_{\text{AX}} = 16.8$); 6.46 (1H, dd, H_X). ^{13}C -NMR (CDCl_3): δ (assignment):

Table 3

¹H-NMR chemical shifts of selected monosubstituted cynichrodene^a, ferrocene^b and benzene^c from tetramethylsilane and Δ^d

R	(CO) ₂ (NO)Cr(C ₅ H ₄ -R)			(C ₅ H ₅)Fe(C ₅ H ₄ -R)			C ₆ H ₅ -R					
	δ (ppm)		Δ (ppm)	δ (ppm)		Δ (ppm)	δ (ppm)			Δ (ppm)		
	H(2,5)	H(3,4)		H(2,5)	H(3,4)		H(2)	H(3)	H(4)			
<i>Electron-withdrawing substituents by resonance</i>												
-CHO	3	5.77	5.27	0.50	9	<u>4.70</u>	4.47	0.23	<u>7.80</u>	7.44	7.55	0.25
-C(O)CH ₃	4	5.72	5.16	0.56		<u>4.66</u>	4.36	0.30	<u>7.91</u>	7.38	7.48	0.43
<i>Electron-donating substituents by resonance</i>												
-NH ₂	5	4.6	<u>4.81</u>	-0.21		<u>3.83</u>	3.7	0.13	6.57	7.09	<u>6.68</u>	-0.11
-CH=CH ₂	6	5.2	5.01	0.19	11	<u>4.36</u>	4.21	0.15	<u>7.51</u>	7.41	7.35	0.16
-CH=CH-R'	7	5.25	5.05	0.20	10	<u>4.38</u>	4.24	0.14	<u>7.52</u>	7.36	7.26	0.26
(E)	R' = (C ₅ H ₄)Cr(CO) ₂ (NO)				R' = (C ₅ H ₄)Fe(C ₅ H ₅)				R' = C ₆ H ₅			

^a From [2].^b From [13].^c From [17].^d Δ = δ[H(2,5)] - δ[H(3,4)] for ferrocene and cynichrodene derivatives; δ[H(2)] - δ[H(4)] for benzene derivatives. The lower-field chemical shift of each pair is underlined.

Table 4

¹³C-NMR chemical shifts of selected monosubstituted cynichrodene^a, ferrocene^b and benzene^c from tetramethylsilane and Δ^d

R	(CO) ₂ (NO)Cr(C ₅ H ₄ -R)			(C ₅ H ₅)Fe(C ₅ H ₄ -R)			C ₆ H ₅ -R					
	δ (ppm)		Δ (ppm)	δ (ppm)		Δ (ppm)	δ (ppm)			Δ (ppm)		
	C(2,5)	C(3,4)		C(2,5)	C(3,4)		C(2)	C(3)	C(4)			
<i>Electron-withdrawing substituents by resonance</i>												
-CHO	3	<u>93.5</u>	92.8	0.70	9	68.0	<u>72.6</u>	-4.6	129.80	129.1	<u>134.0</u>	-4.20
-C(O)CH ₃	4	<u>93.6</u>	92.0	1.6		69.2	<u>71.8</u>	-2.6	128.6	128.5	<u>132.7</u>	-4.10
<i>Electron-donating substituents by resonance</i>												
-NH ₂	5	73.9	<u>85.1</u>	-11.2		58.8	<u>63.0</u>	-4.2	115.3	129.4	<u>118.7</u>	-3.40
-CH=CH ₂	6	88.1	<u>89.7</u>	-1.6	11	66.7	<u>68.6</u>	-1.9	126.1	128.3	<u>127.6</u>	-1.50
-CH=CH-R'	7	88.6	<u>89.9</u>	-1.3	10	68.5	<u>68.5</u>	-2.3	126.8	128.9	<u>127.8</u>	-1.00
(E)	R' = (C ₅ H ₄)Cr(CO) ₂ (NO)				R' = (C ₅ H ₄)Fe(C ₅ H ₅)				R' = C ₆ H ₅ -R			

^a From [2,18].^b From [14].^c From [17,19].^d Δ = δ[C(2,5)] - δ[C(3,4)] for ferrocene and cynichrodene derivatives; δ[C(2)] - δ[C(4)] for benzene derivatives. The lower-field chemical shift of each pair is underlined.

63.53 (Cp¹, C(1)); 66.69 (Cp¹, C(2,5)); 68.63 (Cp¹, C(3,4)); 69.21 (Cp²(Fe)); 111.02 (=CH₂); 134.63 (-CH=).

2.5. X-ray diffraction analysis of **7**

The intensity data were collected on a CAD-4 diffractometer with a graphite monochromator (Mo-K_α radiation). θ-2θ scan data were collected at room temperature (24°C). The data were corrected for absorption, Lorentz and polarization effects. The absorption correction is according to the empirical psi rotation. The details of crystal data and intensity collection are summarized in Table 1.

The structure was solved by direct methods and was

refined by full matrix least squares refinement based on *F* values. All of the non-hydrogen atoms were refined with anisotropic thermal parameters. All of the hydrogen atoms were positioned at calculated coordinates with a fixed isotropic thermal parameter (*U* = *U*(attached atom) + 0.01 Å²). Atomic scattering factors and corrections for anomalous dispersion were from [10]. All calculations were performed on a DEC alpha workstation using the NRCVAX programs [11].

3. Results and discussion

Acid-catalyzed dehydration of alcohol **8** in refluxing benzene, using *p*-toluene sulfonic acid, on the presence

of hydroquinone as a radical inhibitor, gave olefin **6** in yield as high as 84%. Using anhydrous cupric sulfate instead of *p*-toluene sulfonic acid, vinylferrocene (**11**) was obtained from the corresponding alcohol. By reacting with low-valent titanium coupling reagent, prepared from two equivalents of titanium tetrachloride and one equivalent of lithium aluminum chloride, formylcynichrodene (**3**) was transformed into **7** in 36% yield. An analogous method using titanium trichloride as coupling agent was employed to prepare compound **10** from **9** with 20% yield.







Table 5
Atomic parameters *x*, *y*, *z* and B_{eq}^a of **7**, estimated S.D.s refer to the last digit printed

	<i>x</i>	<i>y</i>	<i>z</i>	B_{eq}
Cr	0.25739(9)	0.22020(5)	0.07397(5)	3.40(3)
N1	0.4863(5)	0.2069(3)	0.1757(3)	4.73(17)
C2	0.3740(6)	0.1406(4)	-0.0326(3)	3.88(19)
C3	0.1365(6)	0.0874(4)	0.1225(3)	3.86(19)
C5	0.4368(7)	0.4630(3)	-0.0293(4)	4.13(22)
C6	0.2558(6)	0.4036(3)	0.0092(3)	3.39(18)
C7	0.1003(7)	0.3375(4)	-0.0618(4)	4.28(21)
C8	-0.0516(7)	0.2978(4)	0.0019(5)	5.01(25)
C9	0.0069(7)	0.3390(4)	0.1130(5)	4.83(24)
C10	0.1941(7)	0.4036(4)	0.1175(4)	4.10(21)
O1	0.6377(5)	0.1979(3)	0.1433(3)	6.82(18)
O2	0.4499(5)	0.0924(3)	-0.10142(25)	6.69(18)
O3	0.0628(5)	0.0039(3)	0.1540(3)	6.92(19)
H5	0.519(6)	0.546(3)	0.104(3)	6.1(11)
H7	0.097(5)	0.325(3)	-0.142(3)	3.2(8)
H8	-0.177(6)	0.252(4)	-0.024(3)	7.0(12)
H9	-0.057(5)	0.332(3)	0.180(3)	4.2(9)
H10	0.260(5)	0.437(3)	0.172(3)	2.7(9)

^a B_{eq} is the mean of the principal axes of the thermal ellipsoid.

Table 6
Selected bond distances (Å) and angles (°) of **7**

Bond distances			
Cr–C6	2.211(4)	Cr–C7	2.206(4)
Cr–C8	2.203(4)	Cr–C9	2.193(4)
Cr–C10	2.188(4)		
C6–C7	1.411(6)	C6–C10	1.410(6)
C7–C8	1.397(7)	C8–C9	1.399(8)
C9–C10	1.393(7)	Cr–N1	1.750(4)
Cr–C2	1.812(4)	Cr–C3	1.823(4)
N1–O1	1.161(4)	C2–O2	1.154(5)
C3–O3	1.144(5)	C5–C6	1.471(6)
C5–C5	1.288(8)		
Cr⋯centroid(Cp)	1.849	Cr⋯C5	3.287
C5⋯C10	3.116	H(C5)⋯H(C10)	3.79
Bond angles			
C6–C7–C8	109.0(4)	C7–C8–C9	107.8(4)
C7–C6–C10	106.1(4)	C8–C9–C10	107.9(4)
CO–C10–C9	109.1(4)	N1–Cr–C2	93.10(17)
N1–Cr–C3	93.30(17)	C2–Cr–C3	93.26(18)
Cr–N1–O1	179.8(3)	Cr–C2–O2	178.4(4)
Cr–C3–O3	179.1(4)	C5–C6–C7	124.4(4)
C5–C6–C10	129.4(4)	C5–C5–C6	125.5(4)
Centroid(Cp)–Cr–N1	126.8		
Centroid(Cp)–Cr–C2	120.4	centroid(Cp)–Cr–C3	121.6

Compound **7** exhibits two carbonyl stretching bands, the symmetric mode occurring at 2010 cm^{-1} and the asymmetric mode at 1932 cm^{-1} . The nitrosyl stretching band is observed at 1682 cm^{-1} . It is interesting to compare the three stretching frequencies of **7** with the corresponding bands of its unsubstituted parent compound **1** (2025, 1955; 1695 cm^{-1}). The lower-frequency shift by 15, 23 and 13 wave-numbers indicate that the vinyl group is exerting as an electron donating group to the two adjacent cynichrodene moieties. The data is consistent with the result found from the study of the ¹³C-NMR spectra.




Fig. 2. Molecular configuration of **7**.




Fig. 3. View of **7** along the normal of Cp(Cr) ring.


The $^1\text{H-NMR}$ spectra of **6**, **7**, and **10** were consistent with their structures and similar to other metallocenyl systems [12,13]. The strong diamagnetic anisotropic effect of the vinyl group on the ring protons might explain why the protons (2- and 5-positions) closer to it were deshielded to a greater extent than those (3- and 4-positions) further away from it.

From Table 2, the contracted 2D HetCOR spectra of **2–7** (Fig. 1), it is interesting to find that compounds **2–4** exhibit positive slopes, while **6** and **7** exhibit negative slopes. The opposite slopes may be interpreted as follows. First, the strong diamagnetic anisotropic effects of the carbonyl and vinyl group on the ring protons might explain why the protons (2- and 5-positions) closer to it were deshielded to the lower field for all compounds of **2–4** and **6–7**. These results are in agreement with the corresponding benzene and ferrocene analogues (Table 3). Second, the shielding of C(2,5) and C(3,4) carbon atoms of **6** and **7** are analogous to **5** instead of **2–4** and suggests that the electron density distribution in the cyclopentadienyl rings of **6–7** is similar to **5**, rather than **2–4**.

The assignment of $^{13}\text{C-NMR}$ spectra of **6**, **7**, and **10** were based on standard $^{13}\text{C-NMR}$ correlations [14], 2D HetCOR (Table 2), the DEPT technique and by comparison with other metallo-aromatic systems [15].

It is interesting to compare the $^{13}\text{C-NMR}$ spectra of **3**, **6**, and **7** with their unsubstituted parent compound **1** (Table 4). For the carbon atoms on Cp(Cr) (C(3,4) and C(2,5)), the chemical shifts of **3** occur at a lower field than the chemical shifts of **1** at $\delta = 90.31$ ppm. However, the chemical shifts of **6** and **7** occur at a higher field than $\delta = 90.31$ ppm. This reflects the strong electron-withdrawing effect of the formyl group on **3** and the strong electron-donating effect of vinyl group on **6** and **7**. It is worth pointing out from Table 4 that the chemical shifts of C(3,4) occur at a higher field than the chemical shifts of C(2,5) for **3** and **4**. On the contrary,

the chemical shifts of C(3,4) occur at a lower field than the chemical shifts of C(2,5) for **5–7**. Earlier, we have reported thorough spectra studies on aminocynchrodenone (**5**) [2], a compound with a strong electron-donating substituent, $-\text{NH}_2$ on Cp. We also compared the chemical shifts of selected monosubstituted cynchrodenone derivatives with the NMR data of their analogues of ferrocene and benzene derivatives (Table 4). The large contribution of canonical form **5a** to **5** explained the relatively large negative difference Δ (-11.2) in C(2,5) and C(3,4). This is understandable in the stabilization of chromium anion because of the overall electron-withdrawing properties of CO and NO ligands.



As is well known, the vinyl group can exert an either electron-donating or electron-withdrawing effect by resonance to its attached group. When an electron-donating group is attached to it, the vinyl group withdraws the electron from the group, and vice versa.

Upon examination of the infrared spectra of **6** and **7** (Tables 2 and 4), the following conclusions may be drawn:

1. the vinyl group exerts the same kind of electronic effect as the amino group, rather than carbonyl group, to its adjacent Cp ring;
2. in compounds **6** and **7**, the vinyl group donates electrons to the adjacent Cp ring(s), which in turn transfers the electron density to the metal chromium. This resulted in an enhanced π -back bonding phenomena observed in IR spectra;
3. canonical **7a**, rather than **7b**, to some extent contributes to the structure of **7**.


Table 7
Selected structural data of **7**, **12** and **13**^a

Compound	Bond length (Å)									
	Cr–C(ring)	Cr–NO	Cr–CO	N=O	C=O	Cr–N–O	Cr–C–O	ω_{Cr}^b (°)	θ_{Cr}^c (°)	Cr...exocyclic atom
7	2.200(4)	1.750(4)	1.812(4) 1.823(4)	1.161(4)	1.154(5) 1.144(5)	179.8(3)	178.4(4) 179.1(4)	46.5	–1.2	3.287
12	2.195(22)	1.747(18)	1.798(18) 1.828(23)	1.187(23)	1.126(27) 1.160(22)	174.5(16)	177.0(18) 177.2(19)	8.3	–2.0	3.275
	2.177(25)	1.807(33)	1.745(21) 1.758(17)	1.189(40)	1.130(20) 1.149(25)	179.2(27)	173.8(25) 179.8(25)	9.7	–2.1	3.258
13	2.205(5)	1.712(4)	1.864(4) 1.846	1.178(5)	1.135(5) 1.135(5)	179.4(3)	179.0(4) 177.2(4)	177.2	1.0	3.223

^a **12** (CO)₂(NO)Cr(η^5 -C₅H₄)–NH–C(O)–NH–(η^5 -C₅H₄)Fe(η^5 -C₅H₅); **13** (CO)₂(NO)Cr(η^5 -C₅H₄)C(O)(η^5 -C₅H₄)Fe(η^5 -C₅H₅).

^b ω_{Cr} (°): the twist angle is defined as the torsional angle between the nitrosyl nitrogen atom, the chromium atom, the Cp center and the ring carbon atom bearing the exocyclic carbon atom.

^c θ_{Cr} (°): the angle θ is defined as the angle between the exocyclic bond and the corresponding Cp ring with positive angle towards metal and negative angles away from the metal.



The molecular structure of **7** is shown in Fig. 2. The atomic coordinates of the non-hydrogen atoms are listed in Table 5. Selected bond distances and angles are given in Table 6.

Compound **7** adopts a *transoid* conformation at the organic vinyl carbons. The coordination geometry about the Cr center is approximately a distorted tetrahedron with two carbonyl groups, the Cp group and nitrosyl group as the four coordination sites. The nitrosyl group is located at the side toward the exocyclic carbon atom of Cp(Cr) with a twist angle of 46.5° (Fig. 3). The twist angle is defined as the torsional angle between the nitrosyl nitrogen atom, the chromium atom, the Cp center and the ring carbon atom bearing the exocyclic carbon atom.

In the cycindrodene moiety, the observed average bond length of Cr–C(ring) is 2.200(4) Å. The Cr–N length of 1.750(4) (Cr–N1) is closer to those found in (CO)₂(NO)Cr(η^5 -C₅H₄)–NH–C(O)–NH–(η^5 -C₅H₄)

Fe(η^5 -C₅H₅) (**12**) [2] (1.747(18), 1.803(33) Å) than the values found in (CO)₂(NO)Cr(η^5 -C₅H₄)C(O)(η^5 -C₅H₄)Fe(η^5 -C₅H₅) (**13**) (1.712(4) Å) [1]. The Cr–C (carbonyl) distances of 1.812(4) Å (Cr–C2) and 1.823(4) Å (Cr–C3) are between those found in **12** (average 1.782(23) Å) [2] and those found in **13** (average 1.855(4) Å) [1]. The Cr–N–O angle of 179.8(3)° is consistent with the NO⁺ formalism typical of the linear M–NO linkage. The Cr–C–O angles of 178.4(4)° (Cr–C2–O2) and 179.1(4)° (Cr–C3–O3) indicate the usual mode of bonding in the terminal metal carbonyl complexes. The Cr–centroid (Cp(Cr)) distance of 1.849 Å agrees with the values of 1.843 Å in **12** [2] and 1.846 Å in **13** [1]. The average C–C distance in the ring (Cp(Cr)) is 1.402 Å. Selected structural data of **7**, **12** and **13** are listed in Table 7.

The exocyclic C=C bond length is 1.228(8) Å (C5–C5). The exocyclic carbon atom C5 is bent away from the corresponding Cr atom, with an angle $\theta = -1.2^\circ$. The angle θ is defined as the angle between the exocyclic C6–C5 bond and the corresponding Cp ring with positive angle towards metal and negative angle away from the metal. The vinyl plane turns away from the ring planes by 9.1°. The two Cp planes are coplanar.

Upon examination of the bonds of Cr–NO, Cr–CO, N=O, and C=O, it is interesting to find that **7** has the longer bond distances of Cr–NO and C=O, while it has the shorter bond distances of Cr–CO and N=O than **13** (Table 5). The contributions of canonical form **7ii** to **7** and **13i** to **13** may explain such results.

- [8] C.U. Pittman Jr., T.D. Rounsefell, E.A. Lewis, J.E. Sheata, M.D. Rausch, E.A. Mintz, *Macromolecules* 11 (1978) 560.
- [9] M.D. Rausch, A. Siegel, *J. Organomet. Chem.* 11 (1968) 317.
- [10] Scattering factors from *Int. Tab. Vol. 4: International Tables for X-ray Crystallography*, Vol. IV, Kynoch, Birmingham, England, 1974.
- [11] Full system reference: E.J. Gabe, Y. LePage, J.-P. Charland, F., L. Lee, P.S. White, *J. Appl. Crystallogr.* 22 (1989) 384.
- [12] M.D. Rausch, E.A. Mintz, D.W. Macomber, *J. Org. Chem.* 45 (1980) 689.
- [13] E.W. Slocum, C.R. Ernst, *Adv. Organomet. Chem.* 10 (1972) 79.
- [14] J.B. Stotter (Ed.), *Carbon-13 NMR Spectroscopy*, Academic Press, New York, 1972.
- [15] B.E. Mann, *Adv. Organomet. Chem.* 12 (1974) 135.
- [16] P.A. Downton, B.G. Sayer, M.J. McGlinchey, *Organometallics* 11 (1992) 3281.
- [17] R.M. Silverstein, G.C. Bassler, T.C. Morrill, *Spectrometric Identification of Organic Compounds*, Wiley, New York, 1981.
- [18] M.D. Rausch, D.J. Kowalski, E.A. Mintz, *J. Organomet. Chem.* 342 (1988) 201.
- [19] E. Breitmaier, W. Voelter, *Carbon-13 NMR Spectroscopy*, VCH, New York, 1987.