


Using Molecular Iodine in Direct Oxidative Condensation of Aldoses with Diamines: An Improved Synthesis of Aldo-benzimidazoles and Aldo-naphthimidazoles for Carbohydrate Analysis

Chunchi Lin,[†] Po-Ting Lai,[‡] Sylvain Kuo-Shiang Liao,[†] Wei-Ting Hung,[†]
Wen-Bin Yang,^{*,†} and Jim-Min Fang^{*,†,‡}

The Genomics Research Center, Academia Sinica, Taipei, 11529, Taiwan, and Department of Chemistry,
National Taiwan University, Taipei 106, Taiwan

wbyang@gate.sinica.edu.tw; jmfang@ntu.edu.tw

Received January 29, 2008




A practical method has been developed for conversion of unprotected and unmodified aldoses to aldo-benzimidazoles and aldo-naphthimidazoles. Using iodine as an oxidant or promoter in acetic acid solution, a series of mono-, di-, and trialdoses, including those containing carboxyl and acetamido groups, undergo an oxidative condensation reaction with *o*-phenylenediamine or 2,3-naphthalenediamine at room temperature to give the aldo-benzimidazole and aldo-naphthimidazole products in high yields. No cleavage of the glycosidic bond occurs under such mild reaction conditions. The composition analysis of saccharides is realized by the HPLC analysis of the fluorescent naphthimidazole derivatives.

Introduction

In nature, 5,6-dimethyl-1-(α -D-ribofuranosyl)benzimidazole exists as a moiety of vitamin B₁₂.¹ The furanosyl benzimidazole can be considered as a C-nucleoside mimic.¹ Due to its structural similarity to purine, benzimidazole is also an important chemical entity in pharmaceuticals.² In one approach, aldo-benzimidazoles are prepared by condensation of aldonic acids or aldonic δ -lactones with *o*-phenylenediamines (Scheme 1).³ The condensation reactions are often conducted in strong acidic conditions at elevated temperature (e.g., 135 °C).^{3a} The tetritol-1-yl

SCHEME 1. Previous Synthetic Routes to Saccharide Benzimidazoles



[†] The Genomics Research Center.

[‡] National Taiwan University.

(1) Townsend, L. B.; Revankar, G. R. *Chem. Rev.* **1970**, *70*, 389–438.

(2) Freeman, G. A.; Selleseth, D. W.; Rideout, J. L.; Harvey, R. J. *Nucleosides Nucleotides Nucleic Acids* **2000**, *19*, 155–174.

(3) (a) Moore, S.; Link, K. P. *J. Biol. Chem.* **1940**, *133*, 293–311. (b) El Ashry, E. S. H.; Awad, L. F.; Hamid, H. A.; Atta, A. I. *J. Carbohydr. Chem.* **2007**, *26*, 329–338.

(4) Sallam, M. A. E.; Ibrahim, E. I.; El-Eter, K. A. A.; Cassady, J. M. *Carbohydr. Res.* **1997**, *298*, 93–104.

(5) Moore, S.; Link, K. P. *J. Org. Chem.* **1940**, *5*, 637–644.

(6) (a) Wistler, R. L.; Wolfrom, M. L., Eds. In *Methods in Carbohydrates Chemistry*, Academic Press: New York, 1963; Vol. 2, pp 13–14. (b) Zhang, T. H.; Marchant, R. E. *Macromolecules* **1994**, *27*, 7302–7308. (c) Auzély-Velty, R.; Cristea, M.; Rinaudo, M. *Biomacromolecules* **2002**, *3*, 998–1005. (d) Yang, B. Y.; Montgomery, R. *Carbohydr. Res.* **2005**, *340*, 2698–2705.

benzimidazoles can be further converted to furanosyl benzimidazoles by a Lewis acid promoted dehydrative cyclization.⁴

In another approach, aldoses are subjected to oxidative condensation with *o*-phenylenediamine to give aldo-benzimidazoles in a direct manner.⁵ This approach mitigates the

TABLE 1. Oxidative Condensation of Aldoses with *o*-Phenylenediamines or 2,3-Naphthalenediamine^a

entry	sugar	diamine ^a (equiv)	iodine (equiv)	solvent	imidazole (yield, %) ^b	peracetate (yield, %) ^c
1	D-glucose, 1a	C ₆ H ₄ (NH ₂) ₂ , 2a (1.1)	0.1	AcOH	3a (98)	
2	1a	Me ₂ C ₆ H ₂ (NH ₂) ₂ , 2b (3)	1.0	aq. AcOH		3b-Ac (58)
3	1a	Cl ₂ C ₆ H ₂ (NH ₂) ₂ , 2c (2)	0.1	AcOH	3c (98)	
4	1a	MeC ₆ H ₃ (NH ₂) ₂ , 2d (2)	0.1	AcOH	3d (55)	
5	1a	O ₂ NC ₆ H ₃ (NH ₂) ₂ , 2e (2)	0.1	AcOH	3e (76) ^d	
6	1a	PhCOC ₆ H ₃ (NH ₂) ₂ , 2f (2)	0.1	AcOH	3f (74) ^d	
7	1a	C ₁₀ H ₆ (NH ₂) ₂ , 2g (1.1)	0.1	AcOH	3g (98)	
8	1a	2g (1.1)	0	AcOH	3g (33)	
9	D-galactose, 1b	2a (3)	1.0	aq. AcOH		4a-Ac (57)
10	1b	2a (2)	1.2	buffer ^e		4a-Ac (88)
11	1b	2b (3)	1.0	aq AcOH		4b-Ac (80)
12	1b	2g (1.1)	0.1	AcOH	4g (85)	
13	D-mannose, 1c	2a (3)	1.0	aq AcOH		5a-Ac (42)
14	1c	2b (3)	1.0	aq AcOH		5b-Ac (74)
15	1c	2g (1.1)	0.1	AcOH	5g (96)	
16	D-arabinose, 1d	2a (3)	1.0	aq AcOH		6a-Ac (60)
17	1d	2a (2)	1.2	buffer ^e		6a-Ac (88)
18	1d	2b (3)	1.0	aq. AcOH		6b-Ac (71)
19	1d	2g (1.1)	0.1	AcOH	6g (90)	
20	D-xylose, 1e	2a (3)	1.0	aq AcOH		7a-Ac (63)
21	1e	2a (2)	1.2	buffer ^e		7a-Ac (83)
22	1e	2b (3)	1.0	aq AcOH		7b-Ac (87)
23	1e	2g (1.1)	0.1	AcOH	7g (98)	
24	D-ribose, 1f	2g (1.1)	0.1	AcOH	8g (97)	
25	L-fucose, 1g	2g (1.1)	0.1	AcOH	9g (99)	
26	L-rhamnose, 1h	2g (1.1)	0.1	AcOH	10g (99)	
27	D-GlcNAc, 1i	2g (1.1)	0.1	AcOH	11g (94)	
28	D-GlcA, 1j	2g (1.1)	0.1	AcOH	12g (97)	
29	D-maltose, 1k	2a (2)	1.2	buffer ^e		13a-Ac (78)
30	1k	2b (3)	1.0	aq AcOH		13b-Ac (66)
31	1k	2g (1.1)	0.1	AcOH	13g (99)	
32	D-lactose, 1l	2a (2)	1.2	buffer ^e		14a-Ac (88)
33	1l	2b (3)	1.0	aq AcOH		14b-Ac (53)
34	1l	2g (1.1)	0.1	AcOH	14g (98)	
35	D-cellobiose, 1m	2g (1.1)	0.1	AcOH	15g (90)	
36	D-maltotriose, 1n	2a (2)	1.0	AcOH	16a (51)	

^a The reaction was conducted at room temperature. ^b The yield of the isolated aldo-benzimidazole product obtained by trituration of the reaction mixture with EtOAc. ^c Overall yield of isolated product after peracetylation with Ac₂O in pyridine. ^d The reaction was conducted at 60 °C for 18 h. ^e Acetic acid buffer solution (pH = 4.38).

laborious work in prior preparation of aldonic acids from the corresponding aldoses.⁶ For example, Moore and Link have previously applied this approach to synthesize two monosaccharide benzimidazoles, D-gluco- and D-galactobenzimidazoles, in 24 and 50% yields from D-glucose and D-galactose, respectively, by heating with cupric acetate (2 equiv) at 53 °C for 14 h in aqueous acetic acid solution (Scheme 1).⁵ Some saccharide bis-Schiff bases and quinoxalines also form as side products under such reaction conditions.^{3,5} However, using catalytic amounts of CuSO₄ or replacing the AcOH solvent with aqueous MeOH does not give any desired aldo-benzimidazoles.⁵ Considering the acid-sensitive internal glycosidic bonds in disaccharides and higher saccharides, it is questionable that the corresponding aldo-benzimidazoles can be obtained by the above-mentioned reactions involving acidic conditions at high reaction temperature.

As a continuation of our study of the novel transformation of aldehydes and aldoses by use of amines and iodine,⁷ we describe herein an improved protocol for the synthesis of various aldo-benzimidazoles and aldo-naphthimidazoles by direct oxidative condensation of aldoses, including mono-, di-, and trisac-

charides, with *o*-phenylenediamines and 2,3-naphthalenediamine in the presence of iodine. We also demonstrate that the composition analysis of carbohydrate molecules is facilitated by incorporating the naphthimidazole moiety as a fluorescent label for sensitive detection.

Results and Discussion

Molecular iodine is a convenient and environmental benign oxidizing agent in organic synthesis.⁸ Two recent studies show that aliphatic and aromatic aldehydes react with 1,2-diamines to give the corresponding imidazolines in the presence of stoichiometric amounts of iodine.⁹ The reactions are conducted at high temperature (70–90 °C) in protic solvents (H₂O or *t*-BuOH) using K₂CO₃ as the base to neutralize the generated hydroiodic acid. In one study,^{9a} KI is also used as a promoter in the reaction. So far, these methods have not been applied to aldoses, presumably because the hydroxyl groups in aldoses may also be oxidized by I₂/K₂CO₃ at high temperature.^{7d,10}

At the outset of our study, we tested the direct oxidative condensation of D-glucose (**1a**) with *o*-phenylenediamine (**2a**)

(7) (a) Talukdar, S.; Hsu, J.-L.; Chou, T.-C.; Fang, J.-M. *Tetrahedron Lett.* **2001**, *42*, 1103–1105. (b) Chen, M.-Y.; Hsu, J.-L.; Shie, J.-J.; Fang, J.-M. *J. Chin. Chem. Soc.* **2003**, *50*, 129–133. (c) Shie, J.-J.; Fang, J.-M. *J. Org. Chem.* **2003**, *68*, 1158–1160. (d) Shie, J.-J.; Fang, J.-M. *J. Org. Chem.* **2007**, *72*, 3141–3144.

(8) Togo, H.; Iida, S. *Synlett* **2006**, *14*, 2159–2175, and references cited therein.

(9) (a) Gogoi, P.; Konwar, D. *Tetrahedron Lett.* **2006**, *47*, 79–82. (b) Ishihara, M.; Togo, H. *Synlett* **2006**, 227–230.

(10) Mori, N.; Togo, H. *Tetrahedron* **2005**, *61*, 5915–5925.





FIGURE 1. Structures of the products of aldo-benzimidazoles, aldo-naphthimidazoles, and their peracetylation compounds.

in the $I_2/KI/K_2CO_3/H_2O$ system (90 °C, 1 h) similar to that reported by Gogoi and Konwar.^{9a} However, this operation failed to give the desired aldo-benzimidazole product. We also surveyed the oxidative condensation reaction of D-glucose with *o*-phenylenediamine and iodine in various solvents, including $CHCl_3$, THF, Me_2CO , CH_3CN , DMF, DMSO, MeOH, H_2O , and concentrated HCl solution. None of them are suitable media for the anticipated reaction. After many attempts, we found that aqueous AcOH solution was the solvent of choice, and the oxidative condensation of saccharides with *o*-phenylenediamines (or 2,3-naphthalenediamine) occurred readily at room temperature without using additives of KI or K_2CO_3 . Thus, D-glucose was treated with *o*-phenylenediamine (1.1 equiv) and iodine (0.1 equiv) in acetic acid at room temperature for 8 h to give D-glucobenzimidazole (**3a**) in 98% yield (entry 1, Table 1). The reaction was monitored by TLC analysis, and complete conversion of glucose to glucobenzimidazole was supported by the ^{13}C NMR spectrum of the reaction mixture. The crude product mixture was then triturated with EtOAc to give a practically pure D-glucobenzimidazole as a solid in nearly quantitative yield.

The sample could be purified by flash column chromatography for further analysis. The ^{13}C NMR spectrum of **3a** in $DMSO-d_6$ solution showed a signal at δ 156.2 attributable to the C-2 of the newly formed benzimidazole ring (referring to the numbering in Figure 1). In the 1H NMR spectrum ($DMSO-d_6$), the phenyl protons in benzimidazole appeared at lower fields (δ 7.71 and 7.48) than those in *o*-phenylenediamine. The proton at the C-1' position of **3a** occurred at δ 5.16 as a doublet with a small coupling constant (5.2 Hz). The benzimidazole **3a** was treated with Ac_2O in pyridine to give the peracetylation derivative **3a-Ac**, which was fully characterized by its physical and spectroscopic properties.

The similar reaction of D-glucose with 4,5-dimethyl-1,2-benzenediamine (**2b**) and iodine (1 equiv) in aqueous AcOH solution, followed by acetylation in Ac_2O /pyridine, gave the peracetylation compound **3b-Ac** in 58% overall yield. The iodine-promoted oxidative condensation reactions of D-glucose with 4,5-dichloro-1,2-benzenediamine (**2c**) and 4-methyl-1,2-benzenediamine (**2d**) were also effectively carried out at room temperature to afford the aldo-benzimidazoles **3c** and **3d**,

SCHEME 2. Oxidative Condensation of D-Glucose with *o*-Phenylenediamine Using I₂ as the Oxidizing Agent.


whereas the reactions with 4-nitro-1,2-benzenediamine (**2e**) and 4-benzoyl-1,2-benzenediamine (**2f**) were conducted at an elevated temperature (60 °C). It appeared that the oxidative condensation reactions were accelerated by the electron-donating substituents and decelerated by the electron-withdrawing groups on the phenyl ring of *o*-phenylenediamine.

The oxidative condensation of glucose with 2,3-naphthalenediamine using I₂ (0.1 equiv) as the promoter gave 98% yield of the desired aldo-naphthimidazole **3g** (entry 7, Table 1). However, the yield of **3g** greatly deteriorated to 33% in the absence of I₂ (entry 8).

On the basis of the experimental results, we speculated that the reaction was initiated by formation of a Schiff base (e.g., the intermediate **A** in Scheme 2) by condensation of the aldose with one of the amino groups in *o*-phenylenediamine (**2a–f**) or naphthalenediamine **2g**. The condensation of an aldehyde with an amine to form the Schiff base is a reversible process, in which formation of the Schiff base is favored under mild acidic conditions (e.g., at pH 4–6). This rationale may support our finding that acetic acid is an appropriate solvent for the transformation of glucose to aldo-imidazoles. The subsequent nucleophilic addition of the other amino group to Schiff base **A** would give an imidazoline intermediate **B**, which could be oxidized in air or by iodine to afford the observed product of aldo-benzimidazole. In deed, a previous report¹¹ has shown that aromatic aldehydes are converted to the corresponding benzimidazoles by heating with *o*-phenylenediamines at 100 °C in dioxane solution using air as the oxidant. The aldo-imidazole products might also form via a different pathway.¹² A prior *N*-iodination of the imine moiety in Schiff base **A** would facilitate the formation of iodoimidazoline intermediate **C**, and the desired product of aldo-benzimidazole would be obtained by the subsequent elimination of an HI molecule. It was noted that another HI molecule was also generated during *N*-iodination of the Schiff base. In the HOAc media, the released I[−] ions might be oxidized in air to regenerate I₂. Our model experiment showed that colorless KI salt in acetic acid solution was gradually turned into a yellow solution of iodine by stirring in air at room temperature. This rationale may explain the




FIGURE 2. HPLC chromatograph of pentose and hexose naphthimidazoles on a pair of HC-C₁₈ column (250 mm × 4.6 mm; 5 μm porosity) at 30 °C. The mobile phase is sodium phosphate buffer (100 mM, pH 5.0) containing 28% methanol and 2% acetonitrile at a flow rate of 0.8 mL/min. The fluorescence detection wavelength is 326 nm, and the excitation wavelength is 232 nm. The sample is a mixture of aldo-naphthimidazoles, including those derived from D-mannose (**5g**, peak 1), D-glucose (**3g**, peak 2), D-galactose (**4g**, peak 3), D-ribose (**8g**, peak 4), D-arabinose/D-xylose (**6g/7g**, peak 5 + 6), and L-fucose (**9g**, peak 7) at a concentration of 10 ppm for each component.

quantitative yield of the aldo-imidazole product that was obtained even with only 10 mol % of iodine as the oxidation promoter.

By using I₂ as the oxidant/promoter in AcOH solution, various mono-, di-, and trisaccharides (**1b–n**) also reacted with diamines **2a–g** to give the corresponding oxidative condensation products (entries 9–36 in Table 1). No racemization at the stereogenic centers or cleavage of the glycoside bonds was observed under such mild reaction conditions. *N*-Acetylglucosamine (**1i**) and glucuronic acid (**1j**) were also smoothly converted to the corresponding naphthimidazoles **11g** and **12g** in 94% and 97% yields, respectively (entries 27 and 28), indicating that the acetamido and carboxyl groups were inert in such reaction conditions. The structure of aldo-benzimidazole **16a** derived from D-maltotriose was rigorously determined by NMR analyses including the two-dimensional HSQC spectrum. Our current method for a direct conversion of trisaccharide to the corresponding imidazole product is unique and likely applicable to the oxidative condensation of oligo- and polysaccharides with various aromatic vicinal diamines. This reaction protocol is particularly useful when the prior preparation and isolation of oligosaccharide acid or δ-lactone are problematic.

Carbohydrates play essential roles in living organisms. However, carbohydrate molecules are not easily tracked due to lack of a responsive chromophore. Thus, properly labeled saccharides are desirable for their composition analysis. Though fluorescent labels can be introduced to aldoses by reductive amination,¹³ this method still has limitations such as low reactivity and low yield on modification of oligosaccharides. Our current method using the iodine-promoted oxidative condensation reactions provides an alternative way to convert aldoses to highly fluorescent aldo-naphthimidazoles in a direct

(11) Lin, S.; Yang, L. *Tetrahedron Lett.* **2005**, *46*, 4315–4319.

(12) Maggio-Hall, L. A.; Dorrestein, P. C.; Escalante-Semerena, J. C.; Begley, T. P. *Org. Lett.* **2003**, *5*, 2211–2213.

(13) (a) Baues, R. J.; Gray, G. R. *J. Biol. Chem.* **1977**, *252*, 57–60. (b) Anumula, K. R. *Anal. Biochem.* **2006**, *350*, 1–23. (c) Larsen, K.; Thygesen, M. B.; Guillaumie, F.; Willats, W. T.; Jensen, K. J. *Carbohydr. Res.* **2006**, *341*, 1209–1234.

and efficient manner. Our preliminary study indicated that several aldo-naphthimidazoles derived from pentoses and hexoses were resolved by HPLC on reversed-phase columns (Figure 2). The detection limit was around 1 ppm. The analysis of aldo-naphthimidazoles can be further investigated by using capillary electrophoresis as a high-resolution and sensitive method.

Conclusion

Using iodine as an oxidant/promoter, various aldoses react readily with *o*-phenylenediamines or 2,3-naphthalenediamine in acetic acid solution to form the corresponding aldo-benzimidazoles and aldo-naphthimidazoles in high yields. Other functional groups such as hydroxyl, carboxyl, and amido groups are inert in the mild reaction conditions. Thus, the oxidative condensation reactions of various aldoses were realized without protection or modification of other functional groups. The success of this method relies on the crucial role of iodine and choice of acetic acid as the solvent. This protocol shows advantages over the previous methods^{3,5} by conducting the oxidative condensation reactions under mild conditions in a one-pot procedure to give high yields of aldo-imidazole products, which can be isolated simply by trituration of the crude reaction mixture with ethyl acetate. No racemization of saccharides or cleavage of the glycoside bonds occurs in our reaction protocol. In contrast to the starting materials of fluorescence-insensitive aldose and 2,3-naphthalenediamine, the aldo-naphthimidazole is highly fluorescent. Such property of enhanced fluorescence is potentially useful to the research of glycochemistry and glycobiology. In a preliminary study, we have demonstrated an application in carbohydrate composition analysis via the aldo-naphthimidazole derivatives.

Experimental Section

General Procedure for Oxidative Condensation of Aldoses with *o*-Phenylenediamines or 2,3-Naphthalenediamine. Method A. An appropriate aldose (1.0 mmol) and *o*-phenylenediamine (192 mg, 3.0 mmol) were dissolved in a solution containing acetic acid (1 mL) and water (7 mL). The mixture was stirred with a solution of iodine (254 mg, 1.0 mmol) in MeOH (2 mL) at room temperature until the aldose was completely consumed as indicated by the TLC analysis. During the reaction period, the deep brown color of reaction mixture remained. The reaction was quenched by addition of Na₂S₂O₃ (2 mL of saturated aqueous solution), and the mixture was concentrated under reduced pressure to give the crude products of aldo-benzimidazoles.

The crude aldo-benzimidazole product was treated with acetic anhydride (2 mL) and pyridine (2 mL) at 0–25 °C for 8 h and then partitioned between 1 N HCl (30 mL) and CH₂Cl₂ (50 mL). The organic phase was washed once with brine (30 mL), concentrated under reduced pressure, and purified by silica gel column chromatography (EtOAc/hexane, 1:2) to afford the desired product of aldo-benzimidazole peracetate.

Method B. The oxidative condensation reactions were also performed in buffer solution. Thus, an appropriate aldose (1.0 mmol) and *o*-phenylenediamine (2.0 mmol, 132 mg) was dissolved in acetic acid buffer solution (4 mL, pH 4.38). Iodine (1.0 mmol, 254 mg dissolved in 2 mL of methanol) was added, and the mixture was stirred at room temperature until the aldose was completely consumed as indication of the TLC analysis. The reaction mixture was quenched by Na₂S₂O₃, concentrated, and subjected to peracetylation with Ac₂O/pyridine to afford the desired product of aldo-benzimidazole peracetate.

Method C. A mixture of aldose (0.02 mmol), 2,3-naphthalenediamine (3.5 mg, 0.022 mmol), and iodine (0.5 mg, 0.002 mmol)

in AcOH (5.0 mL) was stirred at room temperature in open air. The reaction was complete in 6 h as indicated by the TLC analysis. The reaction mixture was triturated with EtOAc to give precipitates, which were collected by filtration using nylon membrane filter. The aldo-naphthimidazole products prepared as such was practically pure for characterization. This protocol is applicable to the reaction of larger scale, e.g., 2 mmol.

HPLC Analysis of Aldo-naphthimidazoles. For UV detection, the detection wavelength is set at 310 nm. For fluorescence detection, the excitation wavelength is 232 nm and the detection wavelength is 326 nm. The sample was eluted on a pair of HC-C₁₈ column (250 mm × 4.6 mm; 5 μm porosity) at 30 °C, and the retention time of sample components was recorded. The mobile phase is sodium phosphate buffer (100 mM, pH 5.0) containing 28% methanol and 2% acetonitrile at a flow rate of 0.8 mL/min.

(1'S,2'R,3'R,4'R)-1-Acetyl-5,6-dimethyl-2-(1,2,3,4,5-pentaacetoxy)-pentylbenzimidazole (3b-Ac). According to the general procedure (method A), D-glucose (180 mg, 1.0 mmol) and 4,5-dimethyl-1,2-benzenediamine (396 mg, 3.0 mmol) were stirred with iodine in aqueous AcOH solution for 11 h at room temperature. The crude product was subsequently treated with Ac₂O in pyridine to give compound **3b-Ac** (319 mg, 58% overall yield). C₂₆H₃₂N₂O₁₁: yellowish solid, mp = 83–85 °C; TLC (EtOAc/hexane, 1:2) R_f = 0.16; [α]_D²⁵ +63.1 (c 1.2, CHCl₃); IR ν_{max} (NaCl) 3456, 1752, 1367, 1222, 1036 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.50 (1 H, s), 7.33 (1 H, s), 6.60 (1 H, d, J = 3.6 Hz), 5.90 (1 H, dd, J = 5.6, 3.6 Hz), 5.55 (1 H, t, J = 5.6 Hz), 5.56–5.54 (1 H, m), 4.38 (1 H, dd, J = 12.4, 2.4 Hz), 4.26 (1 H, dd, J = 12.4, 5.6 Hz), 2.82 (3 H, s), 2.39 (3 H, s), 2.34 (3 H, s), 2.20 (3 H, s), 2.10 (3 H, s), 2.08 (3 H, s), 2.05 (3 H, s), 1.93 (3 H, s); ¹³C NMR (CDCl₃, 100 MHz) δ 169.9, 169.4, 169.2 (2 ×), 169.0, 168.9, 149.3, 140.5, 134.1, 133.1, 129.8, 120.8, 113.6, 69.4, 69.3, 69.2, 69.0, 61.6, 26.9, 21.2, 21.1 (2 ×), 21.0 (2 ×), 20.7, 20.3; HRMS (ESI) calcd for C₂₆H₃₃N₂O₁₁ 549.2079, found m/z 549.2061 [M + H]⁺.

(1'S,2'R,3'R,4'R)-2-(1,2,3,4,5-Pentahydroxyl)pentyl-1H-naphthimidazole (3g). According to the general procedure (method C), D-glucose (3.6 mg, 0.02 mmol) and 2,3-naphthalenediamine (3.5 mg, 0.022 mmol) were stirred with iodine (0.5 mg, 0.002 mmol) in 5.0 mL of AcOH for 6 h at room temperature. The crude product was triturated with EtOAc to give compound **3g** (6.2 mg, 98% yield). C₁₆H₁₈N₂O₅: brownish solid, mp = 175–177 °C; ¹H NMR (DMSO-*d*₆, 600 MHz) δ 8.06 (2 H, s), 7.94 (2 H, d, J = 6.2, 3.2 Hz), 7.46 (2 H, d, J = 6.2, 3.2 Hz), 5.12 (1 H, d, J = 5.3 Hz), 4.20 (1 H, dd, J = 5.3, 1.6 Hz), 3.59 (1 H, dd, J = 8.6, 1.3 Hz), 3.52–3.48 (2 H, m), 3.35 (1 H, dt, J = 11.9, 0.9 Hz); ¹³C NMR (DMSO-*d*₆, 150 MHz) δ 161.1 (2 ×), 139.5 (2 ×), 129.8 (2 ×), 127.9 (2 ×), 123.5 (2 ×), 110.9, 72.2, 71.6, 71.4, 70.4, 63.7; HRMS (ESI) calcd for C₁₆H₁₉N₂O₅ 319.1294, found m/z 319.1285 [M + H]⁺.

(1'S,2'R,3'R,4'R)-1-Acetyl-2-[1,2,4,5-tetraacetoxy-3-*O*-(2,3,4,5-tetraacetoxy-β-D-galactopyranosyl)]pentylbenzimidazole (14a-Ac). According to the general procedure (method B), D-lactose monohydrate (180 mg, 0.5 mmol) and *o*-phenylenediamine (108 mg, 1.0 mmol) were stirred with iodine (127 mg, 0.5 mmol) in aqueous AcOH buffer solution for 58 h at room temperature. The crude product was subsequently treated with Ac₂O in pyridine to give compound **14a-Ac** (359 mg, 88% overall yield) without further purification. C₃₆H₄₄N₂O₁₉: yellowish foam; TLC (EtOAc/hexane, 1:1) R_f = 0.12; [α]_D²⁵ +40.3 (c 2.7, CHCl₃); IR ν_{max} (NaCl) 3443, 1750, 1371, 1227, 1047 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz) δ 7.74–7.72 (1 H, m), 7.66–7.63 (1 H, m), 7.39–7.31 (2 H, m) 6.63 (1 H, dd, J = 2.8, 0.8 Hz), 5.74 (1 H, m), 5.34 (1 H, d, J = 3.6 Hz), 5.22–5.19 (1 H, m), 5.13 (1 H, dd, J = 10, 8 Hz), 5.00 (1 H, m), 4.63 (1 H, m), 4.41 (1 H, t, J = 5.6 Hz), 4.21 (1 H, dd, J = 10.8, 7.2 Hz), 4.02–3.92 (2 H, m), 3.84 (1 H, t, 6.8), 2.88 (3 H, s), 2.27 (3 H, s), 2.14 (3 H, s), 2.13 (3 H, s), 2.09 (3 H, s), 2.06 (3 H, s), 2.00 (3 H, s), 1.98 (3 H, s), 1.93 (3 H, s); ¹³C NMR (CDCl₃, 100 MHz) δ 169.9, 169.8, 169.7 (2 ×), 169.6 (2 ×), 169.3, 169.2, 168.8, 151.1, 142.1, 131.7, 125.0, 124.2, 120.5, 113.4, 100.9, 70.7, 70.6, 70.3, 69.4, 69.3, 68.8, 66.6, 61.3, 60.7, 60.1, 26.5, 20.8, 20.7, 20.6 (2 ×),

20.4 (2×), 20.3, 20.2; HRMS (ESI) calcd for C₃₆H₄₅N₂O₁₉ 809.2613, found *m/z* 809.2611 [M + H]⁺.

Compound 16a Derived from Maltotriose. According to the general procedure (method C), D-maltotriose monohydrate (50.4 mg, 0.1 mmol) and *o*-phenylenediamine (21.6 mg, 0.2 mmol) were stirred with iodine (25 mg, 0.1 mmol) in 3.0 mL of AcOH for 30 h at room temperature. The crude product was purified by C18 reversed-phase silica gel column chromatography (MeOH/H₂O, 1% to 30% as gradient) to afford the desired product **16a** (30 mg, 51% yield). C₂₄H₃₆N₂O₁₅: yellowish foam; TLC (acetone/EtOAc/H₂O/AcOH, 60:30:20:1) *R_f* = 0.44; [α]_D²⁵ +55.40 (*c* 1.0, H₂O); ¹H NMR (D₂O, 400 MHz) δ 7.62–7.60 (2 H, m), 7.32–7.30 (2 H, m), 5.37 (1 H, d, *J* = 4.0 Hz), 5.21 (1 H, d, *J* = 4.4 Hz), 5.09 (1 H, d, *J* = 3.6 Hz), 4.34–3.37 (17 H, m) ¹³C NMR (CD₃OD, 150 MHz) δ

154.9, 121.2 (4×), 115.3 (2×), 100.7, 100.2, 81.2, 77.4, 73.8, 73.6, 73.3, 73.0, 72.2, 71.9, 71.6, 71.3, 69.7, 69.0, 62.6, 60.8, 60.4; the assignments of proton and carbon resonances were supported by the HSQC spectrum; HRMS (ESI) calcd for C₂₄H₃₇N₂O₁₅ 593.2188, found *m/z* 593.2190 [M + H]⁺.

Acknowledgment. We thank the National Science Council for financial support.

Supporting Information Available: Synthetic procedures, physical and spectroscopic data of products; ¹H and ¹³C NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO800234X