

BRIEF COMMUNICATION

The relation between accumulation of abscisic acid and proline in detached rice leaves

C.-W. YANG, J.W. WANG and C.H. KAO*

Department of Agronomy, National Taiwan University, Taipei 106, Taiwan, Republic of China

Abstract

The relation between abscisic acid (ABA) and proline accumulation was investigated in detached rice (*Oryza sativa* L.) leaves. In darkness, proline content increased about 2-, 2.5- and 6-fold after 24, 48 and 72 h. ABA content reached maximum after 48 h. In the light, proline content remained almost unchanged until 48 h and subsequently increased slightly. ABA content in the light was lower than in darkness, but the maximum was also after 48 h. During 12-h exposure to decreased air humidity, proline content gradually increased, but ABA content increased about 25-fold after 4 h and declined thereafter. Exogenous application of ABA resulted in an increase in proline content in detached rice leaves under both light and darkness.

Additional key words: *Oryza sativa*, water stress.

It is well established that proline accumulates in plant tissues under water stress, salt stress, and dark-induced senescence (e.g. Kao 1981, Moftah and Michel 1987, Munns *et al.* 1979, Stewart and Voetberg 1985, Venekamp and Kost 1988, Wang *et al.* 1982). Proline accumulation can be induced by abscisic acid (ABA) in plant tissues (Aspinall *et al.* 1973, Chou *et al.* 1991, Pesci 1987, Stewart 1980, Stewart and Voetberg 1985). However, the lack of proline accumulation in response to ABA treatment has also been reported for spinach, pearl millet, tobacco and sunflower leaves (Aspinall and Paleg 1981, McDonnell *et al.* 1983). ABA is known to increase rapidly not only in intact but also in detached leaves under water stress (Hsiao 1973). In darkness, it has been shown that the content of ABA in leaves of oats and rice increased during dark-induced senescence (Gepstein and Thimann 1980, Wang *et al.* 1999). Therefore, proline accumulation in response to water stress or dark-induced senescence may be a response to accumulated ABA. In fact, Ober and Sharp (1994) reported that increased content of ABA is required for proline accumulation in maize primary roots at low water potentials. However,

proline accumulation in the absence of ABA accumulation has also been reported (Stewart and Voetberg 1985, 1987).

Previously, we have reported that proline accumulates in detached rice leaves during dark-induced senescence (Kao 1981, Wang *et al.* 1982) or exposed to water stress (Kao 1981). In this paper, we have examined the relation between accumulation of ABA and proline in detached rice leaves.

Rice (*Oryza sativa* L. cv. Taichung Native 1) seedlings were planted on a stainless net floating on modified half-strength Johnson's nutrient solution in a 500-cm³ beaker (Kao 1980). The nutrient solution (pH 4.5) was replaced every 3 d. Rice seedlings were grown in a greenhouse with natural daylight at day/night temperature 30/25 °C and relative humidity (RH) about 95 % (for detail see Kao 1980). The apical 3 cm segments were excised from the third leaves of 12-d-old seedlings. Groups of twenty leaf segments were exposed to slightly decreased RH above a solution of 0.5 M NaCl. Under this condition, relative water content of leaf segments after 4, 8, and 12 h was 80, 62, and 60 %, respectively (Yang *et al.* 2000).

Received 27 July 1999, accepted 1 November 1999.

Abbreviations: ABA - abscisic acid; f. m. - fresh mass.

Acknowledgement: This work was supported by the National Science Council, the Republic of China, grant 89-2313-B-002-111.

* Author for correspondence: fax: (+886)-2-2362-0879, e-mail: kaoch@ccms.ntu.edu.tw

Similar segments of leaves were floated on 20 cm³ of distilled water or a solution of 45 μM abscisic acid (ABA) in a Petri dish to serve as controls and ABA-treated samples, respectively. All samples were incubated




Fig. 1. Changes in proline content of detached rice leaves floated on distilled water or 45 μM ABA solution under dark or light for 12 h. Means ± SE, n = 4.

at 27 °C in darkness or under irradiance 40 μmol m⁻² s⁻¹. Proline and ABA were extracted and quantified as described previously (Kao 1981, Chen and Kao 1993). For proline utilization, detached rice leaves were pretreated with 50 mM ornithine [since addition of ornithine has been observed to be more effective than that of glutamic acid or arginine in increasing proline content in rice leaves (Yang *et al.* 2000)] to increase the




Fig. 2. Changes in proline and ABA contents of detached rice leaves floated on distilled water under dark or light for 72 h. Means ± SE, n = 4.

endogenous proline content and then transferred to distilled water, 45 μM ABA, or decreased humidity for 4, 8, and 12 h in the light. Change in proline content were then monitored. The decline in proline content will be considered as that proline is utilized (Yang *et al.* 2000). All data were expressed on the basis of initial fresh mass.

All experiments were repeated three times; within each experiment, treatments were replicated 4 times. Similar results and identical trends were obtained in all experiments. The data reported here are from a single experiment.

To establish the relation between accumulation of ABA and proline in detached rice leaves, we have to know whether proline accumulation can be induced by ABA. Under both light and darkness, ABA treatment indeed resulted in an accumulation of proline in detached rice leaves (Fig. 1).

In darkness, proline content in detached rice leaves incubated in distilled water for 24, 48 and 72 h, increased about 2-, 2.5- and 6-fold, respectively (Fig. 2), which confirmed our previous results (Wang *et al.* 1982). Proline content, however, in detached rice leaves incubated in the light remained relatively unchanged after 48 h and subsequently increased slightly (Fig. 2). It is apparent that proline accumulates more in darkness than in the light in detached rice leaves. In darkness, ABA content did not increase 24 h after incubation, but increased 48 h and remained unchanged 72 h (Fig. 2). ABA content in detached rice leaves incubated in the light increased in a similar pattern but to a lesser extent than in darkness (Fig. 2). It is clear that the increase in ABA content in detached rice leaves in the dark was observed after proline accumulation.




Fig. 3. Changes in proline and ABA contents of detached rice leaves floated on distilled water (H₂O) or exposed to slightly decreased RH above 0.5 M NaCl (WS; RWC 80, 62, and 60 % after 4, 8, and 12 h, respectively) in the light. Means ± SE, n = 4.

In control leaves, proline and ABA contents remained relatively unchanged during 12 h of incubation in the light. Treatment with decreased humidity resulted in a significant increase in proline content in the light (Fig. 3). ABA content in detached rice leaves incubated at decreased humidity increased rapidly from a control level of about 0.1 to over 2.5 nmol g⁻¹ (f. m.) in a period of 4 h, and declined thereafter (Fig. 3). Similar time course has

been observed previously for pearl millet, rice and barley (Henson and Quarrie 1981, Stewart and Voetberg 1985). It is clear that ABA content reached maximum at the time when proline accumulated slightly.




Fig. 4. Changes in proline content in detached rice leaves pretreated with 50 mM ornithine for 3 h in the light and then floated on distilled water (H₂O), 45 μM ABA solution (ABA), or exposed to decreased RH (WS) in the light. Means ± SE, n = 4.

References

- Aspinall, D., Paleg, L.G.: Proline accumulation: physiological aspects. - In: Paleg, L. G., Aspinall, D. (ed.): *Physiology and Biochemistry of Drought Resistance in Plants*. Pp. 206-242. Academic Press, Sydney 1981.
- Aspinall, D., Singh, T.N., Paleg, L.G.: Stress metabolism V. Abscisic and nitrogen metabolism in barley and *Lolium temulentum* L. - *Aust. J. Biol. Sci.* **26**: 319-329, 1973.
- Chen, C.T., Kao, C.H.: Osmotic stress and water stress have opposite effects on putrescine and proline production in excised rice leaves. - *Plant Growth Regul.* **13**: 197-202, 1993.
- Chou, I.T., Chen, C.T., Kao, C.H.: Characteristics of the induction of the accumulation of proline by abscisic acid and isobutyric acid in detached rice leaves. - *Plant Cell Physiol.* **32**: 269-272, 1991.
- Gepstein, S., Thimann, K.V.: Changes in abscisic acid content of oat leaves during senescence. - *Proc. nat. Acad. Sci. USA* **77**: 2050-2053, 1980.
- Henson, I.E., Quarrie, S.A.: Abscisic acid accumulation in detached cereal leaves in response to water stress I. Effects of incubation time and severity of stress. - *Z. Pflanzenphysiol.* **101**: 431-438, 1981.
- Hsiao, T.C.: Plant response to water stress. - *Annu. Rev. Plant Physiol.* **24**: 519-570, 1973.
- Kao, C.H.: Senescence of rice leaves IV. Influence of benzyladenine on chlorophyll degradation. - *Plant Cell Physiol.* **21**: 1255-1262, 1980.
- Kao, C.H.: Senescence of rice leaves VI. Comparative study of the metabolic changes of senescing turgid and water-stressed excised leaves. - *Plant Cell Physiol.* **22**: 683-688, 1981.
- McDonnell, E.M., Coughlan, S.J., Wyn Jones, R.G.: Differential effects of abscisic acid on glycinebetaine and proline accumulation in three plant species. - *Z. Pflanzenphysiol.* **109**: 207-213, 1983.
- Mofiah, A.E., Michel, B.E.: The effect of sodium chloride on solute potential and proline accumulation in soybean leaves. - *Plant Physiol.* **83**: 238-240, 1987.
- Munns, R., Brady, C.J., Barlow, F.W.: Solute accumulation in the apex and leaves of wheat during water stress. - *Aust. J. Plant Physiol.* **6**: 379-389, 1979.
- Ober, E.S., Sharp, R.E.: Proline accumulation in maize (*Zea mays* L.) primary root at low water potentials. - *Plant Physiol.* **105**: 881-987, 1994.
- Pesci, P.: ABA-induced proline accumulation in barley leaf segments: dependence on protein synthesis. - *Physiol. Plant.* **71**: 287-291, 1987.
- Stewart, C.R.: The mechanism of abscisic acid-induced proline accumulation in barley leaves. - *Plant Physiol.* **66**: 230-233, 1980.
- Stewart, C.R., Voetberg, G.: Relationship between stress-induced ABA and proline accumulations and ABA-induced proline accumulation in excised barley leaves. - *Plant Physiol.* **79**: 24-27, 1985.
- Stewart, C.R., Voetberg, G.: Abscisic acid accumulation is not required for proline accumulation in wilted leaves. - *Plant Physiol.* **83**: 747-749, 1987.
- Venckamp, J.H., Kost, J.F.M.: The sources of free proline and asparagine in field bean plants, *Vicia faba* L., during and after a short period of water withholding. - *J. Plant Physiol.* **132**: 102-109, 1988.
- Wang, C.Y., Cheng, S.H., Kao, C.H.: Senescence of rice leaves

- VII. Proline accumulation in senescing excised leaves. - *Plant Physiol.* **69**: 1348-1349, 1982.
- Wang, J.-W., Kao, C.H.: Endogenous abscisic acid levels are not linked to methyl jasmonate-promoted senescence of detached rice leaves. - *Plant Growth Regul.* **28**: 17-20, 1999.
- Yang, C.-W., Lin, C.C., Kao, C.H.: Endogenous ornithine and arginine contents and dark-induced proline accumulation in detached rice leaves. - *J. Plant Physiol.*, in press, 2000.
- Yang, C.-W., Lin, C.C., Kao, C.H.: Proline, ornithine, arginine and glutamic acid contents in detached rice leaves. - *Biol. Plant* **43**: 305-308, 2000.
- Yoshida, K., Kiyosue, T., Nakashima, K., Yamaguchi-Shinozaki, K., Shinozaki, K.: Regulation of levels of proline as an osmolyte in plants under water stress. - *Plant Cell Physiol.* **38**: 1095-1102, 1997.