

行政院國家科學委員會專題研究計畫成果報告

計畫名稱：肌肉缺血再灌流損傷分子機轉 (2/2)

計畫編號：NSC 90-2314-B-002-411-

執行期限：90年8月1日至91年12月31日

主持人：簡雄飛 執行機構及單位名稱：台大醫學院外科

一、中文摘要

肌肉缺血再灌流傷害會造成肌肉移植後肌肉功能不全或因血管阻塞後再灌流造成 compartment syndrome。再灌流本身可能比缺血的傷害還大，這是因缺血時所產生的 xanthine oxidase，再灌流後在有氧的情況下代謝 ATP 分解產物 xanthine 與 hypoxanthine，產生氧自由基，雖然一些去氧自由基的藥物如 superoxide dismutase 及 allopurinol 可減緩一些傷害，但研究者相信中性球粘黏於血管壁，在缺血處產生更多的氧自由基及傷害，是此病態生理中重要的過程。我們在大鼠大腿處以 tourniquet 造成 4 小時缺血，再以 *in vivo* MCLA 冷光測定儀測定不同時間再灌流所造成 oxygen free radicals (ORF) 量，發現所測得最高量在第二天，此時以抗中性球抗體(MCA967)染出最高量的中性球數，以及以 TUNEL 染出最多的肌肉細胞凋亡。故我們應尋求較長期(>2d)有效的 ORF 清除劑，尤其是中性球的聚集因子(chemokine)的去除，才可有效的避免肌肉缺血再灌流傷害，目前初步資料顯示，以抗體阻斷聚集因子的確可減少中性球浸潤及 ORF 產生。

關鍵詞：muscle; oxygen free radicals; ischemia reperfusion injury

Abstract

We studied the ischemia reperfusion injury of skeletal muscle by oxygen free radicals (OFRs) detection with MCLA chemiluminescence, neutrophil staining and TUNEL. The OFRs as well as neutrophils and muscle cell apoptosis peaked by day 2 after release of tourniquet. These findings are interesting because the neutrophil number and the muscle cells showing apoptosis are also peaking on day 2. Neutrophils are likely the major source of free radicals. The real-time oxygen free radical monitoring in the reperfused skeletal muscle was shown peaked on the 2nd day. This piece of information is important since the regimen decreasing the OFRs must be prolonged after 2nd day. Otherwise the clinical result must be poor. The source of this OFR was under investigated by neutrophil staining and real-time PCR of CINC (the chemokine for neutrophils of rats), which were shown to be correlated. Besides, the blocking antibodies to CINC could effectively decrease neutrophil infiltration and OFRs. However the data was only preliminary.

Keywords: muscle; oxygen free radicals; ischemia reperfusion injury

二、Introduction

Reperfusion after a transient ischemia often causes greater injury to the affected organs than ischemia itself[1]. Upon reperfusion, increased amount of xanthine oxidase metabolize xanthine and hypoxanthine, which are metabolites of ATP during

ischemia, to produce oxygen free radicals (OFRs). Although superoxide dismutase (free radical scavenger) or allopurinol (xanthine oxidase inhibitor) are partially effective in reducing injury at this early stage, it is clear that neutrophils adhered to the postcapillary venules release even more OFRs and destructive proteases.

The microvascular endothelium plays an important role in initiating the events associated with IRI by releasing chemotactic factors[2;3]. Many cells are known to produce chemokines after being stimulated with pro-inflammatory stimuli. These chemokines could recruited different subsets of leukocytes as needed. However, in IRI, too many activated neutrophils accumulated into ischemic tissue might turn out to be harmful. There were reports about the expression of cytokine-induced neutrophil chemoattractant (CINC), which attract neutrophils in rat brain IRI[4]. The chemokines in skeletal muscle IRI are still unknown. We had experience dissecting the chemokine profile (Two C-X-C chemokines: KC and IP-10, three C-C chemokines: MCP-1, MIP-1 alpha, and RANTES) in degenerating peripheral nerves. Hypothesizing that CINC mRNA might increase early in the course of IRI of rat skeletal muscle, we tested for the expression of CINC mRNA by quantitative reverse transcription-coupled PCR (RT-PCR), *in situ* hybridization, and for the presence of the CINC protein and neutrophils by immunocytochemistry.

There are many ways of measuring OFRs, but all have to deal with their short half-lives. The most common form of measurement are those of the products of OFRs attack on lipid, such as malondialdehyde (MDA). The thiobarbituric acid (TBA) test for MDA is non-specific. Many data of this method are difficult to interpret and compare. With the advent of very sensitive photon counting technique, ultraweak chemiluminescence (CL) from OFRs and enhanced with luminol, leucigenin or MCLA could be easily detected. A sensitive both *in vitro* and *in vivo* CL detecting system by Tohoku Co. Japan were recently available at our common lab. We would use a rat hind limb ischemia model by tourniquet to create 4 h ischemia, followed by various time of reperfusion.

The CL readings from exposed medial gastrocnemius muscle *in vivo* will be correlated with the following parameters to test the hypotheses: 1) OFRs are the main source of damage, and 2) the main source of OFRs is the neutrophils. The other three parameters are 1) tissue viability as stained with nitroblue tetrazolium, 2) neutrophils density within the

venules or outside the venules as stained with anti-neutrophil antibody (MCA 967, Serotec, UK), 3) apoptosis as studied with DNA ladder on gel electrophoresis and TUNEL reaction. And lastly 4) by applying CINC neutralized antibodies, the neutrophil response and apoptosis and CL readings are compared between experimental and control groups.

三、Materials and Methods

With the advent of very sensitive photon counting technique, ultraweak chemiluminescence (CL) from OFRs and enhanced with MCLA could be easily detected. We used a rat hind limb ischemia model by tourniquet to create 4 h ischemia, followed by various time of reperfusion. The CL readings from exposed medial gastrocnemius muscle *in vivo* will be correlated with the following parameters to test the hypotheses: 1) OFRs are the main source of damage, and 2) the main source of OFRs is the neutrophils. The other parameters are 1) neutrophils density within the venules or outside the venules as stained with anti-neutrophil antibody (MCA 967, Serotec, UK), 2) apoptosis as studied with DNA ladder on gel electrophoresis and TUNEL reaction. Besides we used anti-neutrophil chemokine antibodies to block the chemokine induction after ischemia/reperfusion with 1 mg anti-CINC (Chemicon, USA) [5].

四、Results

1. Animal model

We created this model by catheterizing the iliac arteries via either contralateral femoral artery or ipsilateral jugular artery till the bifurcation of iliac arteries, where the enhancer MCLA was injected totally into the ischemic side. The constant readings were got due to this method as well as accurate dose of MCLA.

2. MCLA Chemiluminescence detection

By applying *in vivo* continuous measuring of MCLA chemiluminescence, we accidentally found the peak was not in the early phase of reperfusion, but rather at 2 day. This is why most of the OFR scavengers work not so good, since most of the drug were used in the early phase.




Fig.1 Intermittent CL measurement of gastrocnemius muscle in a representative animal after ischemia for 4 h and reperfused for various time periods. MCLA was infused at a speed of 0.125 mg/hr. The peaked measurement was on day 2.

3. Neutrophil and TUNEL staining

With fresh tissue on cryostat section, we developed neutrophil staining using Serotec's MCA967 antibodies. With paraformaldehyde fixed tissue, TUNEL was done, constantly showing apoptotic muscle cells, especially at 2 day.




Fig.2 Upper panel showed the neutrophil- antibody immunoreactive cells on day 1 and day 2. Neutrophils were less on day 1, day 3 or day 4 (not shown) when compared with that of day 2. Lower panel showed TUNEL positive nuclei of muscle cells on day 1 and day 2. There were more neutrophil infiltration and apoptotic muscle cells on day 2.

4. Anti-CINC antibodies neutralization

The preliminary data shown the blocking antibodies to CINC could effectively decrease neutrophil infiltration and OFRs. The apoptosis index of muscle is also shown to be less.

四. 討論及自我評估

The real-time oxygen free radical monitoring in reperfused skeletal muscle was done here, which to our surprise, the OFR peaked on the 2nd day. This piece of information is important since the regimen decreasing the OFRs must be prolonged after 2nd day. Otherwise the clinical result must be poor. The source of this OFR was under investigated by neutrophil staining and real-time PCR of CINC, which was preliminarily shown to be correlated. Besides the blocking antibodies to CINC could effectively decrease neutrophil infiltration and OFRs. The apoptosis index of muscle is also shown to be less. Statistical evaluation is under way. These piece of

information is valuable if we can minimize the ischemia/reperfusion injury by simply decreasing neutrophil response through blocking its chemokine.

五、参考文献

Reference List

- [1] Granger DN. Role of xanthine oxidase and granulocytes in ischemia-reperfusion injury. [Review] [46 refs]. *Am J Physiol* 1988; 255(6 Pt 2):H1269-H1275.
- [2] Korthuis RJ, Granger DN. Reactive oxygen metabolites, neutrophils, and the pathogenesis of ischemic-tissue/reperfusion. [Review] [92 refs]. *Clinical Cardiology* 1993; 16(4 Suppl 1):I19-I26.
- [3] Formigli L, Lombardo LD, Adembri C, Brunelleschi S, Ferrari E, Novelli, GP. Neutrophils as mediators of human skeletal muscle ischemia-reperfusion syndrome. *Hum Pathol* 1992; 23(6):627-634.
- [4] Liu T, Young PR, McDonnell PC, White RF, Barone FC, Feuerstein GZ. Cytokine-induced neutrophil chemoattractant mRNA expressed in cerebral ischemia. *Neurosci Lett* 1993; 164(1-2):125-128.
- [5] Hirose J, Yamaga M, Kato T, Ikebe K, Takagi K. Effects of a hydroxyl radical scavenger, EPC-K1, and neutrophil depletion on reperfusion injury in rat skeletal muscle. *Acta Orthop Scand* 2001; 72(4):404-410.