

NEUROLOGY

Teaching NeuroImage: Cervical cord atrophy with dorsal root ganglionopathy in Sjögren syndrome

Chih-Chun Lin and Ming-Jang Chiu

Neurology 2008;70:e27

DOI: 10.1212/01.wnl.0000297221.51829.db

This information is current as of October 9, 2009

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://www.neurology.org/cgi/content/full/70/7/e27>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2008 by AAN Enterprises, Inc. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.




Teaching *NeuroImage*: Cervical cord atrophy with dorsal root ganglionopathy in Sjögren syndrome

Chih-Chun Lin, MD
Ming-Jang Chiu, MD,
PhD

Address correspondence and
reprint requests to Dr. Ming-
Jang Chiu, Department of
Neurology, National Taiwan
University Hospital, No. 7,
Chung-Shan S. Rd., Taipei, 100,
Taiwan
mjchiu@ntu.edu.tw

Figure MRI showing increased signal intensity at the dorsal column (A) and atrophy (B) of the cervical cord on T2-weighted images



The axial image is at the level of C3/C4 and the cord atrophy extends from C2 to C6.

A 59-year-old woman had had asymmetric numbness from her neck to both arms and left leg for 5 years. She had had dry eyes and mouth for many years. Examination showed sensory ataxia, generalized areflexia, positive Romberg test, and hand pseudoathetosis. Vibration sense was reduced in both lower limbs; the severity was more on the left side but was not length dependent. Joint position sense was impaired in all limbs while pain and temperature senses were comparatively preserved. No sensory level was demarcated. Electrophysiologic studies revealed absence of sensory action potentials in all sampled nerves and prolonged central conduction time from median nerves on somatosensory-evoked potentials.^{1,2} Her anti-SSA antibody,

rheumatoid factor, and Schirmer test were positive. MRI (figure) showed cervical cord atrophy with increased T2 signal intensity at the dorsal column which implied a consequence of centripetal Wallerian degeneration from dorsal root ganglionopathy in a patient with Sjögren syndrome.

REFERENCES

1. Malinow K, Yannakakis GD, Glusman SM, et al. Subacute sensory neuronopathy secondary to dorsal root ganglionitis in primary Sjögren's syndrome. *Ann Neurol* 1986;20:535-537.
2. Lauria G, Pareyson D, Sghirlanzoni A. Neurophysiological diagnosis of acquired sensory ganglionopathies. *Eur Neurol* 2003;50:146-152.

From the Departments of Neurology, National Taiwan University Hospital, College of Medicine, National Taiwan University, Taipei, Taiwan.

Disclosure: The authors report no conflicts of interest.

Teaching NeuroImage: Cervical cord atrophy with dorsal root ganglionopathy in Sjögren syndrome

Chih-Chun Lin and Ming-Jang Chiu

Neurology 2008;70:e27

DOI: 10.1212/01.wnl.0000297221.51829.db

This information is current as of October 9, 2009

Updated Information & Services

including high-resolution figures, can be found at:
<http://www.neurology.org/cgi/content/full/70/7/e27>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

MRI

<http://www.neurology.org/cgi/collection/mri> **Autoimmune diseases**

http://www.neurology.org/cgi/collection/autoimmune_diseases **All Spinal Cord**

http://www.neurology.org/cgi/collection/all_spinal_cord **Evoked Potentials/Somatosensory**

http://www.neurology.org/cgi/collection/evoked_potentials-somatosensory

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.neurology.org/misc/Permissions.shtml>

Reprints

Information about ordering reprints can be found online:

<http://www.neurology.org/misc/reprints.shtml>

