

# 行政院國家科學委員會專題研究計畫 成果報告

胃癌致病機轉：以 cytokine 和 chemokine 基因單核苷酸多型  
性探討宿主敏感性角色(3/3)

計畫類別：整合型計畫

計畫編號：NSC93-2321-B-002-031-

執行期間：93年08月01日至94年07月31日

執行單位：國立臺灣大學醫學院一般醫學科

計畫主持人：吳明賢

報告類型：完整報告

處理方式：本計畫可公開查詢

中 華 民 國 94 年 12 月 19 日

# 行政院國家科學委員會專題研究計畫成果報告

胃癌致病機轉：以cytokine和chemokine基因單核苷酸多型性探討宿主敏感性角色(3/3)

Gastric carcinogenesis: the role of host susceptibility studied by single nucleotide polymorphisms of cytokine and chemokine genes (3/3)

計畫類別：個別型計畫 整合型計畫

計畫編號： NSC92-2314-B-002-122

執行期間： 93年 8月 1日至 94年 7月 31日

整合型計畫：總計畫主持人：林肇堂醫師  
子計畫主持人：吳明賢醫師

註：整合型計畫總報告與子計畫成果報告請分開編印各成一冊，彙整一起繳送國科會

處理方式：可立即對外提供參考  
一年後可對外提供參考  
兩年後可對外提供參考  
(必要時，本會得展延發表時限)

執行單位：台大醫學院內科

中華民國 94 年 8 月 26 日

## 中文摘要

胃癌為全世界第二常見惡性腫瘤，在台灣則每年有 2000 以上患者死於胃癌，佔癌症死因第四位。胃癌致病機轉複雜且主要經由遺傳敏感性和環境因子交互作用。過去多數研究著重於胃癌環境危險因子，有不少報告指出幽門桿菌感染和抽煙為主要致病因素，但是只有少數暴露這些因子的人得到胃癌，而且最近新的証據也顯示個體的遺傳因子也佔重要角色，更精確而言是宿主對環境因素的反應，而非細菌和環境因子本身，是造成較易或較不易發生胃癌的主因。

在胃癌的多步驟改癌過程中，胃發炎是胃癌發生的必經之路。因此誘使和調控發炎反應的因子可能在預後的決定上有關鍵作用，而遺傳因子和其它的環境因子也可能是透過胃炎的層次來影響。在此觀點中，cytokines 和 chemokines 由於本身在微生物免疫和發炎反應有一定角色，是最有可能控制胃炎預後的主要因子。故假設不同促發炎和抗發炎 cytokines 間的平衡決定了胃炎的後果，而胃癌可能是由不同 cytokine 基因的表現增加和減少組合成特定的 cytokine 環境下產生。

幽門桿菌和抽煙皆能誘發胃的 cytokine 和 chemokine 表現。而不同個體間 cytokine 產生能力差異頗大。這種個體差異主要是由於 cytokine 基因上的單核苷酸多形性(SNPs)造成。因此，個體在基因(或 transcription)層次上所導致的不同 cytokine 反應或許可解釋對感染或環境毒素的個體敏感性的差異。這些發現促使吾人得以探討 cytokines, chemokines 和其受体等免疫調控主要分子的 SNPs 在個人胃癌相對危險的角色。最近兩個研究也指出 IL-1 $\beta$  的 SNPs 與胃癌危險性有關。雖然如此，由於基因型受種族影響，這些研究必須考慮種族的差異性而有所不同，而且基因—基因和基因—環境的交互作用在這些研究中也付之闕如。

在第一年的計劃裏，吾人收集 220 例胃癌及 230 例健康對照，抽取白血球 DNA，分析 TNF- $\alpha$ ，IL-1，IL-4 和 IL-10 等基因上的 11 處多型性(表一)，並將基因型與 H. pylori 感染和抽煙等環境因子比較，結果顯示 H. pylori 感染 [odds ratio (OR) 1.7, 95% confidence interval (CI) 1.19~2.56]，抽煙 (OR:2.02, 95% CI 1.38~2.95) 和高 IL-10 製造基因型 (OR:2.67, 95%ci 1.29~5.50) 明顯增加胃癌發生的危險性。進一步分析不同亞型發現瀰漫型胃癌 (OR:1.64, 95%CI 1.01~2.61) 或賁門癌 (OR:2.44, 95%CI 1.13~2.67) 與

IL-4 (-590)的 CT/CC 基因型有關，而高 IL-10 基因型則與賁門癌(OR:3.21, 95% CI 1.06~9.73)或進行性胃癌(OR:2.29, 95%CI 1.14~2.52)有關，至於 IL-1 和 TNF- $\alpha$  基因型則和胃癌危險性無關。以迴歸分析則顯示 H. pylori 感染(OR:1.7, 95%CE 1.14~2.52)，抽煙(OR:1.81, 95%CI 1.27~2.96)和 IL-10 基因型(OR:2.54, 95%CI 1.24~5.61)為胃癌發生的獨立危險因子。上述結果明白顯示胃癌的發生受到宿主和環境因子等共同影響。

在第二年的計劃裏，利用 cDNA microarray 分析不同菌株(潰瘍、胃癌、淋巴瘤)對上皮細胞的基因表現影響，初步已發現一些重要基因的改變在不同幽門螺旋桿菌疾病有所不同。另外，持續收集的病案與對照例數已超過 250 對。而在已知基因多型性的分析方面，吾人則進行 TNF- $\alpha$  (-238, -308, -857, -863, -1031)，TNF- $\alpha$  受器基因 TNFR1(-383)及 TNFR2(codon 196)，CD14(-159)與 Toll-like receptor 4(TLR4 Asp299Gly, Thr399Ile)，和 IL-8(-251)等基因多型性的對照病例研究，發現 TNF- $\alpha$  -857C→T 與胃黏膜相關淋巴瘤的危險性有關(OR=0.33, 95%CI: 0.15~0.75)，但與胃癌則並無相關性，這些結果顯示胃癌與胃黏膜相關淋巴瘤致病機轉上之明顯不同。

在第三年的計劃裏，吾人已收集至少 300 例不同表現型胃癌的病例以做為進一步分析之用，並利用這些病例與健康對照進行 IL-6 -634 C/G 及 IL-18 -137 G/C，-607 C/A 基因型的分析。另外，吾人利用 cDNA microarray 之，初步也發現一些具 biologic significance 且為 novel 的 candidate gene，這些基因多形性於宿主敏感性之角色，值得進一步研究。由於吾人在此方面的成果不錯，因此也將 host 及 environment 彼此影響的觀念，受邀寫成 review 於 Cancer Epidemiol Biomark & Prev 發表。

關鍵詞：胃癌，幽門螺旋桿菌，基因型，宿主易感性

## Abstract

Gastric cancer (GC) is the second most common malignancy in the world. In Taiwan, GC remains the fourth leading cause of cancer death with more than 2000 persons dying of this devastating disease annually. The pathogenesis of GC is complex and consists of two interacting elements: genetic susceptibility and environment factors. In the past decades, much attention has been paid to the environmental risks of GC. Considerable evidence has pointed out that *H. pylori* infection and smoking are among the major determinants of GC. However, only a small portion of patients exposed to environmental stimuli develop GC. Recently, new evidence has emerged suggesting that host genetic factors also play a critical role. Specifically, the host response to environmental triggers, rather than bacteria or environmental factors *per se*, may confer susceptibility to or protection against GC.

In the multistage model of gastric carcinogenesis, gastric inflammation is a prerequisite for the development of GC. Therefore, factors involved in initiation and regulation of the inflammatory responses may play a pivotal role in determining outcomes and genetic factors, like various *H. pylori* strains or environmental toxins, may operate at the level of gastric inflammation. In this respect, the intrinsic role of cytokines and chemokines as modulators of anti-microbial immunity and inflammatory responses makes them attractive candidates as a key regulator influencing the outcome of gastritis. It was assumed that the critical balance between pro- and anti-inflammatory cytokines would dictate the outcomes of gastric inflammation and development of GC might be facilitated in a permissive cytokine environment by combination of different unregulated or down regulated cytokine gene expression.

Both *H. pylori* infection and cigarette smoking could induce cytokine and chemokine expression in the stomach. The *in vitro* maximal capacity to produce different cytokines varies among different individuals. Such inter-individual differences can be attributed to single nucleotide polymorphisms (SNPs) in the coding or promoter regions of cytokine genes. SNPs may affect the overall expression and secretion of cytokines. Therefore, different host cytokine responses determined at the genetic (transcription) level may explain why there is a variety of individual susceptibility to a given microbial infection or environmental toxins. These developments allow us the opportunity to investigate the role of SNPs in key immunoregulatory molecules-cytokines, chemokines, and their receptors on the relative risk of GC for a given individual. Two recent studies have indeed provided evidence that SNPs of IL-1  $\beta$  are associated with the risk of GC. However, such study was performed mainly in Caucasians and the

genotype status varies greatly between different ethnic backgrounds. Furthermore, studies concerning gene-environment and gene-gene interactions remain lacking.

In the first year grant period, we analyzed 11 functional polymorphisms in tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin (IL)-1, IL-4 and IL-10 genes (Table 1) in 220 Taiwanese Chinese with GC and in 230 healthy controls. The risk of genotype was adjusted with confounding environmental risks. Our results revealed that the frequency of *Helicobacter pylori* infection [odds ratio (OR) 1.7, 95% confidence interval (CI) 1.19~2.56], cigarette smoking (OR:2.02, 95%CI 1.38~2.95) and high IL-10 producer genotype (OR:2.67, 95%CI 1.29-5.50) was significantly increased in the entire GC patients. Among different subtypes of GC, a higher risk of developing diffuse type (OR:1.64, 95%CI 1.01~2.67) or cardia cancer (OR:2.44, 95%CI 1.13~2.67) was observed for the CT/CC genotype of IL-4 at the position -590, whereas the high IL-10 producer genotype was significantly linked with the risk of cardia cancer (OR:3.21, 95%CI 1.06-9.73) or advanced stage (OR:2.29, 95%CI 1.12-4.64). No association was noted between GC and controls in the distribution of IL-1 and TNF- $\alpha$  genotypes. Logistic regression analyses revealed that *H. pylori* infection (OR:1.7, 95%CI 1.14~2.52), cigarette smoking (OR:1.87, 95%CI 1.27-2.96) and IL-10 genotype (OR:2.54, 95%CI 1.24~5.61) are independent risks for GC. Independent effects of IL-10 genotype, *H. pylori* infection and cigarette smoking indicate that carcinogenesis of GC is influenced by a variety of host and environmental factors.

In the second year grant period, analyses of gene expression in AGS induced by different strains of *H. pylori* has been performed by cDNA microarray. We have found some differences among bacterial strains from various disease (ulcer vs. cancer vs. maltoma). In addition, case and control enrollment keeps continuing and is more than 250 in numbers. Genetic polymorphisms in TNF- $\alpha$  (-238, -308, -857, -863, -1031), TNFR1 (-383), TNFR2 (codon 196), CD14 (-159), TLR4 (Asp299 Gly and Thr 399 Ile) and IL-8 (-251) have been investigated in GC, maltoma and controls. Our results demonstrated TNF- $\alpha$  -857 C $\rightarrow$ T was significantly underrepresented in maltoma compared to controls (6.4% vs. 14.3%,  $p=0.018$ ), conferring a 3-fold decreased in risk (OR=0.33, 95%CI 0.15-0.75). Comparison of allele frequencies between GC and controls failed to show any statistical significance. The differences in genetic background as well as divergent clinicopathologic features between GC and maltoma supports the notion that fundamental mechanistic differences exist in these 2 well-defined *H. pylori*-associated malignancies.

In the third year grant period, we have collected at least 300 cases of

GC with different but well-characterized phenotypes for further genotypic analyses of IL-6 (-634 C/G) and IL-8 (-137 G/C & -607 C/A). In addition, we have found several novel genes with biologic significance in H. pylori-associated diseases disclosed by cDNA microarray. Their roles in host susceptibility of gastric carcinogenesis will be tested in future investigations. With our efforts and achievements in this field, we were invited to write a review article describing the implication of host-environment interaction in gastric carcinogenesis, which has been published in Cancer Epidemiol Biomarker & Prev.

Keywords: Gastric cancer, Helicobacter pylori, Genotype, Host Susceptibility

Table I. Primer sequences and methods used for detection of cytokine gene polymorphisms

Gene	Primers	PCR condition	Detection method
TNF- $\alpha$ (-238,-308)	5'-CAAACACAGGCCTCAGGACTC-3' 5'-AGGGAGCGTCTGCTGGGCTG-3'	95°C 40s 65°C 1min 72°C 30s 35 cycles	direct sequencing with primer 5'-TCTGGAAGTTAGAAGGAAAC-3'
TNF- $\alpha$ (-857,-863, -1031)	5'-GCTTGTGTGTGTGTGTCTGG-3' 5'-GGACACACAAGCATCAAGG-3'	94°C 1min 55°C 2min 72°C 3min	Direct sequencing with primer 5'-TGTGGCCATATCTTCTTAAA-3'
TNFR1 (-383)	5'-TTATTGCCCTTGGTGTGGTTG-3' 5'-GGAGGGGAAGAGTGAGGCACTGTT-3'	95°C 30sec 65°C 30sec 72°C 30sec 35 cycles	BglII digestion, 2.5% agarose gel electrophoresis
TNFR2 (codon 196)	5'-ACTCTCCTATCCTGCCTGCT-3' 5'-TTCTGGAGTTGGCTGCGTGT-3'	95°C 30sec 72°C 30sec 35 cycles	NIaIII digestion 4% agarose gel electrophoresis
IL-1 $\beta$ (-31,-511)	5'-CTCAGAGGCTCCTGCAATTG-3' 5'-AGATAAGCAGTATCCATTCCC-3'	95°C 45s 55°C 1min 72°C 1min 35 cycles	direct sequencing with primer 5'-TCGTTCTGCAGTTGATGTCCA-3'
IL-1RN (intron 2 VNTR)	5'-CTCAGCAACACTCCTAT-3' 5'-TCCTGGTCTGCAGGTAA-3'	95°C 1min 58°C 1min 72°C 1min 35 cycles	size fractionation in 2.5% agarose gel electrophoresis
IL-4(-590)	5'-ACTAGGCCTCACCTGATACG-3' 5'-GTTGTAATGCAGTCCTCCTG3'	95°C 30s 55°C 30s 72°C 30s 35 cycles	Bsm FI digestion, 6% polyacrylamide gel electrophoresis
IL-4R (Ile50 Val)	5'-GGCAGGTGTGAGGAGCATCC-3' 5'-GCCTCCGTTGTTCTCAGGTA-3'	95°C 30s 55°C 30s 72°C 1min 35 cycles	Rsa I digestion, 3% agarose & 1% NuSieve agarose gel electrophoresis
IL-4R(Q576R)	5'-GCCCCACCAGTGGCTACC-3' 5'-GCCTTGTAACCAGCCTCTCCT-3'	94°C 30s 55°C 30s 72°C 1min 40 cycles	Msp I digestion, 6% polyacrylamide gel electrophoresis
IL-10(-592,-819,-1082)	5'-ATCCAAGACAACACTACTAA-3' 5'-TAAATATCCTCAAAGTTCC-3'	94°C 40s 54°C 45s 72°C 30s 35 cycles	direct sequencing with primer 5'-TAAATATCCTCAAAGTTCC-3' and 5'-TTGGCCTTAGAGTTTCTTTT-3'



## 參考文獻

1. Fuchs CS, Mayer RJ. Gastric carcinoma. *N Engl J Med* 1995; 333:32-41.
2. Wu MS, Lin JT, Lee WJ, Yu SC, Wang TH. Gastric cancer in Taiwan. *J Formos Med Assoc* 1994; 93:S77-S89.
3. Correa P. Human gastric carcinogenesis: a multistep and multifactorial process. *Cancer Res* 1992; 52:6735-6740.
4. Huang JQ, Sridhar S, Chen Y, Hunt R. Metanalysis of the relationship between *Helicobacter pylori* seropositivity and gastric cancer. *Gastroenterology* 1998;114:1169-1179.
5. Parkin DM, Bray FI, Devesa SS. Cancer burden in the year 2000. The global picture. *Eur J Cancer* 2001;37:S4-S66.
6. Palli D, Masala G, Russo A, Saieva C, Bazzoli F. When enthusiasm is infectious. *Gastroenterology* 2000;119:274-275.
7. Blaser MJ, Perez-Perez GI, Kleanthous H, Cover TL, Peek RM, Chyou PH, Stemmermann GN, Nomura A. Infection with *Helicobacter pylori* strains possessing *cag A* associated with an increased risk of developing adenocarcinoma of the stomach. *Cancer Res* 1995;55:2111-2115.
8. Parsonnet J, Friedman GD, Orentreich N, Vogelman H. Risk for gastric cancer in people with CagA positive or CagA negative *Helicobacter pylori* infection. *Gut* 1997;40:297-301.
9. Yang JC, Wang TH, Wang HJ, Kuo CH, Wang JH, Wang WC. Genetic analysis of the cytokine-associated gene and the vacuolating toxin gene in *Helicobacter pylori* strains isolated from Taiwanese patients. *Am J Gastroenterol* 1997;92:1316-1321.
10. Sakagami T, Dixon M, O'Rourke J, Howlett R, Alderuccio F, Vella J, Shimoyama T, Lee A. Atrophic gastritis changes in both *Helicobacter felis* and *Helicobacter pylori* infected mice are host dependent and separate from antral gastritis. *Gut* 1996;39:639-648.
11. Moss SF, Sordillo EM, Abdalla AM, Makarov V, Hanzley Z, Perez-Perez GI, Blaser MJ, Holt PR. Increased gastric epithelial cell apoptosis associated with colonization with *cagA+* *Helicobacter pylori* strain. *Cancer Res* 2001;61:1406-1411.

12. Malaty H, Engstrand L, Pedersen N, Graham DY. Helicobacter pylori infection: genetic and environmental influences. *Ann Intern Med* 1994;120:982-986.
13. Magnusson PKE, Enroth H, Eriksson I, Held M, Nyren D, Engstrand L, Hansson LE, Gyllensten UB. Gastric cancer and human leukocyte antigen: distinct DQ and DR alleles are associated with development of gastric cancer and infection by Helicobacter pylori. *Cancer Res* 2001;61:2684-2689.
14. Nedrud JG, Czinn SJ. Host, heredity and Helicobacter. *Gut* 1999;45:323-324.
15. Ernst P. The role of inflammation in the pathogenesis of gastric cancer. *Aliment Pharmacol Ther* 1999;13:13-18.
16. Crabtree JE. Gastric mucosal inflammatory responses to H. pylori. *Aliment Pharmacol Ther* 1996;10 (suppl 1):29-37.
17. El-Omar EM. The importance of interleukin 1 $\beta$  in Helicobacter pylori associated disease. *Gut* 2001;48:743-747.
18. Imanishi J. Expression of cytokines in bacterial and viral infections and their biochemical aspects. *J Biochem* 2000;127:525-530.
19. Berg DJ, Lynch NA, Lynch RG, Lauricella DM. Rapid development of severe hyperplastic gastritis with gastric epithelial dedifferentiation in Helicobacter felis-infected IL-10 (-/-) mice. *Am J Pathol* 1998;152:1377-1386.
20. Mohammadi M, Nedrud J, Redline R, Lycke N, Czinn SJ. Murine CD4 T-cell response to Helicobacter infection: TH1 cells enhance gastritis and TH2 cells reduce bacterial load. *Gastroenterology* 1997;113:1848-1857.
21. Shimoyama T, Everett SM, Fukuda S, Axon AT, Dixon MF, Crabtree JE. Influence of smoking and alcohol on gastric chemokine mRNA expression in patients with Helicobacter pylori infection. *J Clin Pathol* 2001;54:332-334.
22. Westendorp RG, Langermans JA, Huizinga TW, Elouali AH, Verweij CL, Boomsma DI, Vandenbrouke JP. Genetic influence on cytokine production and fatal meningococcal disease. *Lancet* 1997;349:170-173.
23. Hogan CM, Hammer SM. Host determinants in HIV infection and disease. *Ann Intern Med* 2001;134:978-996.

24. Thio CL, Thomas DL, Carrington M. Chronic viral hepatitis and the human genome. *Hepatology* 2000;31:819-827.
25. El-Omar EM, Chow WH, Rabkin CS. Gastric cancer and H. pylori: Host genetics open the way. *Gastroenterology* 2001;21:1002-1004.
26. El-Omar EM, Carrington M, Chow WH, McColl KEL, Bream JH, Young HA, Herrera J, Lissowska J, Yuan CC, Rothman N, Lanyon G, Martin M, Fraumeni JF, Jr., Rabkin CS. Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature* 2000;404:398-402.
27. Machado JC, Pharoah P, Sousa S, Carvalho R, Oliveira C, Figueiredo C, Amorim A, Seruca R, Caldas C, Carneiro F, Sobrinho-Simoes M. Interleukin-1  $\beta$  and interleukine 1RN polymorphisms are associated with increased risk of gastric carcinoma. *Gastroenterology* 2001;121:823-829.
28. Kunstmann E, Epplen C, Elitok E, Harder M, Suerbaum S, Peitz U, Schmiegel W, Epplen JT. Helicobacter pylori infection and polymorphisms in the tumor necrosis factor region. *Electrophoresis* 1999;20:1756-1761.
29. Stadtlander CT, Waterbor JW. Molecular epidemiology, pathogenesis and prevention of gastric cancer. *Carcinogenesis* 1999;20:2195-2208.
30. Kato S, Onda M, Yamada S, Matsuda N, Tokunaga A, Matsukwa N. Association of the interleukin-1 beta genetic polymorphism and gastric cancer risk in Japanese. *J Gastroenterol* 2001;36:696-699.
31. Tseng LH, Chen PJ, Lin MT, Shau WY, Chuang SM, Martin PJ, Hansen JA. Single nucleotide polymorphisms in intron 2 of the human interleukin-1 receptor antagonist (IL-1 Ra) gene: further definition of the IL-1 $\beta$  and IL-1 Ra polymorphisms in North American Caucasians and Taiwanese Chinese. *Tissue Antigens* 2001;57:318-324.
32. Manger ID, Relman DA. How the host sees pathogens: global gene expression responses to infection. *Curr Opin Immunol* 2000;12:215-218.
33. Cumming CA, Relman DA. Using DNA microarrays to study host-microbe interactions. *Emerg Infect Dis* 2000;6:513-525.
34. Chen JJ, Peck K, Hong TM, Yang SC, Sher YP, Shih JY, Wu R,

Chen JL, Roffler SR, Wu CW, Yang PC. Global analysis of gene expression in invasion by a lung cancer model. *Cancer Res* 2001;61:5223-5230.

35. Wu MS, Shun CT, Wang HP, Sheu JC, Lee WJ, Wang TH, Lin JT. Genetic alterations of gastric cancer relation to histological subtypes, tumor stage, and *Helicobacter pylori* infection. *Gastroenterology* 1997;112:1457-1465.

36. Wu MS, Lee WC, Shun CT, Wang HP, Lee WJ, Sheu JC, Lin JT. Clinicopathologic significance of altered loci of replication error and microsatellite instability associated mutations in gastric cancer. *Cancer Res* 1998;58:1494-1497.

#### 已發表或投稿之論文

1. Wu MS, Wu CY, Chen CJ, Lin MT, Shun CT, Lin JT. Interleukin-10 genotypes associated with the risk of gastric carcinoma in Taiwanese Chinese. *Int J Cancer* 2003;104:617-623.
2. Wu MS, Chen CJ, Lin JT. Genetic alterations and polymorphisms in gastric cancer. *J Formos Med Assoc* 2003;102:447-458.
3. Wu MS, Chen LC, Shun CT, Huang SP, Chiu HM, Wang HP, Lin MT, Cheng AL, Lin JT. Promoter polymorphisms of tumor necrosis factor- $\alpha$  are associated with risk of gastric mucosa-associated lymphoid tissue lymphoma. *Int J Cancer* 2004;110:695-700.
4. Wu MS, Cheng TY, Shun CT, Lin MT, Chen LC, Lin JT. Functional polymorphisms of CD14 and toll-like receptor 4 in Taiwanese Chinese with *Helicobacter pylori*-related gastric malignancies. *Hepatogastroenterology* (in press)
5. Chang YT, Wu MS, Chang YC, Yang PC, Chen CC, Lin JT. Distinct gene expression profiles in gastric epithelial cells induced by different clinical isolates of *Helicobacter pylori*-implication of bacteria and host interaction in gastric carcinogenesis. *Hepatogastroenterology* (in press)
6. Wu MS, Chen CJ, Lin JT. Host-environmental interactions: Their impact on progression from gastric inflammation to carcinogenesis and on development of new approaches to prevent and treat gastric cancer. *Cancer Epidemiol Biomark & Prev* 2005; 14:1878-1882. [SCI]