Coupling of Carbomethoxy and Vinylidene Ligand in Molybdenum Complexes

Jung-Yen Yang, Shou-Ling Huang, Ying-Chih Lin,* Yi-Hong Liu, and Yu Wang

Department of Chemistry, National Taiwan University, Taipei, Taiwan, 106 Republic of China

Received September 13, 1999

Treatment of [Cp(dppe)(CO)Mo=C=C(Ph)CH₂CH=CH₂]I (2) with MeONa caused nucleophilic attack to occur at the terminal CO ligand. This is followed by a coupling reaction of the resulting carbomethoxy group with $C\alpha$ of the vinylidene ligand accompanied with coordination of the terminal olefin to afford Cp(dppe)MoC(COOMe)=C(Ph)CH₂CH=CH₂ (5). Similar coupling was observed when $[Cp(dppe)(CO)Mo=C=C(Ph)CH_2Ph]I$ (3) was treated with MeONa, but the reaction afforded the neutral allylic complex Cp(dppe)Mo[CH-(COOMe)C(Ph)CHPh] (6). The structures of complexes 5 and 6 have been determined by X-ray diffraction analysis.

Introduction

We previously described a new type of deprotonation reaction of several cationic vinylidene complexes leading to a rare class of metal complexes containing various cyclic ligands.¹ Using this synthetic strategy, we successfully prepared several ruthenium cyclopropenyl complexes containing various substituents as well as ruthenium furanyl complexes. For the cyclopropenyl complex, the cyclization reaction also results in formation of a chiral carbon center in the three-membered ring. These features render this cyclization process potentially useful for organic synthesis.² Therefore, we set out to study the chemical reactivity of a molybdenum vinylidene system of Cp(Ph₂PCH₂CH₂PPh₂)(CO)Mo.³ Such a system was chosen because the presence of donor phosphine ligands is known to assist the preparation of cationic metal vinylidene complexes,⁴ and the presence of a CO ligand could be utilized to study carboncarbon bond formation. Surprisingly, treatment of these vinylidene complexes containing a terminal CO ligand with sodium methoxide affords unexpected products via an unprecedented coupling of a carbomethoxy group with Ca of the vinylidene ligand.⁵ Coupling reactions

(3) (a) Allen, S. R.; Beevor, R. G.; Green, M.; Norman, N. C.; Orpen, A. G.; Williams, I. D. J. Chem. Soc., Dalton Trans. **1985**, 435. (b) Beevor, R. G.; Green, M.; Orpen, A. G.; Williams, I. D. J. Chem. Soc., Dalton Trans. **1987**, 1319. (c) Nickias, P. N.; Seleague, J. P.; Young, B. A. Organometallics **1988**, 7, 2248. (d) Chang, K. H.; Lin, Y. C. Chem. Commun. 1998, 1441.

(4) (a) Bruce, M. I. Chem. Rev. 1991, 91, 197. (b) Bruce, M. I.;

involving $C\alpha$ of a vinylidene ligand on the metal complex are limited in the literature. Herein we report preparation of a number of cationic molybdenum vinylidene complexes Cp(Ph₂PCH₂CH₂PPh₂)(CO)Mo=C=C(Ph)-CH₂R and nucleophilic addition of methoxide to the CO ligand followed by a coupling reaction leading to carboncarbon formation at $C\alpha$ of the vinylidene ligand.

Results and Discussion

Preparation of the Acetylide Complex 1. The acetylide complex [Mo](CO)C=CPh (1, [Mo] = $(\eta^5$ - C_5H_5)(dppe)Mo, dppe = Ph₂PCH₂CH₂PPh₂) was prepared by two different methods. The reaction of [Mo]-(CO)Cl with LiCCPh in THF, which gave 1 in reasonable yield, required handling of an air-sensitive lithium reagent. We therefore pursued an alternative synthesis, namely, the reaction of [Mo](CO)Cl with HC=CPh. This reaction gave {[Mo](CO)=C=CHPh}Cl, which underwent thermal decarbonylation to afford $\{[Mo](\eta^2-HC)=$ CPh)}Cl. Then the η^2 -phenylacetylene complex was treated with CO in the presence of MeONa to give 1. The tautomerization of alkyne to vinylidene on the bisdimethylphenylphosphine molybdenum analogue has been reported before.^{3c} Our scheme involves more steps but gives higher overall yield.

The ³¹P NMR spectrum of **1** exhibits two sets of doublet resonances at δ 90.4 and 78.7 with $J_{P-P} = 37.2$ Hz. In the ¹H NMR spectrum, the resonance for the Cp ligand is a doublet at δ 4.49 with $J_{\rm H-P}$ = 1.6 Hz, indicating coupling with only one of the phosphorus atoms of the dppe. In the ¹³C NMR spectrum, the signal for the terminal CO ligand is also a doublet at δ 245.0 with $J_{C-P} = 22.2$ Hz, indicating the same coupling pattern. By the use of selective heteronuclear decoupling techniques, i.e., ¹H{³¹P} and ¹³C{³¹P} NMR spectra, we are able to determine that both ¹H-³¹P coupling of the Cp ligand and ¹³C-³¹P coupling of the CO ligand originate from the same phosphorus atom displaying the downfield resonance (at δ 90.4) in the ³¹P NMR spectrum. Coupling between carbon and phosphorus nuclei of the complex with a piano-stool coordination geometry

^{(1) (}a) Ting, P. C.; Lin, Y. C.; Cheng, M. C.; Wang, Y. Organome-tallics **1994**, *13*, 2150. (b) Ting, P. C.; Lin, Y. C.; Lee, G. H.; Cheng, M. C.; Wang, Y. J. Am. Chem. Soc. **1996**, *118*, 6443. (c) Chang, C. W.; Ting, P. C.; Lin, Y. C.; Lee, G. H.; Wang, Y. J. Organomet. Chem. **1998**, *553*, 417. (d) Lo, Y. H.; Lin, Y. C.; Lee, G. H.; Wang, Y. Organometallics 1999, 18, 982.

^{(2) (}a) Kubota, K.; Mori, S.; Nakamura, M.; Nakamura, E. J. Am. *Chem. Soc.* **1998**, *120*, 1334. (b) Muller, P.; Imogai, H. *Tetrahedron: Asymmetry* **1998**, *9*, 4419. (c) Plemenkov, V. V.; Rul, S. V.; Gubaidullin, A. T.; Litvinov, I. A.; Karaseva, I. P.; Nuretdinov, I. A. *Russ. J. Org.* Chem. 1998, 34, 971.

⁽a) Diuce, M. I. Chenh. Rev. 1991, 91, 197. (b) Bruce, M. I.;
Swincer, A. G. Adv. Organomet. Chem. 1982, 20, 159.
(5) (a) Buil, M. L.; Esteruelas, M. A.; Lòpez, A. M.; Oñate, E. Organometallics 1997, 16, 3169. (b) Bianchini, C.; Casares, J. A.;
Peruzzini, M.; Romerosa, A.; Zanobini, F. J. Am. Chem. Soc. 1996, 118, 4585. (c) Fauro M.; Montetta, L. Darredina, D. Linger, 2016, 2016, 2017. 4585. (c) Faure, M.; Maurette, L., Donnadieu, B.; Lavigne, G. Angew. Chem., Int. Ed. Engl. 1999, 34, 518.



depends on the orientation.⁶ In the ¹H NMR spectrum of {[Mo](η^2 -HC=CPh)}Cl, the characteristic triplet resonance at δ 10.02 with $J_{H-P} = 13.8$ Hz is assigned to the terminal CH. The corresponding ¹H signal of the vinylidene proton of {[Mo](CO)=C=CHPh}Cl appears as a singlet at δ 4.78.

Preparation of Vinylidene Complexes. Treatment of 1 with $BrCH_2R$ (R = CH=CH₂, C₆H₅, and CN) afforded cationic brown-red vinylidene complexes {[Mo]- $(CO)=C=C(Ph)CH_2R$ }Br (2, R = CH=CH₂; 3, R = C₆H₅; **4**, $\mathbf{R} = \mathbf{CN}$). All reactions gave the desired products in 70-80% yields. Use of organic iodides gave lower yields. All these vinylidene complexes are soluble in CHCl₃, CH₂Cl₂, and CH₃OH but are insoluble in hexane. Complex 2 is air- and moisture-sensitive but is stable under nitrogen. The ¹H NMR spectrum of **2** displays multiplet resonances centered at δ 5.00, 4.96 and 2.87, 2.78 assignable to the olefinic and saturated diastereotopic CH₂ units, respectively, and the multiplet resonance at δ 5.62 to the allylic CH unit. The geminal coupling constant of the diastereotopic methylene protons is 15.6 Hz, and interestingly, the long-range P–H coupling with one of the phosphorus atoms (${}^{5}J_{P-H} = 2.8$ and 2.0 Hz) is also observed for these protons. The ¹H NMR spectra of complexes 3 and 4 display the same features. In the ¹³C NMR spectrum, the characteristic doublet of doublet resonance centered at δ 358.4 with ${}^{2}J_{P-C} = 36.0$ and 6.5 Hz is assigned to the vinylidene Ca. In the ³¹P NMR spectrum, two doublet resonances at δ 71.5 and 67.0 with $J_{P-P} = 37.5$ Hz are assigned to the dppe ligand. In the ³¹P NMR spectra of **3** and **4**, the ³¹P resonances fall in a similar region with comparable coupling constants (δ 69.0 and 66.2 with $J_{P-P} = 36.3$ Hz for **3** and δ 66.8 and 63.7 with $J_{P-P} = 30.7$ Hz for **4**). Other spectroscopic features are consistent with their formulations.

Reactions of 2 and 3 with MeONa. Having established the cyclopropenation reaction of several cationic vinylidene complexes of ruthenium, we examined the deprotonation reaction of the molybdenum complex 2 using various bases with an expectation to see similar results. However, much to our surprise, treatment of 2 with n-Bu₄NOH, DBU, and n-Bu₄NF gave no reaction. In the reaction of 2 with MeONa the coupling product $[Mo](\eta^3-C(COOMe)=C(Ph)CH_2CH=CH_2)$ (5) (Scheme 1) was obtained in high yield. Formation of 5 is rationalized by nucleophilic addition of MeO- to the terminal CO ligand⁷ giving the carbomethoxy group followed by a coupling reaction⁸ of the carboxymethoxy group with $C\alpha$ of the vinylidene ligand to afford **5** (Scheme 1) in high yield. The terminal olefin group, acting as a twoelectron donor, fills the vacant site left from the coupling reaction. It is interesting to note the difference in spectral characteristics of 2 and 5. Signals of the terminal vinyl group in the ¹H NMR spectrum of 2 appear at δ 5.62 (=CH) and 5.00, 4.96 (=CH₂), while those of **5** occur at δ 1.78 (=CH) and 2.58, 3.66 (=CH₂). These assignments are determined by the use of 2D COSY and HSQC experiments. While most of H-H

^{(6) (}a) Bainbridge, A.; Craig, P. J.; Green, M. J. Chem. Soc. (A) **1968**, 2713. (b) Faller, J. W.; Anderson, A. S. J. Am. Chem. Soc. **1970**, 92, 5852. (c) Beach, D. L.; Barnett, K. W. J. Organomet. Chem. **1975**, 97, C27. (d) Sakaba, H.; Ishida, K.; Horino, H. Chem. Lett. **1995**, 1145.

^{(7) (}a) Nakazawa, H.; Kadoi, Y.; Mizuta, T.; Miyoshi, K.; Yoneda, H. J Organomet. Chem. **1989**, *366*, 333. (b) Gibson, D. H.; Ong, T. S.; Ye, M.; Franco, J. O.; Owens, K. Organometallics **1988**, *7*, *2569*. (c) Busetto, L. L.; Zanotti, V.; Norfo, L.; Palazzi, A.; Albano, V. G.; Braga, D. Organometallics **1993**, *12*, 190. (d) Nakazawa, H.; Yamaguchi, M.; Kubo, K.; Miyoshi, K. J Organomet. Chem. **1992**, *428*, 145. (e) Werner, H.; Hofmann, L.; Zolk, R. Chem. Ber. **1987**, *120*, 379. (f) Kiel, W. A.; Buhro, W. E.; Gladysz, J. A. Organometallics **1984**, *3*, 879. (g) Carlos, F.; Barrientos-Penna, A.; Hugo, K. O.; Derek, S. Organometallics **1985**, 4, 367. (h) Bao, Q. B.; Rheingold, A. L.; Brill, T. B. Organometallics **1986**, *5*, *2259*.

^{(8) (}a) Wiedemann, R.; Steinert, P.; Gevert, O.; Werner, H. J. Am. Chem. Soc. **1996**, *118*, 2495. (b) Braun, T.; Meuer, P.; Werner, H. Organometallics **1996**, *15*, 4075.



Figure 1. ORTEP drawing of **5** with thermal ellipsoids shown at the 30% probability level. For the dppe phenyl groups, only the ipso carbons are shown.

Table 1. Selected Bond Distances (Å) and Angles(deg) of

Cp(dppe)MoC(COOMe)=C(Ph)CH ₂ CH=CH ₂ (5)					
Mo-P1	2.4844(11)	C3-C4	1.494(5)		
Mo-P2	2.4794(11)	C4-C5	1.338(5)		
Mo-C1	2.216(3)	C4-C8	1.493(5)		
Mo-C2	2.295(3)	C5-C6	1.473(5)		
Mo-C5	2.248(3)	C8-C9	1.383(5)		
P1-C14	1.855(4)	C8-C13	1.380(5)		
P1-C16	1.842(4)	C9-C10	1.382(6)		
P1-C22	1.841(4)	C10-C11	1.367(7)		
P2-C15	1.829(4)	C11-C12	1.345(7)		
P2-C28	1.824(4)	C12-C13	1.383(6)		
P2-C34	1.844(4)	C14-C15	1.521(5)		
O1-C6	1.199(5)	C40-C41	1.400(6)		
O2-C6	1.355(4)	C40-C44	1.413(6)		
O2-C7	1.433(5)	C41-C42	1.380(6)		
C1-C2	1.406(5)	C42-C43	1.401(6)		
C2-C3	1.503(5)	C43-C44	1.395(6)		
P1-Mo-P2	77.36(4)	Mo-C2-C3	114.54(22)		
P1-Mo-C1	126.81(11)	C1-C2-C3	119.9(3)		
P1-Mo-C2	90.56(10)	C2-C3-C4	110.2(3)		
P1-Mo-C5	83.71(9)	C3-C4-C5	116.7(3)		
P2-Mo-C1	86.08(10)	C3-C4-C8	116.1(3)		
P2-Mo-C2	74.37(9)	C5-C4-C8	127.1(3)		
P2-Mo-C5	140.73(10)	Mo-C5-C4	121.01(25)		
C1-Mo-C2	36.28(14)	Mo-C5-C6	121.51(24)		
C1-Mo-C5	78.22(12)	C4-C5-C6	117.4(3)		
C2-Mo-C5	71.72(12)	O1-C6-O2	121.1(3)		
C6-O2-C7	115.6(3)	O1-C6-C5	126.7(3)		
Mo-C1-C2	74.89(20)	O2-C6-C5	112.2(3)		
Mo-C2-C1	68.84(20)				

coupling constants in the allylic group are measurable for **2**, only the coupling constant of 17.2 Hz between mutually trans-protons in the coordinated olefin is attainable for 5. However, significant upfield shifts indicate π -coordination of the terminal double bond in 5. Surprisingly, in the ³¹P NMR spectrum of 5, two singlet resonances at δ 90.5 and 71.2 with equal intensity are observed possibly because of a particular orientation of the dppe ligand.⁹ In the ¹³C NMR spectrum of **5**, resonances of π -coordinated olefinic carbon atoms appeared at δ 60.3 (dd) and 47.6 (d) with $J_{C-P} =$ 5.4, 4.0, and 5.9 Hz, respectively. To fully characterize this product, single crystals of 5 were grown from a hexane/CH₂Cl₂ solution, and the molecular structure was determined by X-ray diffraction analysis. An ORTEP drawing is shown in Figure 1, and selected bond distances and angles are listed in Table 1. The environment about the metal center corresponds to a structure of distorted four-legged piano-stool geometry. The carbomethoxy group resulting from nucleophilic attack of



Figure 2. ORTEP drawing of **6** with thermal ellipsoids shown at the 30% probability level. For the dppe phenyl groups, only the ipso carbons are shown.

MeO⁻ to the terminal CO ligand is bound to C α . And it is clear that the internal olefin C4-C5 is in an E configuration with the phenyl group and the Mo fragment in a trans disposition. The terminal olefin is oriented such that the C=C bond is contained in the plane defined by the metal, the center of the Cp ring, and the trans phosphorus atom. The olefinic ligand coordinates to the metal center with Mo-C1 = 2.216-(3) Å and Mo-C2 = 2.295(3) Å. The Mo-C5 bond length of 2.248(3) Å is typical of a Mo-C single bond, and the C4-C5 bond length of 1.338(5) Å typical of a C=C double bond. The C1–C2 bond length of 1.406(5) Å, resulting from coordination to the metal center, is slightly longer that that of the C4–C5 double bond. The C=C double bond of 1.437(9) Å in the cationic olefin complex $Cp[P_2(Me)_4C_2H_4](CO)Mo(C_2H_3Ph)^{10}$ is slightly longer possibly due to stronger back-bonding to the antibonding orbital of the olefin.

Treatment of **3** with MeONa similarly causes nucleophilic addition of MeO⁻ to the terminal CO ligand followed by an analogous coupling reaction and affords the neutral allylic product [Mo][η^3 -CH(COOMe)C(Ph)-CHPh] (**6**) in high yield. Due to the lack of a terminal olefin donor group, the vinylidene ligand in **3** transforms to the η^3 -allylic ligand of **6** possibly by coupling followed by a 1,3-hydrogen shift. The ¹³C resonance attributed to the ester carbon appears at δ 183.2, and C–P coupling constants of three multiplet resonances, appearing at δ 48.7 (d), 90.3 (t), and 49.2 (d) with $J_{C-P} = 4.5$, 3.1, and 2.7 Hz, respectively, clearly indicate the η^3 -allylic bonding mode. Two ³¹P resonances at δ 71.3 and 72.2 with $J_{P-P} = 36.8$ Hz are assigned to the dppe ligand.

The structure of **6** has also been confirmed by an X-ray diffraction analysis. An ORTEP drawing is shown in Figure 2, and selected bond distances and angles are listed in Table 2. The carbomethoxy group is again bound to the allylic terminal carbon. The trisubstituted allylic ligand is in an endo conformation with a synphenyl group and an anti-carbomethoxy group at two terminal carbon atoms. The two allylic C–C bond lengths are about equal (C1–C4 = 1.463(4) Å, C4–C5 = 1.446(4) Å), with the Mo–C1 bond slightly shorter

⁽⁹⁾ Morton, M. S.; Lachicotte, R. J.; Vicic, D. A.; Jones, W. D. Organometallics 1999, 18, 227.

^{(10) (}a) Kegley, S. E.; Bergstrom, D. T.; Crocker, L. S.; Weiss, E. P.; Berndt, W. G.; Rheingold, A. L. *Organometallics* **1991**, *10*, 573. (b) Abugideiri, F.; Kelland, M. A.; Poli, R. *Organometallics* **1992**, *11*, 1311. (c) Kegley, S. E.; Walter, K. A.; Bergstrom, D. T.; MacFarland, D. K.; Young, B. G.; Rheingold, A. L. *Organometallics* **1993**, *12*, 2339.

 Table 2. Selected Bond Distances (Å) and Angles
 (deg) of Cp(dppe)MoCH(COOMe)C(Ph)CHPh (6)

•			
Mo-P1	2.4992(8)	P2-C38	1.839(3)
Mo-P2	2.4951(8)	O1-C2	1.210(4)
Mo-C1	2.259(3)	O2-C2	1.376(4)
Mo-C4	2.328(3)	O2-C3	1.435(4)
Mo-C5	2.324(3)	C1-C2	1.462(4)
P1-C19	1.851(4)	C1-C4	1.463(4)
P1-C20	1.858(3)	C4-C5	1.446(4)
P1-C26	1.858(3)	C4-C6	1.510(4)
P2-C18	1.857(3)	C5-C12	1.483(4)
P2-C32	1.846(3)		
P1-Mo-P2	77.63(3)	C1 - C2 - O1	130.3(3)
P1-Mo-C1	87.27(8)	C1 - C2 - O2	109.0(3)
P1-Mo-C4	100.36(8)	01 - C2 - O2	120.7(3)
P1-Mo-C5	134.73(8)	C2 - O2 - C3	115.3(3)
P2-Mo-C1	132.36(8)	C1-C4-C5	113.1(3)
P2-Mo-C4	101.46(7)	C1-C4-C6	117.7(3)
P2-Mo-C5	96.09(8)	C5-C4-C6	126.5(3)
C2-C1-C4	124.2(3)	C4-C5-C12	129.5(3)
	(-)		

than the other two Mo–C(allylic) bonds (Mo–C1 = 2.259(3) Å, Mo–C4 = 2.328(3) Å, and Mo–C5 = 2.324-(3) Å). The different configuration of the carbomethoxy group relative to the neighboring phenyl group (cis in **5** and trans in **6**) could possibly be attributed to the additional hydrogen shift process in the formation of **6**.

Treatment of **4** with MeONa in MeOH gave, in moderate yield, an unstable green complex, which decomposed at room temperature in 2 h in CDCl₃. This solid product was isolated by precipitation from the CH₂-Cl₂ solution of the product via addition of ether. The FAB mass spectrum of this complex displays peaks that could be attributed to the cyclopropenyl complex [Mo]-

 $(CO)\dot{C}=C(Ph)\dot{C}HCN$ (7). In the IR spectrum of 7, the absorption at 1846 cm⁻¹ is assigned to the vibrational stretching of the terminal CO ligand, indicating that there is no nucleophilic addition. And in the ³¹P NMR spectrum, chemical shifts of two singlet resonances at δ 105.4 and 103.9 differ significantly from the range (δ 90.5 and 71.2) observed for **5** and **6**. Unfortunately, we can obtain spectroscopic data only for this green compound, and there is no established data for a molybdenum cyclopropenyl complex for comparison. This is somewhat surprising since in the ruthenium system¹ the presence of an electron-withdrawing CN substituent seemed to stabilize a number of cyclopropenyl complexes. In this Mo vinylidene system with a terminal CO ligand, the CN group provides no similar effect. In our attempts to carry out addition reactions of 4 using nucleophilic reagents other than MeO⁻, we do not observe any coupling products as in the reactions of **2** and 3.

In summary, we report high-yield preparation of three cationic molybednum vinylidene complexes {[Mo](CO)= C=C(Ph)CH₂R}Br (**2**, R = CH=CH₂; **3**, R = C₆H₅; **4**, R = CN). In the presence of MeONa, the molybdenum vinylidene complexes **2** and **3**, each containing a terminal carbonyl ligand, undergo nucleophilic addition at the CO ligand followed by a C-C coupling reaction at C α of the vinylidene ligand to give addition products. The site preference of this coupling reaction is possibly due to the relatively strong bond of Ru=C α and proximity of the two groups on the ruthenium metal center. For the analogous CN-substituted vinylidene complex **4**, the same reaction yielded an unstable cyclopropenyl complex through deprotonation.

Experimental Section

General Procedures. All manipulations were performed under nitrogen using vacuum-line, drybox, and standard Schlenk techniques. CH₂Cl₂ was distilled from CaH₂ and diethyl ether and THF from Na/diphenylketyl. All other solvents and reagents were of reagent grade and were used as received. NMR spectra were recorded on Bruker AM-300WB and DMX-500 FT-NMR spectrometers at room temperature (unless states otherwise) and are reported in units of δ with residual protons in the solvents as a standard (CDCl₃, δ 7.24; C_2D_6O , δ 2.04). FAB mass spectra were recorded on a JEOL SX-102A spectrometer. Complex [Mo](CO)Cl ([Mo] = $(\eta^{5}-\eta^{5})$ C_5H_5 (dppe)Mo, dppe = $Ph_2PCH_2CH_2PPh_2$) was prepared according to the methods reported in the literature.¹¹ Elemental analyses and X-ray diffraction studies were carried out at the Regional Center of Analytical Instrumentation located at the National Taiwan University.

Preparation of [Mo](CO)C=CPh (1). To a sample of [Mo]-(CO)Cl (0.22 g, 0.35 mmol) dissolved in 30 mL of THF at room temperature was added a THF solution of lithium phenylacetylide (1.0 M, 0.40 mL) via a syringe. The resulting mixture was heated under reflux for 1 h. After removal of all volatile substances in vacuo, 20 mL of Et₂O was added to the residue under nitrogen, and the extract was filtered. The solvent of the filtrate was removed under vacuum to afford the yellow product 1, which was further recrystallized from a 1:1 mixture of hexane/CH₂Cl₂ to afford crystals of 1 (0.14 g, 61% yield). Spectroscopic data for 1: IR (cm⁻¹, CH₂Cl₂) 2074 (s, $\nu_{C=C}$), 1850 (s, ν_{CO}); ¹H NMR (CDCl₃) δ 7.95–6.64 (m, 25H, Ph), 4.49 (d, $J_{\rm H-P} = 1.6$ Hz, 5H, Cp), 2.40, 1.78 (m, 4H, CH₂CH₂); ¹³C NMR (CDCl₃) δ 245.0 (d, $J_{C-P} = 22.2$ Hz, CO), 141.9–119.2 (Ph), 91.0 (Cp), 31.8 (dd, $J_{C-P} = 28.0$ Hz, $J_{C-P} = 18.6$ Hz, PCH₂), 28.6 (dd, $J_{C-P} = 22.0$ Hz, $J_{C-P} = 17.3$ Hz, PCH₂); ³¹P NMR (CDCl₃) δ 90.4 (d, $J_{P-P} = 37.2$ Hz), 78.7 (d, $J_{P-P} = 37.2$ Hz); MS (FAB, m/z, Mo⁹⁸) 690 (M⁺), 662 (M⁺ - CO). Anal. Calcd for C₄₀H₃₄OP₂Mo: C, 69.77; H, 4.98. Found: C, 70.02; H, 4.79.

Reaction of [Mo](CO)Cl with HC=CPh. To a sample of [Mo](CO)Cl (0.20 g, 0.32 mmol) dissolved in 50 mL of MeOH at room temperature was added phenylacetylene (0.21 mL, 1.82 mmol) via a syringe. The resulting mixture was heated under reflux for 2.5 h to give a green solution. After removal of all volatile substances in vacuo, 20 mL of CH₂Cl₂ was added to the residue, and the extract was filtered. The solvent of the filtrate was reduced under vacuum to about 5 mL and the solution added to a stirred solution of Et₂O to give green precipitates. The solid was collected by filtration and washed with Et₂O to give the green product {[Mo](η^2 -HC=CPh}Cl (0.20 g, 90% yield). Spectroscopic data: ¹H NMR (CDCl₃) δ 10.02 (t, $J_{H-P} = 13.8$ Hz, CH); 7.50–6.82 (m, 25H, Ph), 5.03 (s, 5H, Cp), 2.70 (m, 4H, PCH₂CH₂P); ³¹P NMR (CDCl₃) δ 79.3 (s); MS (FAB, m/z, Mo⁹⁸) 663 (M⁺). Anal. Calcd for C₃₉H₃₅P₂MoCl: C, 67.20; H, 5.06. Found: C, 67.21; H, 5.23.

Reaction of [Mo](CO)Cl with HC=CPh in the Presence of CO. To a sample of [Mo](CO)Cl (0.10 g, 0.16 mmol) dissolved in 30 mL of MeOH at room temperature was added phenylacetylene (0.11 mL, 0.92 mmol) via a syringe. The resulting mixture was heated under reflux for 2.5 h to give a green solution. After cooling to room temperature the solution was treated with 1 atm of CO gas and the solution turned yellow. After removal of all volatile substances in vacuo, 20 mL of CH₂-Cl₂ was added to the residue, and the extract was filtered. The solvent of the filtrate was reduced under vacuum to about 5 mL and the solution added to a stirred solution of hexane to give yellow precipitates, which were collected by filtration and washed with hexane to afford {[Mo](CO)(=C=CHPh}Cl, (0.12

^{(11) (}a) Staerker, K.; Curtis, M. D. *Inorg. Chem.* **1985**, *24*, 3006. (b) Lau, Y. Y. Huckabee, W. W.; Gipson, S. L. *Inorg. Chim. Acta* **1990**, *172*, 41.

g, 87% yield). Spectroscopic data: IR (cm⁻¹, CH₂Cl₂) 1909 (s, $\nu_{\rm CO}$); ¹H NMR (CDCl₃) δ 7.50–6.82 (m, 25H, Ph), 5.05 (s, 5H, Cp), 4.78 (s, 1H, =CH), 2.70 (m, 4H, PCH₂CH₂P); ³¹P NMR (CDCl₃) δ 58.5 (d, $J_{\rm P-P}$ = 18.0 Hz), 15.5 (d, $J_{\rm P-P}$ = 18.0 Hz); MS (FAB, m/z, Mo⁹⁸) 691 (M⁺), 663 (M⁺ – CO). Anal. Calcd for C₄₀H₃₅OP₂MoCl: C, 66.26; H, 4.87. Found: C, 66.41; H, 4.98. This reaction is reversible. If the MeOH solution of the product was heated under refluxing, decarbonylation occurred and the η^2 -acetylide complex was obtained in quantitative yield. The carbonylation reaction in the presence of MeONa (54 mg, 0.10 mmol in 30 mL of MeOH) gave the acetylide complex **1** in 92% yield.

Reaction of 1 with Allyl Bromide. To a sample of 1 (0.60 g, 0.87 mmol) dissolved in 50 mL of CHCl₃ at room temperature was added excess BrCH₂CH=CH₂ (0.10 mL, 1.16 mmol) via a syringe. The resulting mixture was stirred for 36 h in the dark. The color of the reaction mixture changed from yellow to brown within this period. After removal of all volatile substances in vacuo, 10 mL of CH₂Cl₂ was added to the residue under nitrogen, and the extract was filtered. The solvent of the filtrate was removed under vacuum to afford a brown residue. The residue was redissolved in 5 mL of CH₂Cl₂ and was then added dropwise to a stirred Et₂O solution to cause precipitation of brown solid, which was collected by filtration and washed with Et₂O under nitrogen and dried under vacuum to afford [[Mo](CO)=C=C(Ph)CH2CH=CH2]Br (2) (0.57 g, 81% yield). Spectroscopic data for **2**: IR (cm⁻¹, CH₂Cl₂) 1931 (s, ν_{CO}); ¹H NMR (CDCl₃) δ 7.72–6.82 (m, 25H, Ph), 5.62 (m, J_{H-H} = 17.1, 10.1, 6.3, 6.0 Hz, 1H, =CH); 5.20 (s, 5H, Cp), 5.00 (m, $J_{H-H} = 17.1$, 1.5 Hz, 1H, =CH), 4.96 (m, $J_{H-H} = 10.1$, 1.5 Hz, 1H, =CH), 3.44, 3.22 (m, 2H, PCH₂), 2.87 (ddd, $J_{H-H} = 15.6$, 6.3 Hz, $J_{H-P} = 2.0$ Hz, 1H of CH₂), 2.78 (ddd, $J_{H-H} = 15.6$, 6.0 Hz, $J_{P-H} = 2.8$ Hz, 1H of CH₂), 2.42 (m, 2H, PCH₂); ¹³C NMR (CDCl₃) δ 358.4 (dd, J_{C-P} = 36.0, 6.5 Hz, Ca), 227.4 (d, J_{C-P} = 21.8 Hz, CO), 135.0 (CH), 134.3-126.5 (Ph), 116.8 (=CH₂), 95.7 (Cp), 33.9 (CH₂), 30.1 (dd, J_{C-P} = 27.1, 13.1 Hz, PCH₂), 28.9 (dd, $J_{C-P} = 28.5$, 12.3 Hz, PCH₂); ³¹P NMR (CDCl₃) δ 71.5 (d, $J_{P-P} = 37.5 \text{ Hz}$), 67.0 (d, $J_{P-P} = 37.5 \text{ Hz}$); MS (FAB, m/z, Mo⁹⁸) 731 (M⁺), 703 (M⁺ – CO). Anal. Calcd for $C_{43}H_{39}OP_2MoBr:\ C,$ 63.79; H, 4.86. Found: C, 64.01; H, 4.97.

Reaction of 1 with Benzyl Bromide. A mixture of 1 (0.30 g, 0.43 mmol) and BrCH₂C₆H₅ (0.11 mL, 0.84 mmol) was dissolved in 40 mL of CHCl₃ at room temperature, and the resulting mixture was stirred for 36 h in the dark. The color of the reaction mixture changed from yellow to red. After removal of the solvent in vacuo, 10 mL of CH₂Cl₂ was added to the residue under nitrogen, and the extract was filtered. The solvent of the filtrate was reduced in volume to ca. 5 mL. The solution was added dropwise to a stirred Et₂O solution to give a brown precipitate. The solid was collected by filtration and washed with 5 mL of Et₂O under nitrogen. The solid was dried under vacuum to afford the brown product [[Mo](CO)= C=C(Ph)CH₂Ph]Br (3) (0.27 g, 73% yield). Spectroscopic data for 3: IR (cm⁻¹, CH₂Cl₂) 1931 (s, ν_{CO}); ¹H NMR (CDCl₃) δ 7.98– 6.73 (m, 30H, Ph), 5.09 (s, 5H, Cp), 3.42 (dd, $J_{H-H} = 14.5$ Hz, $J_{H-P} = 2.0$ Hz, 1 H on CH₂Ph), 3.27 (dd, $J_{H-H} = 14.5$ Hz, J_{H-P} = 3.7 Hz, 1 H on CH₂Ph), 3.20 (m, 2H, CH₂), 2.38 (m, 2H, CH₂); ¹³C NMR (CDCl₃) δ 356.7 (dd, J_{C-P} = 37.8 Hz, 5.6 Hz, $C\alpha$), 227.8 (d, $J_{C-P} = 20.6$ Hz, CO), 134.1–126.8 (Ph), 95.9 (Cp), 36.0 (CH_2Ph), 30.3 (dd, $J_{C-P} = 28.3$, 13.6 Hz, PCH₂), 28.5 (dd, $J_{C-P} = 28.7, 11.4 \text{ Hz}, \text{PCH}_2$; ³¹P NMR (CDCl₃) δ 69.0 (d, J_{P-P} = 36.3 Hz), 66.2 (d, J_{P-P} = 36.3 Hz); MS (FAB, m/z, Mo⁹⁸) 781 (M⁺), 753 (M⁺ – CO). Anal. Calcd for $C_{47}H_{41}OP_2MoBr$: C, 65.67; H, 4.81. Found: C, 65.60; H, 4.99.

Reaction of 1 with Bromoacetonitrile. To a sample of **1** (0.61 g, 0.87 mmol) dissolved in 40 mL of CHCl₃ at room temperature was added excess $BrCH_2CN$ (0.1 mL, 1.44 mmol) via a syringe. The resulting mixture was stirred for 24 h in the dark. The color of the reaction mixture changed from yellow to brown within 12 h. After removal of all volatile substances in vacuo, 10 mL of CH_2Cl_2 was added to the residue

under nitrogen, and the extract was filtered. The solvent of the filtrate was reduced in volume under vacuum to ca. 5 mL. The solution was added dropwise to a stirred Et₂O solution to give brown precipitate. The solid was collected by filtration and washed with Et₂O under nitrogen. The solid was dried under vacuum to afford [[Mo](CO)=C=C(Ph)CH₂CN]Br (4) (0.52 g, 74% yield). Spectroscopic data for 4: IR (cm⁻¹, CH₂-Cl₂) 1977 (s, ν_{CO}); ¹H NMR (CDCl₃) δ 7.74–6.77 (m, 25H, Ph), 5.21 (s, 5H, Cp), 3.58, 3.29, 2.52, 2.33 (m, 4H, PCH₂CH₂P), 2.95 (m, J_{H-H} = 18.2 Hz, J_{H-P} = 5.2 Hz, 1H of CH₂CN), 2.84 (m, J_{H-H} = 18.2 Hz, J_{H-P} = 5.5 Hz, 1H of CH₂CN); ³¹P NMR (CDCl₃) δ 66.8, 63.7 (2d, J_{P-P} = 30.7 Hz); MS (FAB, *m/z*) 730 (M⁺), 702 (M⁺ – CO). Anal. Calcd for C₄₂H₃₆NOBrP₂Mo (808.5): C, 62.39; H, 4.49; N, 1.73. Found: C, 62.17; H, 4.68; N, 1.76.

Reaction of 2 with NaOMe in Methanol. To a sample of 2 (0.33 g, 0.41 mmol) dissolved in 10 mL of methanol at room temperature was added a methanol solution of sodium methoxide (0.15 g, 2.78 mmol) via a cannula. An orange powder precipitated after 1 h, and the resulting solution was stirred for 4 h as the color of the reaction mixture changed from red to yellow. The product was collected by filtration and washed with 20 mL of methanol under nitrogen. The powder was dried under vacuum to afford the product $[Mo][\eta^3-C(CO_2Me)=C(Ph)-C(Ph)]$ CH₂CH=CH₂] (5) (0.25 g, 80% yield). The powder was further recrystallized from a mixture of hexane/CH2Cl2 to afford yellow crystals of 5 for X-ray diffraction analysis. Spectroscopic data for 5: IR (cm⁻¹, CH₂Cl₂) 1673 (s, ν_{CO}); ¹H NMR (C₆D₆) δ 8.25– 6.71 (m, 25H, Ph), 4.71 (s, 5H, Cp), 3.66 (d, $J_{H-H} = 17.2$ Hz, 1H, =CH₂), 3.41 (s, 3H, OCH₃), 2.82 (br, 1H, PCH₂), 2.58 (br, 1H, =CH₂), 2.52 (br, 1H, PCH₂), 2.35 (br, 1H, PCH₂), 2.02 (br, 1H, PCH₂), 1.97 (br, 1H, CH₂), 1.78 (br, 1H, =CH), 1.70 (br, 1H, CH₂); ¹³C NMR (C₆D₆) δ 180.6 (CO₂), 144.4–124.7 (Ph), 87.6 (Cp), 60.3 (dd, $J_{C-P} = 5.4$, 4.0 Hz, =CH), 49.4 (O CH_3), 47.6 (d, $J_{C-P} = 5.9$ Hz, $=CH_2$), 34.6 (CH_2), 29.1 (dd, $J_{C-P} =$ 29.9 Hz, $J_{C-P} = 15.0$ Hz, PCH₂), 27.9 (dd, $J_{C-P} = 25.8$, 14.4 Hz, PCH₂); ³¹P NMR (C₆D₆) δ 90.5 (s), 71.2 (s); MS (FAB, m/z, Mo⁹⁸) 762 (M⁺). Anal. Calcd for C₄₄H₄₂O₂P₂Mo: C, 69.47; H, 5.57. Found: C, 69.32; H, 5.79.

Reaction of 3 with NaOMe in Methanol. A mixture of 3 (0.40 g, 0.46 mmol) and sodium methoxide (0.20 g, 3.70 mmol) was dissolved in 30 mL of methanol at room temperature. The solution was stirred for 1 h, and a red product precipitated. The product was collected by filtration, washed with methanol under nitrogen, and dried under vacuum to afford the red compound [Mo](η^3 -CH(CO₂CH₃)C(Ph)CHPh) (**6**) (0.31 g, 83%) yield). The solid was further recrystallized from a mixture of Et₂O/CH₂Cl₂ to afford red crystals for diffraction analysis. Spectroscopic data for 6: IR (cm⁻¹, CH₂Cl₂) 1674 (s, ν_{CO}); ¹H NMR (CD₂Cl₂) δ 7.98-6.14 (m, 30H, Ph), 4.62 (s, 1H, syn H), 3.88 (d, $J_{H-P} = 11.4$ Hz, 1H, anti H), 3.73 (s, 5H, Cp), 3.71 (s, 3H, OCH₃), 2.68 (m, 1H, PCH₂), 2.53 (m, 1H, PCH₂), 2.47 (m, 1H, PCH₂), 1.90 (m, 1H, PCH₂); 13 C NMR (CD₂Cl₂) δ 183.2 (d, $J_{C-P} = 3.5$ Hz, CO_2), 149.0–125.0 (Ph), 92.4 (Cp), 90.3 (t, J_{C-P} = 3.1 Hz, C(Ph)), 50.2 (O CH_3), 49.2 (d, $J_{C-P} = 2.7$ Hz, CHPh), 48.7 (d, $J_{C-P} = 4.5$ Hz, $CHCO_2CH_3$), 33.7 (dd, $J_{C-P} = 28.1$, 14.3 Hz, PCH₂), 24.2 (dd, $J_{C-P} = 23.1$, 12.9 Hz, PCH₂); ³¹P NMR (CD₂Cl₂) δ 71.3 (d, $J_{P-P} = 36.8$ Hz), 72.2 (d, $J_{P-P} = 36.8$ Hz); MS (FAB, m/z, Mo⁹⁸) 812 (M⁺). Anal. Calcd for C₄₈H₄₄O₂P₂-Mo: C, 71.11; H, 5.47. Found: C, 71.06; H, 5.59.

Deprotonation of [Cp(dppe)(CO)Mo=C=C(Ph)CH₂CN]-Br by NaOMe. To a sample of **4** (0.11 g, 0.14 mmol) dissolved in 15 mL of MeOH at room temperature was added 5 mL of a MeOH solution of sodium methoxide (0.05 g, 1.9 mmol) via cannula. The color of the solution changed from red to green immediately. After removal of the solvent in vacuo, 2 mL of CH₂Cl₂ and 20 mL of hexane were added to the residue under nitrogen, and the extract was filtered. The solvent of the filtrate was removed under vacuum to afford a green residue. The residue was redissolved in 2 mL of CH₂Cl₂. The solution was added dropwise to a stirred Et₂O solution to give green

Table 3. Crystal and Intensity Collection Data for Cp(dppe)MoC(COOMe)=C(Ph)CH₂C=CH₂ (5) and Cp(dppe)MoCH(COOMe)C(Ph)CHPh (6)

	5	6
mol formula	C45H44O2P2Cl2M0	C48H44O2P2M0
mol wt	845.63	810.71
space group	$P2_{1}/c$	$P\bar{1}$
a, Å	17.998(5)	9.6178(1)
<i>b</i> , Å	8.647(3)	10.8305(2)
<i>c</i> , Å	26.080(5)	21.3254(3)
α, deg		96.911(1)
β , deg	100.30(2)	94.388(1)
γ , deg		113.258(1)
V, Å ³	3993.6(19)	2007.22(5)
Z	4	2
cryst dimens, mm ³	$0.30\times0.35\times0.40$	$0.20\times0.25\times0.35$
Mo K α radiation: γ , Å		0.71073
θ range for data collection	0.55 - 25.0	0.97 - 25.0
limiting indices (<i>h</i> , <i>k</i> , <i>l</i>)	-21, 21; 0, 10; 0, 30	-12, 12; -14, 14; -28, 28
no. of reflns collected	7015	16793
no. of ind reflns	4949	7071
max. and min. transmn	0.868 and 0.819	0.492/0.408
refinement method	full-matrix leas	t-squares on F^2
no. of data/restraints/	4949/0/470	7071/0/479
params		
GÓF	1.55	1.196
final R indices		
$[I > 2\sigma(I)]$	0.034/0.037	0.0411/0.1386
all data		0.0451/0.1434
$\Delta \rho$ (in final map), e/Å	-0.470, +0.490	-0.846, +0.507

precipitates. The solid was collected by filtration, washed with Et₂O under nitrogen, and dried under vacuum to afford a green

solid, Cp(dppe)(CO)Mo-C=C(Ph)CHCN (7) (0.052 g, 53% yield). Spectroscopic data for 7: IR (cm⁻¹, CH₂Cl₂) 1846 (s, ν_{CO}); ¹H NMR (CDCl₃) δ 7.81–6.91 (m, 25H, Ph), 4.78 (s, 5H, Cp), 3.73, 3.06 (2m, 4H, PCH₂CH₂P), 2.51 (s, 1H, CHCN); ³¹P NMR (CDCl₃) δ 105.4 (s), 103.9 (s); MS (FAB, m/z) 729 (M⁺), 701 (M⁺ – CO). Anal. Calcd for C₄₂H₃₅NOP₂Mo: C, 69.33; H, 4.85; N, 1.93. Found: C, 69.97; H, 5.35, N, 2.64.

X-ray Structure Determination of 5 and 6. Yellow crystals of 5 suitable for X-ray diffraction study were grown directly from a mixture of hexane/CH₂Cl₂. A single crystal of dimensions $0.30 \times 0.35 \times 0.40 \text{ mm}^3$ was glued to a glass fiber and mounted on an Nonius CAD-4 diffractometer. Data were collected and processed, and crystallographic computations were carried out using the NRCC structure determination package.¹² The structure was solved using direct methods¹³ and was refined on intensities of 4949 reflections to give R =0.034, $R_{\rm w} = 0.037$ ($I > 2\sigma(I)$). For **6**, a single crystal of dimensions $0.20 \times 0.25 \times 0.35 \text{ mm}^3$ was mounted on a glass fiber with epoxy. Data were collected at room temperature on a Siemens SMART CCD area detector system employing a 3 kW sealed tube X-ray source operating at 1.5 kW. The total data collection was approximately 6 h, yielding 7071 independent data after integration using SAINT.¹⁴ Unit cell parameters were determined from the least-squares refinement of three-dimensional centroids of unique reflections. Data were corrected for absorption with the SADABS program.¹⁵ The space group was assigned as P1, and the structure was solved and refined using direct methods included in the SHELXTL package.¹⁶ In the final model, non-hydrogen atoms were refined anisotropically, with hydrogen atoms included in idealized locations. The structure was refined to R1 = 0.0411 and wR2 = 0.1386 for $I > 2\sigma(I)$ and to R1 = 0.0451 and wR2 = 0.1434for all data.¹⁷ Crystal and intensity collection data for 5 and 6 are given in Table 3, and fractional coordinates and thermal parameters are given in the Supporting Information.

Acknowledgment. Financial support from the National Science Council, Taiwan, is gratefully acknowledged.

Supporting Information Available: Details of the structural determination for complexes 5 and 6, including crystal and intensity collection data, positional and anisotropic thermal parameters, and all of the bond distances and angles. This material is available free of charge via the Internet at http://pubs.acs.org.

OM9907167

(12) Gabe, E. J.; Lee, F. L.; Lepage, Y. In *Crystallographic Computing 3*; Sheldrick, G. M., Kruger, C., Goddard, R., Eds.; Clarendon Press: Oxford, England, 1985; p 167.
(13) Sheldrick, G. M. *SHELXS-86, Program for Crystal Structure Solution*; University of Gottingen: Gottingen, Germany, 1986.

(14) SAINT (Siemens Area Detector Integration) program; Siemens Analytical X-ray: Madison, WI, 1995.

Analytical X-ray: Madison, WI, 1995. (15) The SADABS program is based on the method of Blessing; see: Blessing, R. H. *Acta Crystallogr., Sect. A* **1995**, *51*, 33. (16) *SHELXTL*: Structure Analysis Program, version 5.04; Siemens Industrial Automation Inc.: Madison, WI, 1995. (17) GOF = $[\sum [w(F_o^2 - F_o^2)^2]/(n - p)]^{1/2}$, where *n* and *p* denote the number of data and parameters. R1 = $(\sum ||F_o| - |F_c||)/\sum |F_o|$, wR2 = $[\sum [w(F_o^2 - F_o^2)^2]/\sum [w(F_o^2)^2]^{1/2}$, where $w = 1/[\sigma^2(F_o^2) + (aP)^2 + bP]$ and $P = [(max; 0, F_o^2) + 2F_o^2]/3$.