

A Novel Photochromic System of 4,5-Dialkenylthiophenes Constructed by the Samarium Diiodide Promoted Coupling Reactions of Thiophene-2-carboxylate with Aryl Ketones


Jiun-Jie Shie, Shyh-Ming Yang, Chao-Tsen Chen, and Jim-Min Fang*

Department of Chemistry, National Taiwan University, Taipei 106,
Taiwan, Republic of China

jmfang@ccms.ntu.edu.tw

Received December 20, 2001

ABSTRACT



The SmI₂-promoted coupling reaction of ethyl thiophene-2-carboxylate with aryl ketones (2 equiv), followed by acid-catalyzed dehydration and oxidative aromatization, gave dialkenylthiophenes **1b–d**, which underwent electrocyclizations upon irradiation with 300-nm light in CH₃CN solution to give the corresponding closed-ring species with absorption $\lambda_{\text{max}} \approx 425$ nm. The interconversion between dialkenylthiophenes and their corresponding closed-ring species constitutes a novel photochromic system bearing an ester group for potential uses in linkage and wavelength tuning.

Photochromism is a light-induced reversible isomerization between two forms having different absorption spectra. The photochromic system based on the interconversion of 1,3,5-hexatriene to cyclohexadiene has been extensively investigated,¹ partly as a result of their potential applications to optical memories and switches. The photocyclization of stilbene to dihydrophenanthrene is a well-known photochromic system.² According to the Woodward–Hoffmann rule,³ the double bond of stilbene should have (*Z*)-configuration

to conform a concerted conrotatory cyclization in the photochemical conditions. Dihydrophenanthrene can return rapidly to stilbene in the dark; however, it can also undergo oxidative aromatization in the presence of oxygen.² By replacing the phenyl rings with thiophene rings, 1,2-di(3-thienyl)ethene undergoes a photocyclization reaction to give the closed-ring species with an improved stability.⁴ This


(1) For reviews, see: (a) Laarhoven, W. H. In *Photochromism: Molecules and Systems*; Dürr, H., Bouas-Laurent, H., Eds.; Elsevier: Amsterdam, 1990; Chapter 7, pp 270–313. (b) Irie, M.; Uchida, K. *Bull. Chem. Soc. Jpn.* **1998**, *71*, 985. (c) *Organic Photochromic and Thermochromic Compounds*; Crano, J. C., Guglielmetti, R. J., Eds.; Plenum Press: New York, 1999. (d) Irie, M. *Chem. Rev.* **2000**, *100*, 1685.

(2) Muszkat, K. A.; Fischer, E. *J. Chem. Soc. C* **1967**, 662. (e) Repinec, S. T.; Sension, R. J.; Szarka, A. Z.; Hochstrasser, R. M. *J. Phys. Chem.* **1991**, *95*, 10380.


(3) (a) Dewar, M. J. S. *Angew. Chem., Int. Ed. Engl.* **1971**, *10*, 761. (b) Zimmerman, H. E. *Acc. Chem. Res.* **1971**, *4*, 272. (c) Nakamura, S.; Irie, M. *J. Org. Chem.* **1988**, *53*, 6136.

(4) Kellogg, R. M.; Groen, M. B.; Wynberg, H. *J. Org. Chem.* **1967**, *32*, 3093.

result is attributed to the low aromatic stabilization energy difference between the open- and closed-ring isomers of thiophene by comparison with that of benzene.^{1b,3c} Irie and co-workers further demonstrate that the light-induced cyclization of 1,2-bis(2,4-dimethylthien-4-yl)perfluorocyclopentene gives a stable colored species.^{1b} The cyclopentene ring incorporates the requisite double bond of the (Z)-configuration, and the methyl groups at the 2- and 2'-positions prevent the closed-ring species from oxidative aromatization.



We report herein a novel photochromic system (eq 1) based on the interconversion between 4,5-dialkenylthiophenes **1a–d** and their closed-ring isomers **2a–d**. Dialkenylthiophene **1a–d** were readily prepared by a three-step sequence (Scheme 1): (i) coupling of ethyl thiophene-




^a Reagents and conditions: (i) SmI₂, THF, HMPA, 25 °C, 12–24 h. **8**, 50% yield; **9**, 43%; **10**, 32%; **11**, 37%. (ii) Cat. *p*-TsOH, PhH, reflux, 4–12 h. **12**, 95% yield; **13**, 87%; **14**, 67%; **15**, 73%. (iii) For **1a**, DDQ, PhH, 60 °C, 12 h; 83% yield. For **1b–d**, DDQ, PhCH₃, reflux, 24 h; **1b**, 69% yield; **1c**, 80%; **1d**, 84%.

2-carboxylate (**3**) with aryl ketones **4–7** (2 equiv) by the promotion of SmI₂, giving diols **8–11**,⁵ (ii) acid-catalyzed dehydration, giving dienes **12–15**, and (iii) oxidation by DDQ (1 equiv), giving dialkenylthiophenes **1a–d**. The

central double bonds of **1a–d** were confined in the thiophene ring to adopt a (Z)-configuration required by the concerted electrocyclizations.

The UV–vis spectrum of dialkenylthiophene **1a**⁶ exhibited λ_{max} at 242 nm (ε = 38800) and 289 nm (ε = 13800). Upon irradiation of **1a** (colorless solution) with 300-nm light, an orange color immediately appeared, indicating the existence of new species. The reaction course of **1a** in CDCl₃ (ca. 0.1 M) was recorded by the ¹H NMR spectra.⁷ After irradiation for 10 min, closed-ring species **2a** (ca. 30%) was produced as inferred from a new set of proton signals, in which a singlet at δ 2.99 was ascribed to the methylene protons of tetrahydrobenzothiophene moiety. The ratio of **2a** increased as the irradiation progressed to 60 min. However, several other compounds including benzothiophene **16**⁸ and 6,7-dihydrobenzothiophene **17**⁹ were also produced at the expense of **2a** (Scheme 2).



Compound **16** was likely obtained by an oxidative aromatization of **2a** similar to that proposed in the conversion

(5) (a) Yang, S.-M.; Fang, J.-M. *Tetrahedron Lett.* **1997**, 38, 1589. (b) Yang, S.-M.; Nandy, S. K.; Selvakumar, A. R.; Fang, J.-M. *Org. Lett.* **2000**, 2, 3719.

(6) Compounds **1a–d** and **8–18** were fully characterized by spectral methods (IR, UV, MS, HR-MS, ¹H and ¹³C NMR). The details are reported in Supporting Information, whereas some pertinent physical and spectral properties are herein listed. Compound **1a**: solid, mp 109–111 °C; UV (CHCl₃) λ_{max} (ε) 242 nm (38800), 289 nm (13800); FL (CHCl₃, c = 2 × 10^{−4} M) λ_{em} 415 nm on excitation at 364 nm; ¹H NMR (CDCl₃, 300 MHz) δ 7.77 (1 H, s), 7.73–7.70 (2 H, m), 7.58–7.54 (4 H, m), 7.44–7.36 (6 H, m), 7.16 (1 H, dd, *J* = 8.5, 1.5 Hz), 7.10 (1 H, dd, *J* = 8.5, 1.5 Hz), 5.57 (1 H, s), 5.50 (1 H, s), 5.41 (1 H, s), 5.35 (1 H, s), 4.38 (2 H, q, *J* = 7.3 Hz), 1.39 (3 H, t, *J* = 7.3 Hz). Compound **1b**: UV (CHCl₃) λ_{max} (ε) 290 nm (35400). Compound **1c**: UV (CH₃CN) λ_{max} (ε) 249 nm (34400), 300 nm (16800). Compound **1d**: UV (CH₃CN) λ_{max} (ε) 251 nm (34900), 300 nm (13700).

(7) The photochemical reactions were conducted in a Rayonet photochemical reactor using 300 or 450-nm lamps. The corresponding NMR and UV–vis time course spectra are included in Supporting Information.

of dihydrophenanthrene to phenanthrene. When the photochemical reaction of **1a** was conducted in deoxygenated CD₃CN solution, the formation of **16** was inhibited. Compound **17** might be derived from **2a** by an H-shift. Attempts to isolate **2a** failed, presumably because of its instability. Attempts to trap the transient closed-ring species **2a**, as an *o*-thiophenequinodimethane, by *N*-phenylmaleimide in the photochemical reaction of dialkenylthiophene **1a** (300-nm, CH₃CN, 25 °C, 48 h) also failed, even though the existence of **2a** could be inferred from a newly occurring NMR signal at δ 2.99 (s). However, **2a** was alternatively generated by the thermal electrocyclic of **1a** in refluxing toluene and successfully trapped by *N*-phenylmaleimide to give a [4 + 2] cycloaddition product **18**.¹⁰ Although the trapping experiment under thermal conditions should not be strictly correlated to the photochemical reaction of **1a**, this result lent a clue to intermediate **2a** in support of the newly NMR signal at δ 2.99 found under photochemical conditions.

The drawbacks of oxidative aromatization and H-shift would be avoided in the photochemical cyclization of trienes **1b–d** as they have C-5 and C-6 positions substituted by methyl groups. When a colorless acetonitrile solution (1.1×10^{-5} M) of **1b** was irradiated with 300-nm light at room temperature, the 290-nm absorption of **1b** decreased along with the growth of a new 429-nm absorption attributed to the formation of the yellow closed-ring species **2b** (Figure 1). The corresponding ¹H NMR spectra⁷ also showed a newly




Figure 1. The UV-vis spectral change of compound **1b** on irradiation with 300-nm light at 25 °C in CH₃CN solution.

occurring singlet (δ 6.64) ascribed to the H-3 of **2b**. According to the NMR study, the maximal content of the

closed-ring species **2b** was estimated to be 15% in the reaction mixture under the photochemical conditions (30 min at -20 °C in CD₃CN solution).

The photochemical properties of **1c** (Ar = Ph) and **1d** (Ar = *p*-MeOC₆H₄) were similar to that of **1b** (Ar = 2-naphthyl), even though they had different aryl groups. Irradiation of **1c** and **1d** with 300-nm light in CH₃CN solution afforded the closed-ring species **2c** and **2d**, which showed absorption maxima at 422 and 425 nm, respectively.⁷ On the basis of these results, the moiety of *o*-thiophenequinodimethine carboxylate in **2b–d** was likely the main contributing chromophore. The absorption change in **2b–d** was trivial, as their aryl substituents might not be coplanar to the chromophore.

We also demonstrate that the closed-ring species **2b–d** are photochemically reversible.¹¹ When a mixture of **1b/2b** and **1d/2d** was irradiated with 450-nm light, respectively, **2b** and **2d** disappeared and returned rapidly (within 1 min) to the open-ring species (**1b** and **1d**). The interconversion between **1c** and **2c** was realized by alternate irradiation with UV (300-nm) and visible (419-nm) lights at 10-s intervals. In a preliminary experiment, the absorption intensity at 424 nm showed a significant change after irradiation of **1c/2c** for several cycles (see Supporting Information).

The ring-closure and ring-opening reactions as shown in eq 1 should be both photochemically and thermally allowed. The closed-ring species were unstable on standing at room temperature, and they returned to the open-ring species in the dark. For example, a mixture of **1b/2b** (ca. 85:15) obtained by irradiation of **1b** with 300-nm light for 40 s in CH₃CN was removed from the light source and allowed to stand at room temperature in the dark. According to the absorption intensity changes at 290 and 429 nm, **2b** returned to **1b** to a considerable degree (~20%) in the first minute and **2b** completely disappeared in 40 min.⁷ A mixture of **1c/2c** (ca. 82:18) also behaved similarly to give exclusively **1c** after standing at room temperature for 24 h in the dark. This interconversion was monitored by ¹H NMR spectra, in which the characteristic H-3 signal of **1c** (singlet at δ 7.52) increased as that of **2c** (singlet at δ 6.22) disappeared.⁷

In summary, the interconversion between dialkenylthiophenes **1b–d** and their corresponding closed-ring species constitutes a novel photochromic system. The preparation of **1b–d** is simple by using readily available starting materials via a three-step sequence (Scheme 1), though the yield is not yet optimized. The ester group in trienes **1b–d** may serve as a linker to polymeric supports or undergo functional group transformation to tune the photochemical properties. The closed-ring species **2b–d** are unstable and return to **1b–d** in the dark even at room temperature. The photochemical and thermal stability of these substrates must be improved in order to reach a practical application of this novel dialkenylthiophene/dihydrobenzothiophene system.

The elegant diarylethene photochromic system developed

(8) For comparison, an authentic sample of **16** was also prepared by treatment of triene **1a** with DDQ in refluxing toluene. Compound **16** exhibited a characteristic H-3 at δ 8.27 (s).

(9) Compound **17** was isolated by chromatography of the reaction mixture at the interval of 90-min irradiation of **1a**. The structure of **17** was tentatively assigned according to the ¹H NMR, HR-MS, and NOE analyses. There is no enhancement of H-3 at δ 7.68 (s) on irradiation of H-7 at δ 4.53 (t, *J* = 9 Hz).

(10) Compound **18** had the *endo*-configuration as its NOESY spectrum showed a correlation of H-4a/H-7a (δ 4.34–4.33, m) with the ethylene-bridge protons (δ 2.15–2.11, m).

(11) Miyasaka, H.; Murakami, M.; Itaya, A.; Guillaumont, D.; Nakamura, S.; Irie, M. *J. Am. Chem. Soc.* **2001**, *123*, 753.

by Irie and co-workers^{1b,d} has demonstrated high quantum yield, thermal stability, and fatigue resistance. This system incorporates a central ring of perfluorocyclopentene with substitution of two aromatic rings. When the open isomer changes to the closed isomer upon irradiation, the two aromatic rings must change simultaneously to form another conjugated chromophore. In contrast, our current photochromic system utilizes a thiophenyl ring as the central moiety. The synthesis of the photo-/thermochromic substrates of dialkenylthiophenes is relatively easy by using the commercially available starting materials of thiophene-2-carboxylate and aryl ketones. Furthermore these two aryl groups are not incorporated into the contributing *o*-thiophenequinodimethane chromophore of the closed-ring species. This unique structure bearing two independent chromophores thus provides a good opportunity to build the photochromic system having a fluorescent sensing property. For example,

one can replace the naphthyl ring in **1b** by an appropriate coumarin as the acceptor chromophore for the light emitted from the closed-ring species, which is generated by UV-light irradiation. Modification of the ketone moieties in our current system would not be difficult. We are currently engaged in this endeavor to demonstrate such energy-transfer mechanism in a photochromic system.

Acknowledgment. We thank the National Science Council for financial support and Professor Pi-Tai Chou (National Taiwan University) for helpful discussion.

Supporting Information Available: Detailed experimental procedures, NMR and UV–vis spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0172830