



Cite this: *Food Funct.*, 2019, **10**, 1767

DOI: 10.1039/c9fo90006c

rsc.li/food-function

Correction: From white to beige adipocytes: therapeutic potential of dietary molecules against obesity and their molecular mechanisms

Siyu Wang,^a Min-Hsiung Pan,^b Wei-Lun Hung,^c Yen-Chen Tung^b and Chi-Tang Ho^{*a}

Correction for 'From white to beige adipocytes: therapeutic potential of dietary molecules against obesity and their molecular mechanisms' by Siyu Wang *et al.*, *Food Funct.*, 2019, DOI: 10.1039/c8fo02154f.

In the original manuscript, an incorrect version of Fig. 1 was shown. The correct figure and caption are as follows.




Fig. 1 Distribution of three types of adipocytes and their molecular characteristics in rodents and humans. The dotted lines indicate inducible characteristics of beige adipocytes.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aDepartment of Food Science, Rutgers University, New Brunswick, NJ 08901, USA. E-mail: ctho@sebs.rutgers.edu; Tel: +848-932-5553

^bInstitute of Food Science and Technology, National Taiwan University, Taipei 10617, Taiwan

^cSchool of Food Safety, Taipei Medical University, Taipei 11031, Taiwan

