



Black tea in chemo-prevention of cancer and other human diseases

Min-Hsiung Pan^{a,*}, Ching-Shu Lai^a, Hong Wang^b, Chih-Yu Lo^c, Chi-Tang Ho^d, Shiming Li^{e,**}

^a Department of Seafood Science, National Kaohsiung Marine University, Kaohsiung 811, Taiwan, China

^b Shaklee Corporation, 4747 Willow Road, Pleasanton, CA 94588, USA

^c Department of Food Science, National Chiayi University, Chiayi City 60004, Taiwan, China

^d Department of Food Science, Rutgers University, New Brunswick, NJ 08901, USA

^e Hubei Key Laboratory of Economic Forest Germplasm Improvement and Resources Comprehensive Utilization, Huanggang Normal University, Huanggang 438000, China

Received 21 February 2013; received in revised form 1 March 2013; accepted 16 March 2013

Abstract

Tea is the most popular functional beverage in the world and has been gaining more and more attention for its health beneficial properties. Among common teas, black tea is consumed more than green tea and oolong tea worldwide. Numerous studies have shown the biological activities of black tea and its polyphenols that include anti-oxidant, anti-tumor, anti-inflammation and metabolic regulation. Tea polyphenols such as theaflavins and catechins are considered to be multifunctional compounds that could be effective in the prevention or treatment of various cancers, cardiovascular disease, chronic inflammation, obesity, metabolic syndrome, and neurodegenerative diseases. In this review, we summarized the up-to-date research and underlying molecular mechanisms of black tea and its polyphenols.

© 2013 Beijing Academy of Food Sciences. Production and hosting by Elsevier B.V. All rights reserved.

Keywords: Black tea; Theaflavins; Catechins; Human disease; Molecular mechanism; Cancer

1. Introduction

As tea drinking has become part of a phenomenal global culture nowadays, it is mainly attributed to its biological functionality in contrast to the traditional tea drinking, which is mostly driven by its flavor and stimulant effect. Indeed, polyphenolic compounds in tea have many health promoting activities due to their diverse biological actions including antioxidant, anti-inflammatory, anti-tumor and metabolic regulatory effects that could lead to cancer prevention and protect against metabolic, cardiovascular and inflammatory diseases.

Since black tea consumption accounts for 78% of the overall tea beverage industry [1], we directed our attention to the studies of black tea in both its chemistry and biological activities. In our previous review, we focused on the chemistry of black tea: the formation, composition, analysis and stability of black tea [1]. From the reported data and our analysis, we concluded that the major polyphenols in black tea are composed of theaflavins (Table 1) and catechins (Table 2). Black tea is different from other tea such as green tea, white tea and oolong tea not only in their processing and flavor, but also in their chemical compositions. Chemical analysis showed that black tea also contains high amount of green tea catechins in addition to theaflavins. Therefore, both catechins and theaflavins contribute to the bioactivity of black tea either by additive and/or synergistic effects.

Numerous articles have summarized the biological properties of green tea and its catechins and a few have reviewed bioactivity of black tea in general. However, few articles have attributed the bioactivity data of black tea to its chemical profiles. In this review, we summarized the health beneficial properties of black tea and also listed the bioactivities of individual theaflavins (TFs), the key identity marker of black tea. By describing various health promoting effects of black tea and the potential mechanisms from its major ingredients, we would like to advocate the importance of chemical profiling in natural product research and functional food products development.

* Corresponding author at: Department of Seafood Science, National Kaohsiung Marine University, No. 142, Haijhuang Rd., Nanzih District, Kaohsiung 81143, Taiwan, China. Tel.: +886 7 361 7141x3623.

** Corresponding author.

E-mail addresses: mhpan@mail.nkmu.edu.tw (M.-H. Pan),


shiming3702@yahoo.com (S. Li).

Peer review under responsibility of Beijing Academy of Food Sciences.




Production and hosting by Elsevier

Table 1
Chemical structure of theaflavins.



Compound	R	R'
Theaflavin (TF1)	H	H
Theaflavin-3-monogallate (TF2a)	Galloyl	H
Theaflavin-3'-monogallate (TF2b)	H	Galloyl
Theaflavin-3,3'-digallate (TF3)	Galloyl	Galloyl

Table 2
Chemical structure of catechins.



Compound	R	R'
Epicatechin (EC)	H	H
Epigallocatechin (EGC)	H	OH
Epicatechin gallate (ECG)	Galloyl	H
Epigallocatechin gallate (EGCG)	Galloyl	OH

2. Cancer prevention and anti-tumor effects of black tea

Cancer is the most common cause of human mortality worldwide. Cancer cells are capable of limitless replication potential due to their self-sufficient growth signals and their resistant to anti-growth signals from host defense that enable them to evade apoptosis. Furthermore, they can sustain angiogenesis that can lead to tissue invasion and metastasis [2]. Targeting the regulation of key molecules and cellular signaling pathways in tumorigenesis by natural products was effective in preventing or reducing risk of cancer [3]. There are many reports on the efficacy of the anti-tumor and cancer prevention activity of black tea. For instance, studies from Baker et al. [4] and Doral et al. [5] support the cancer chemo-preventive effect of black tea in the development of prostate cancer, ovarian cancer and rectal cancer. However, others [6,7] do not. The discrepancy of the effects could be a result of actual dosage differences among studies with unspecified amount ingestion of bioactive tea polyphenols. Daily consumption of black tea in women lowered the concentration of 17 β -estradiol (E2), which may reduce hormone-related cancer risk [8] (Table 3). Regular black tea consumption is also associated with the reduced risk of ovarian and bladder cancer in female subjects [9–11]. Furthermore, several research groups

have attempted to elucidate the molecular mechanisms of black tea and its polyphenols.

Drinking black tea reduced incidence and number of skin papilloma in 7,12-dimethylbenz[a]anthracene (DMBA)-treated mice through activation of detoxification enzymes and decreased lipid peroxidation [12]. Oral administration of black tea polyphenols delayed tumorigenesis, reduced tumor number and volume in DMBA-induced mouse skin carcinogenesis through induction of apoptosis in tumor cells [13]. Topical application of combined black tea polyphenols and resveratrol synergistically inhibited DMBA/TPA-induced skin carcinogenesis by reducing tumor incidence, number and volume. Mechanistic study showed that this combination down-regulated mitogen-activated protein kinases (MAPKs) and increased tumor suppressor gene *p53* and apoptosis [14]. Consistent with the results in skin cancer model, oral intake of black tea polyphenols or extract also suppressed DMBA-induced mammary tumors and oral tumors by scavenging reactive oxygen species (ROS) that reduced the oxidative stress [15] and down-regulating cyclooxygenase-2 (COX-2), nuclear factor kappa-B (NF- κ B) and protein kinase B (Akt) [15], and interfering with the activity of carcinogen metabolizing enzymes [16].

In DMH (1,2-dimethylhydrazine)-induced colorectal tumor model, consumption of tea with high content polymeric black tea polyphenols inhibited the tumorigenesis *via* down-regulation of Wnt/ β -catenin pathway and proliferative gene expression [17]. Oral administration of black tea polyphenols was also effective against arsenic-induced formation of 8-hydroxy-2'-deoxyguanosine (8-OHdG) through up-regulation of DNA repair enzymes in Swiss albino mice [18].

One common feature in the effects of black tea in antagonizing various chemical-induced carcinogenesis is the activation of the detoxification enzymes. The detoxifying enzyme system plays an important role in determining the final fate of carcinogens/procarcinogens and their subsequent impact on carcinogenesis [19]. Regulation of many detoxifying enzymes is mediated by the transcription factor nuclear factor E2-related factor 2 (Nrf2) that binds to the antioxidant response element (ARE)/electrophile response element (EpRE), which is located in the promoter region of related genes to initiate gene expression. Ingredients in black tea including EGCG activate Nrf2 and up-regulate the protective enzymes [19,20]. In addition to Nrf2 activation, there are many other important mechanisms that contribute to the anti-carcinogenic effects by black tea polyphenols.

Theaflavins (TFs) are major bioactive components in black tea and are classified as theaflavin (TF1), theaflavin-3-*O*-gallate (TF2a), theaflavin-3'-*O*-gallate (TF2b) and theaflavin-3,3'-*O*-digallate (TF3) (Table 1). TFs' treatment of human leukemic U937 and K562 cells suppressed Akt signaling, thus blocked Wnt/ β -catenin signaling, cyclin D1 level and increased levels of FOXO1 and p27 that contribute to induce G0/G1 cell-cycle arrest [21]. In HeLa cervical cancer cells, TFs caused apoptosis by generation of ROS, increase of p53 and down-regulation of COX-2 and cyclin D1 *via* blockage of Akt and NF- κ B activation [22].

While some of the actions may come from the combinations of TFs, an individual TF has its individual activities against cancer cells. TF1 induced ROS-mediated apoptosis

Table 3

Anti-tumor and cancer chemopreventive effect of black tea and its bioactive compounds.

	Molecular mechanism	Ref.
Catechins	Reduced the concentration of 17 β -estradiol (E2) in women	[8]
	Inhibited 7,12-dimethylbenz[a]anthracene (DMBA)-induced skin tumorigenesis through activation of superoxide dismutase (SOD) and catalase (CAT) as well as induced apoptosis in mouse skin tumors	[12]
Black tea	Induction of apoptosis by tea polyphenols mediated through mitochondrial cell death pathway in mouse skin tumors	[13]
	Inhibited DMBA/12- <i>O</i> -tetradecanoylphorbol-13-acetate (TPA)-induced skin carcinogenesis by down-regulation of mitogen-activated protein kinases (MAPKs) and increase of p53 in a resveratrol-synergistically manner	[14]
	Suppressed DMBA-induced mammary tumors by scavenging of reactive oxygen species (ROS), down-regulation of cyclooxygenase-2 (COX-2), nuclear factor- κ B (NF- κ B) and Akt	[15]
	Suppressed 1,2-dimethylhydrazine (DMH)-induced colonic tumorigenesis by inhibition of cyclin D1, c-myc and COX-2 gene expression through blockage of Wnt/ β -catenin pathway	[17]
	Against arsenic-induced oxidative damage and up-regulation of DNA repair enzymes in Swiss albino mice	[18]
Black tea extract	Against oxidative stress, interfered with the activity of carcinogen metabolizing enzymes and inhibited DMBA-induced tumor in hamster buccal pouch carcinogenesis model	[16]
	Induction of apoptosis through increasing reactive oxygen species (ROS) production and activation of c-Jun N-terminal kinases (JNK)-p38 MAPK signaling.	[23]
	Suppression of MMP-2 gene expression and activity by down-regulation of epidermal growth factor receptor (EGFR) and NF- κ B signaling	[24]
TF1	Induced apoptosis through Fas death receptor/caspase-8 cascade and inhibited pAkt/pBad survival pathway in p53-mutated human breast cancer cells	[25]
	Inhibited gene and protein expression of fatty acid synthase (FAS) in MCF-7 breast cancer cells	[26]
	Against lung carcinogenesis through induction of apoptosis and inhibition of COX-2 expression in mice	[27,28]
	Inhibited bronchiolar cell proliferation and tumor formation in 4-(methyl nitrosamino)-1-(3-pyridyl)-1-butanone (NNK)-induced lung carcinogenesis in A/J mice	[29]
	Inhibited cell proliferation by suppression of proteasome activity	[30]
	Induction of apoptosis in human colon cancer cell line and repressed TPA-induced ear edema in mice	[31]
TF2	Inhibited cell proliferation, induced apoptosis and down-regulated COX-2 gene expression in cancer cells	[32]
	Induced apoptosis through up-regulation of p53 and Bax in WI38VA and HeLa cells	[31]
	Induction of apoptosis through cytochrome c-mediated cascade in U937 cells	[33]
	Inhibited cell proliferation through suppression of phosphorylated EGFR and platelet-derived growth factor (PDGF)	[35]
TF3	Repressed EGFR expression through protein endocytosis and degradation	[36]
	Induced apoptosis by increasing oxidative stress in human oral squamous cells	[37]
	Induced cell cycle arrest combined with ascorbic acid in human lung adenocarcinoma cells	[38]

in human malignant melanoma A375 cells through c-Jun N-terminal kinases (JNK) – p38 mitogen-activated protein kinase (MAPK) cascade [23]. In a human melanoma cell line, TF1 suppressed enzyme activity and gene and protein expression of metalloproteinases (MMP)-2 by down-regulating epidermal growth factor receptor (EGFR), extracellular regulated protein kinase (ERK) and NF- κ B signaling pathways [24]. Mutation or inactivation of tumor suppressor gene p53 is frequently found in breast cancer cells and contributes to drug resistant. TF1 induced apoptosis through activating the Fas death receptor/caspase-8 pathway and inhibiting the pAkt/pBad survival pathway in p53-mutated human breast cancer cells [25]. Moreover, TF1 inhibited both gene and protein expression of fatty acid synthase (FAS) that could reduce cell lipogenesis and proliferation in MCF-7 breast cancer cells [26]. Drinking TF1 in water daily suppressed benzo(a)pyrene (BP)-induced lung carcinogenesis through induction of apoptosis and inhibition of cyclooxygenase-2 (COX-2) expression [27,28] and reduced 4-(methyl nitrosamino)-1-(3-pyridyl)-1-butanone (NNK)-induced cell proliferation and tumor formation in mice [29]. TF1 was effective in inhibiting 20S proteasome and in reducing proliferation in human multiple myeloma (Arp and Opm1) cells [30].

TF2, which includes TF2a and TF2b, triggered mitochondria-mediated apoptosis signaling in human colon cancer cells. It also suppressed TPA (12-*O*-tetradecanoylphorbol-13-acetate) induced COX-2 gene expression *in vitro* and ear edema *in vivo* [31]. Treatment with TF2 in SV40 transformed WI38 human cells (WI38VA) caused cell growth inhibition, induction of apoptosis as well as down-regulation of COX-2 gene expression [32]. TF2 up-regulated p53 and Bax gene expression that resulted in apoptosis in WI38VA and HeLa cells [31]. TF2 also exhibited strong growth inhibitory effect in human histolytic lymphoma U937 cells [33].

Among theaflavins, theaflavin digallate (TF3) also has potent anti-cancer and chemo-preventive properties. TF3 is a potent inhibitor of DNA methyltransferases (DNMT3a), an enzyme in which its upregulation is involved in silencing tumor suppressor genes [34]. TF3 showed stronger activity in inhibiting cell proliferation than EGCG in A431 cells and mouse NIH3T3 fibroblast cells by reducing the auto-phosphorylation of EGF and platelet-derived growth factor (PDGF) receptors [35]. It also reduced EGF-induced EGFR expression by induction of protein endocytosis and degradation [36] and acted as an inducer of oxidative stress that caused apoptosis in human oral squamous HSC-2 cells [37]. In combination with ascorbic acid, TF3 showed a

Table 4

Protective effect of black tea and its bioactive compounds on cardiovascular disease (CAD).

Compound	Molecular mechanism	Ref.
Black tea	Association with reduced risk of CAD by reduction of C-reactive protein, glucose, triglyceride and increase of plasma antioxidant	[41,42]
TF1, TF2a, TF2b, TF3	Inhibited Cu ²⁺ -mediated low-density-lipoprotein (LDL) oxidation <i>in vitro</i>	[45]
TF1	Attenuates atherosclerosis in apolipoprotein E (ApoE)-knockout mice through reduction ROS and inflammatory cytokines production.	[47]
	Recovered cardiac function after ischemia/reperfusion through inhibition of mitochondrial permeability transition pore (mPTP) opening	[46]
Black tea extract	Potent inhibitors of the acetyltransferase and 1-alkyl-2-acetyl-sn-glycero-3-phosphocholine (PAF) biosynthesis that inhibited platelet aggregation.	[44]
	Protected tert-butyl hydroperoxide (t-BHP) induced malondialdehyde (MDA) and dityrosine level in human vein endothelial cells (HUVEC)	[48]
TF3	Increased nitric oxide (NO) production and endothelial NO synthase (eNOS) activity in bovine aortic endothelial cells	[49]
	Inhibited plasminogen activator inhibitor type 1 (PAI-1) activity	[105]

synergistic effect in inducing G0/G1 cell cycle arrest in human lung adenocarcinoma cells [38]. Overall, various polyphenolic compounds in black tea exert multiple actions against cancer.

3. Effects of black tea on cardiovascular diseases

Cardiovascular disease (CAD) is becoming the leading cause of death in the world. Atherosclerosis is considered the major etiology in CAD. In vascular endothelium, low density lipoproteins are oxidized by free radicals forming ox-LDL, which activate and alter the adhesive property of monocytes, leading to their attachment to endothelium [39]. The attached monocytes are further differentiated into macrophages, which express scavenger receptors and can uptake ox-LDL. They finally become foam cells and result in the formation of fatty streaks and atheroma plaques [39]. The infiltration of monocytes/macrophages within vascular endothelium causes damage to endothelial cells that in turn results in endothelial dysfunction and reduction in vasodilation capacity in CAD [40].

Epidemiological studies have documented many benefits for cardiovascular health from drinking black tea. Black tea consumption lowered plasma uric acid and C-reactive protein (CRP) levels in both man and women [41] (Table 4) and it also reduced serum glucose, triglyceride, LDL/HDL plasma cholesterol ratio and increased plasma antioxidant that may contribute to reduce risk of CAD [42]. A clinical study reviewed that short-term (2 h) or long-term (4 weeks) of black tea consumption reversed endothelial dysfunction and improved flow-mediated dilation of brachial artery in patients with coronary arterial disease [43]. Black tea extract is also found to inhibit platelet aggregation in rabbit platelet polymorphonuclear leukocyte (PMN) system [44] that may prevent heart attack or stroke.

Theaflavins, TF1, TF2a, TF2b and TF3, all showed dose-dependent antioxidant activity in Cu²⁺-mediated LDL oxidation *in vitro* [45]. TF1 protected against ischemia/reperfusion injury by inhibiting mitochondrial permeability transition pore (mPTP) opening in isolated heart [46]. Dietary administration of TF1 significantly attenuated atherosclerotic lesion size and reduced

vascular superoxide, leukotriene B4 (LTB4), and plasma P-selectin concentrations in apolipoprotein E (ApoE)–/– gene-knockout mice [47]. TF3 also inhibited plasminogen activator inhibitor type 1 (PAI-1) activity that contributes to pathogenesis of vascular remodeling and is implicated in the development of myocardial infarction [47,48].

Oxidative stress is involved in endothelial dysfunction that has been considered a risk factor of cardiovascular disease [40]. Pre- and post-treatment of black tea extract both showed a protective effect against tert-butyl hydroperoxide (t-BHP) induced malondialdehyde (MDA) and dityrosine level in human vein endothelial cells (HUVEC) that exhibited beneficial effect on endothelial function [48]. Consumption of black tea increased antioxidant capacity in blood in patients with coronary arterial disease [43]. In bovine aortic endothelial cells, black tea polyphenols including theaflavins (TF3) and catechins (EGCG) significantly increased endothelial NO synthase (eNOS) activity in bovine aortic endothelial cells and vaso-relaxation activity in rat aortic rings [49]. These findings are consistent with the clinical study that black tea consumption reversed the endothelial dysfunction in humans [43] and further demonstrated the effectiveness of tea consumption in combating CVD.

4. Effects of black tea in chronic inflammatory diseases

Chronic inflammation is a key component in many human chronic diseases and conditions, including rheumatoid arthritis, inflammatory bowel disease (IBD), osteoporosis and allergy [50–53]. Activation and infiltration of inflammatory cells as well as excessive production of pro-inflammatory cytokines/chemokines are major characteristics and etiology in chronic inflammatory diseases. Inhibition of inflammatory cells recruitment and pro-inflammatory mediator production are considered the important steps targeted by natural dietary compounds in prevention or treatment of chronic inflammatory diseases [54].

Table 5
Effect of black tea and its bioactive compounds on chronic inflammatory diseases.

Targeted disease	Compound	Molecular mechanism	Ref.
–	Black tea	Reduced carrageenan and dextran-induced paw edema in SD rats	[55]
–	Black tea extract	Reduced lipopolysaccharide (LPS)-induced NO and O ₂ [–] production as well as inducible nitric oxide synthase (iNOS) expression in murine macrophage	[59]
–		Reduced ratings of delayed onset muscle soreness (DOMS), increased glutathione (GSH)/oxidized glutathione (GSSG) ratio and performance	[76]
–	TF1	Suppressed tumor necrosis factor (TNF)- α -induced interleukin (IL)-8 gene expression through blockage of NF- κ B activation and reduced DNA binding activity of activator protein (AP)-1	[62]
–		Inhibited LPS-induced IL-6, monocyte chemoattractant protein-1 (MCP-1), and intercellular adhesion molecule-1 (ICAM-1) expression by suppression of NF- κ B as well as MAPKs signaling	[63]
–	TF2	Suppressed TPA-induced inflammatory genes and COX-2 promoter activity in Caco-2 cells	[31]
–	TF3	Inhibited inflammatory enzyme expression and cytokines production in murine macrophage	[66]
–		Suppressed TPA-induced ear edema by inhibition of arachidonic acid metabolism	[67]
Inflammatory bowel disease	Black tea extract	Reduced DSS-induced colitis through inhibition of NF- κ B activation and increase of apoptosis	[56]
	TF1	Inhibited LPS-induced adhesion molecule expression by suppression of NF- κ B and JNK	[64]
	TF3	Against trinitrobenzene sulfonic acid (TNBS)-induced colitis and pro-inflammatory cytokines production in mice	[68]
Gastric ulcer	Black tea	Against indomethacin-induced gastric ulcer by down-regulation of myeloperoxidase (MPO) activity, iNOS and eNOS levels	[57]
Gastric inflammation	Black tea extract	Suppressed IL-1 β -induced IL-8 production and secretion by inhibition of NF- κ B activation	[60]
Allergy	TF2a	Decreased pro-inflammatory cytokines production in type IV allergy mice	[65]
Periodontal disease	TF3	Inhibited inflammatory cytokines production in human gingival fibroblasts	[69,70]
Osteoporosis	Black tea extract	Maintain skeletal health through reduction of active osteoclasts, inflammatory cytokines production and oxidative stress	[72]
		Against high fat diet (HFD) fed non-alcoholic steatohepatitis (NASH)-induced bone skeletal dysfunctional changes in Wistar rats	[74]
	TF3	Inhibited differentiation of osteoclasts through down-regulation of MMP-2 and MMP-9	[75]

There are many well-documented anti-inflammatory effects of feeding black tea or its extract in animal models. Black tea effectively reduced carrageenan/dextran-induced paw edema [55] and dextran sulfate sodium (DSS)-induced colitis [56] and improved healing against indomethacin-induced stomach ulceration [57] (Table 5). One key feature of these anti-inflammatory effects of black tea is *via* the inhibition of NF- κ B [56]. NF- κ B is a transcription factor that plays a central role in inflammation through its ability to induce transcription of pro-inflammatory genes. NF- κ B is highly activated at sites of inflammation in diverse diseases and can induce transcription of pro-inflammatory cytokines, chemokines, adhesion molecules, MMPs, COX-2, and inducible nitric oxide (iNOS) [58]. Inhibition of NF- κ B can effectively reduce the production of various pro-inflammatory mediators and enzymes including COX-2 and iNOS and switch off inflammation and therefore is considered a key target for anti-inflammatory therapeutics [58]. The anti-inflammatory effects of black tea are further confirmed by cell model that using black tea extract reduced nitric oxide (NO) and O₂[–] production [59] and inhibited the effects of pro-inflammatory cytokines [60].

Consistent with the results using black tea or its extract, major ingredients in black tea including TFs and EGCG inhibited the induction of NF- κ B [61] and thus were able to reduce inflammation mediators. By blocking NF- κ B, TF1 inhibited tumor necrosis factor (TNF)- α -mediated interleukin (IL)-8 gene [62] and the expression of LPS-induced interleukin-6 (IL-6), monocyte chemoattractant protein-1 (MCP-1), intercellular adhesion molecule-1 (ICAM-1) and vascular cell adhesion molecule (VCAM)-1 *via* the modulation of MAPKs signaling [63,64].

By modulation multiple transcription factors including NF- κ B, AP-1 and CREB, TF2 treatment suppressed TPA-induced inflammatory gene expression in Caco-2 cells, such as COX-2, TNF- α , ICAM-1, and iNOS. It also reduced COX-2 promoter activity [31]. Oral administration of TF2a reduced levels of pro-inflammatory cytokines, including IL-12, IFN- γ , and tumor TNF- α in type IV allergy in male ICR mice [65].

Various *in vitro* and *in vivo* models demonstrated the anti-inflammatory effects of TF3. Similar to TF1 and TF2, suppression of NF- κ B by TF3 inhibited LPS-induced iNOS expression in RAW264.7 macrophage [66]. In TPA-induced mouse ear edema model, TF3 showed stronger inhibitory effect

Table 6

Effect of black tea and its bioactive compounds on obesity and metabolic syndrome.

Targeted disease	Compound	Molecular mechanism	Ref.
Obesity, hyperglycemia	Black tea	Reduced HFD-induced body weight gain, improved hyperglycemia and glucose intolerance <i>via</i> increasing the level of glucose transporter type 4 (GLUT4) in male C57BL/6J mice	[79]
Obesity	Black tea extract	Suppressed HFD-induced body weight increase, adipose tissue mass and liver lipid content may through inhibition of intestinal lipid absorption in C57BL/6N mice	[81]
–	TF2a	Reduced lipid and triglyceride levels in fatty acid-overload HepG2 cells through induction of AMP-activated protein kinase (AMPK) signaling	[85]
–	TF1	Inhibited gene and protein expression of FAS in MCF-7 breast cancer cells	[26]
–	TF2a, TF2b	Reduced the solubility of cholesterol into micelles	[86,87]
Hypertriacylglycerolemia	TF3	Suppressed hypertriacylglycerolemia through inhibition pancreatic lipase activity in rats	[88]

than other tea polyphenols in reducing of IL-6, prostaglandin E2 (PGE2), and LTB4 levels [67]. Oral administration of TF3 decreased both gene and protein levels of TNF- α , IL-12, interferon (IFN)- γ and iNOS in trinitrobenzene sulfonic acid (TNBS)-induced colitis animal model [68]. TF3 suppressed IL-16 and CXC chemokine ligand 10 (CXCL10) production by targeting multiple signaling pathways in human gingival fibroblasts [69,70], which play a critical role in sustaining inflammatory response in the pathogenesis of periodontal disease [71].

Mononuclear cell infiltration and osteoclast activation have been linked to bone loss [52]. Black tea extract feeding effectively preserved and restored skeletal health by reducing osteoclast activation and oxidative stress caused by mononuclear cells in rats [72]. Osteoporosis is a common complication associated with chronic liver disease [73]. In high fat diet (HFD)-induced non-alcoholic steatohepatitis (NASH) model, consumption of black tea extract ameliorated bone skeletal dysfunctional changes, including bone turnover and density [74]. In addition, it was found that TF3 and EGCG inhibited the formation and differentiation of osteoclasts through decreasing gene expression of MMP-2 and MMP-9 as well as their enzymatic activities. These results supported the role of black tea polyphenols in preventing the degeneration of the matrix in bone and cartilage [75].

Delayed onset muscle soreness (DOMS) frequently occurs in athlete after high intensity exercise. Muscle soreness, which is initiated by muscle damage, is usually noted at 24 h post-exercise and can last as long as 5–7 d post-exercise. The muscle damage elicits inflammatory and oxidative responses that may exacerbate muscle injury and prolong the time to regeneration [76]. In a human trial study of healthy athletes engaging in high intensity anaerobic exercise, supplement with TF2-enriched black tea extract significantly reduced DOMS, increased glutathione (GSH)/oxidized glutathione (GSSG) ratio and performance [76]. These results are consistent with the molecular actions of black tea polyphenols that they activate transcription factor Nrf2, which up-regulates protective detoxifying and antioxidant enzymes including enzyme for glutathione biosynthesis and metabolism to maintain glutathione levels [19]. Activation of Nrf2 also attenuates the NF- κ B mediated inflammatory response [77], which may play a significant role in anti-inflammatory effects of black tea.

5. Effects of black tea in obesity and metabolic syndrome

Obesity is a highly prevalent condition and regarded as a risk factor for cardiovascular disease, hypertension and metabolic syndrome, fatty liver disease, type 2 diabetes and chronic kidney disease [78]. Obesity is currently considered a chronic inflammatory disorder that promotes the development of insulin resistance and diabetes [77]. In a high fat diet (HFD) induced obesity animal model, drinking black tea or its extract decreased body fat [79], and improved hyperglycemia and glucose intolerance [79] (Table 6). Consumption of black tea up-regulated glucose transporters type 4 (GLUT4) in muscle that resulted in increased glucose uptake and improved glucose intolerance [79]. It also suppressed C/EBP β expression in perirenal fat that implicated in adipocyte differentiation and fat accumulation [80]. Dietary feeding black tea polyphenol extract also suppressed HFD- or high sucrose-induced increased liver lipid content or dyslipidemia in animals [81]. The potential mechanisms include decreased hepatic fatty acid synthase (FAS) and increased AMP-activated protein kinase (AMPK) activity [82]. AMPK is the key cellular energy sensor that regulates glucose and lipid metabolisms and is the drug target of anti-diabetic drug such as metformin [83,84] and lipid-lowering drug, statins [84]. Activation of AMPK leads to increased glucose uptake *via* up-regulation of glucose transporters, turning off anabolic pathways including decreased fatty acid synthase for lipid synthesis and increased catabolic pathways in cells [83]. In agreement with the *in vivo* results, *in vitro* study showed that theaflavins and EGCG treatment strongly reduced intracellular total lipid, triglyceride, and cholesterol levels in fatty acid (FA)-overloaded liver cell line through activating AMPK signaling pathway [85], and TF1 was found to inhibit both gene and protein expression of FAS *in vitro* that may reduce cell lipogenesis [26].

Other possible mechanism of black tea against obesity and dyslipidemia also has been proposed. Both TF2a and TF2b are found to reduce micellar solubility of cholesterol *in vitro* [86,87]. Moreover, TF3 is more effective in inhibiting pancreatic lipase activity than that of EGCG by reducing triacylglycerol absorption and consequently suppressing postprandial hypertriacylglycerolemia [88]. These actions contributed to the prevention of black tea polyphenols against diet-induced obesity in animals [81].

Table 7
Effect of black tea and its bioactive compounds on neurological disorders.

Targeted disease or longevity	Compound	Molecular mechanism	Ref.
Alzheimer's disease	Black tea extract	Inhibited lipid membrane destabilization induced by amyloid β peptide ($A\beta$) 42 peptides <i>in vitro</i>	[101]
		Against $A\beta$ fibrils-induced toxicity in primary cultures of rat hippocampal cell	[102]
	TF2b, TF3	Potent inhibitor of PAI-1 in human plasma that may slow down progression of Alzheimer's disease	[105,106]
	TF3	Inhibited $A\beta$ fibril formation <i>via</i> reduction of ThT signals and assembly of $A\beta$ fibrils into nontoxic, spherical aggregates <i>in vitro</i>	[103]
Parkinson's disease	Black tea	Reduced risk of Parkinson's disease in the Singapore Chinese Health Study	[96]
	Black tea extract	Protected 6-hydroxydopamine (6-OHDA)-induced dopaminergic neurons damage, improved motor and neurochemical deficits in Wistar rats	[97]
		Attenuated 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine (MPTP/p) induced apoptosis and neurodegeneration in C57BL/6 mice through increase of dopamine transporter (DAT) and reduction of apoptosis in the substantia nigra	[98]
		Improved MPTP-induced damage of dopaminergic neurons by reduction of oxidative stress	[99]
		Attenuates MPTP-induced neuroinflammation and apoptosis by down-regulation of inflammatory cytokines	[100]

In addition to the effects of reducing postprandial lipids absorption, drinking black tea also reduced postprandial blood glucose levels and enhanced insulin response in healthy human subjects [89]. Solutions of black tea increased insulin activity in rat adipocytes [90]. One of the potential mechanisms includes theaflavins mimicked the effects of insulin and insulin-like growth factor-1 (IGF-1) by initiating phosphorylation of Forkhead transcription factor O1a (FOXO1a) and repressing phosphoenolpyruvate carboxykinase (PEPCK) promoter activity. FOXO1a and PEPCK are both important components of the insulin-signaling pathway [91]. The phosphorylation of FOXO1a may be mediated by AMPK [92,93], which is the key metabolic regulatory enzyme activated by black tea polyphenols.

6. Potential use of black tea in neurodegenerative disorders

Parkinson's disease (PD) and Alzheimer's disease (AD) are age-dependent neurodegenerative disorders [94]. Parkinson's disease is characterized by accumulation of α -synuclein protein of Lewy bodies and loss of dopaminergic neurons [95]. Consumption of black tea was associated with the reduced risk of Parkinson's disease in an epidemiology study. It was concluded that ingredients of black tea other than caffeine were responsible for the beverage's inverse association with Parkinson's disease [96]. In 6-hydroxydopamine (6-OHDA)-lesioned rat model of PD, oral administration of black tea extract before or after 6-OHDA treatment reduced dopaminergic neuron damage and improved motor and neurochemical deficits [97]. In another neurotoxin 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine (MPTP)-induced PD animal model, oral treatment of TF1 was effective in reduction of oxidative stress-induced neurodegeneration and apoptosis by suppressing Bax down-regulation and increasing Bcl-2. It also effectively reduced inflammatory mediators (IL-1 β , IL-6, TNF- α , IL-10)

and ameliorated dopamine transporters' reduction and behavioral deficits [98–100].

Black tea may also be beneficial in the treatment of AD. In AD, it is believed that the neurodegeneration is caused by the cleavage product of amyloid precursor protein (APP) and the aggregation of amyloid β peptide ($A\beta$) in brain. Black tea extract inhibited $A\beta$ 42 peptides-induced lipid membrane destabilization that may protect membrane damage [101] (Table 7). In primary culture of rat hippocampal cell, black tea extract displayed neuroprotective effect by the inhibition of $A\beta$ -induced cytotoxicity [102]. Black tea polyphenols including theaflavins and EGCG inhibited formation of toxic amyloid- β ($A\beta$) and α -synuclein (α S) fibrils *in vitro* by stimulating the assembly of $A\beta$ and α S into nontoxic spherical aggregates that are independent of their antioxidant effects [103]. TF3 was less susceptible to oxidation and had an increased efficacy under oxidizing conditions than EGCG [103]. EGCG was effective in reducing the β -amyloid mediated cognitive impairment in Alzheimer transgenic mice [104]. It would be interesting to find out if TF3 is more potent in same animal model. In addition to the direct effects on $A\beta$ aggregation, both TF2a and TF3 were found to inhibit PAI-1 activity, which has been implicated in the accumulation of $A\beta$ plaque in brain [105,106]. Results from all these *in vitro* studies look promising and further *in vivo* and clinical studies are needed to confirm the efficacy of black tea polyphenols in the treatment of PD and AD.

7. Summary and perspective

In summary, black tea and its extracts possess many health promoting properties supported by a myriad of data from *in vitro* and *in vivo* experiments as well as human clinical trials. The major black tea polyphenols, theaflavins and catechins should be recognized as biomolecular markers of black tea and its extract. Delineating biological mechanisms that are associated with a

well-characterized chemical profile will provide crucial information for product development in targeting specific conditions.

The key molecular action of black tea polyphenols include the activation of Nrf2 transcription factor that up-regulates detoxifying and antioxidant enzymes. This action may also contribute to the inhibition of NF- κ B, the master transcription factor that regulates inflammation. Black tea polyphenols also activate AMPK the key cellular energy sensor that regulates glucose and lipid metabolisms. All these diverse actions provide the scientific foundation in supporting the consumption of black tea or its extract for the prevention of cancer and treatment of many inflammatory and metabolic diseases.

References

- [1] S. Li, C.Y. Lo, M.H. Pan, et al., Black tea: chemical analysis and stability, *Food & Function* 4 (2013) 10–18.
- [2] D. Hanahan, R.A. Weinberg, The hallmarks of cancer, *Cell* 100 (2000) 57–70.
- [3] M.H. Pan, C.T. Ho, Chemopreventive effects of natural dietary compounds on cancer development, *Chemical Society Reviews* 37 (2008) 2558–2574.
- [4] J.A. Baker, K. Boakye, S.E. McCann, et al., Consumption of black tea or coffee and risk of ovarian cancer, *International Journal of Gynecological Cancer* 17 (2007) 50–54.
- [5] I. Dora, L. Arab, A. Martinchik, et al., Black tea consumption and risk of rectal cancer in Moscow population, *Annals of Epidemiology* 13 (2003) 405–411.
- [6] N. Tang, Y. Wu, B. Zhou, et al., Green tea, black tea consumption and risk of lung cancer: a meta-analysis, *Lung Cancer* 65 (2009) 274–283.
- [7] R.A. Goldbohm, M.G. Hertog, H.A. Brants, et al., Consumption of black tea and cancer risk: a prospective cohort study, *Journal of the National Cancer Institute* 88 (1996) 93–100.
- [8] M. Kapiszewska, M. Miskiewicz, P.T. Ellison, et al., High tea consumption diminishes salivary 17 β -estradiol concentration in Polish women, *British Journal of Nutrition* 95 (2006) 989–995.
- [9] C.M. Nagle, C.M. Olsen, C.J. Bain, et al., Tea consumption and risk of ovarian cancer, *Cancer Causes and Control* 21 (2010) 1485–1491.
- [10] A.H. Lee, D. Su, M. Pasalich, et al., Tea consumption reduces ovarian cancer risk, *Cancer Epidemiology* 37 (2013) 54–59.
- [11] S. Wu, F. Li, X. Huang, et al., The association of tea consumption with bladder cancer risk: a meta-analysis, *Asia Pacific Journal of Clinical Nutrition* 22 (2013) 128–137.
- [12] P. Saha, S. Das, Elimination of deleterious effects of free radicals in murine skin carcinogenesis by black tea infusion, theaflavins & epigallocatechin gallate, *Asian Pacific Journal of Cancer Prevention* 3 (2002) 225–230.
- [13] P. Roy, N. Nigam, J. George, et al., Induction of apoptosis by tea polyphenols mediated through mitochondrial cell death pathway in mouse skin tumors, *Cancer Biology and Therapy* 8 (2009) 1281–1287.
- [14] J. George, M. Singh, A.K. Srivastava, et al., Resveratrol and black tea polyphenol combination synergistically suppress mouse skin tumors growth by inhibition of activated MAPKs and p53, *PLoS ONE* 6 (2011) e23395.
- [15] P. Roy, J. George, S. Srivastava, et al., Inhibitory effects of tea polyphenols by targeting cyclooxygenase-2 through regulation of nuclear factor kappa B, Akt and p53 in rat mammary tumors, *Investigational New Drugs* 29 (2011) 225–231.
- [16] K.V. Chandra Mohan, Y. Hara, S.K. Abraham, et al., Comparative evaluation of the chemopreventive efficacy of green and black tea polyphenols in the hamster buccal pouch carcinogenesis model, *Clinical Biochemistry* 38 (2005) 879–886.
- [17] R. Patel, A. Ingle, G.B. Maru, Polymeric black tea polyphenols inhibit 1,2-dimethylhydrazine induced colorectal carcinogenesis by inhibiting cell proliferation via Wnt/beta-catenin pathway, *Toxicology and Applied Pharmacology* 227 (2008) 136–146.
- [18] D. Sinha, M. Roy, Antagonistic role of tea against sodium arsenite-induced oxidative DNA damage and inhibition of DNA repair in Swiss albino mice, *Journal of Environmental Pathology, Toxicology and Oncology* 30 (2011) 311–322.
- [19] C. Chen, A.N. Kong, Dietary chemopreventive compounds and ARE/EpRE signaling, *Free Radical Biology and Medicine* 36 (2004) 1505–1516.
- [20] Y.J. Surh, J.K. Kundu, H.K. Na, Nrf2 as a master redox switch in turning on the cellular signaling involved in the induction of cytoprotective genes by some chemopreventive phytochemicals, *Planta Medica* 74 (2008) 1526–1539.
- [21] B. Halder, G.S. Das, A. Gomes, Black tea polyphenols induce human leukemic cell cycle arrest by inhibiting Akt signaling: possible involvement of Hsp90, Wnt/beta-catenin signaling and FOXO1, *FEBS Journal* 279 (2012) 2876–2891.
- [22] M. Singh, R. Singh, K. Bhui, et al., Tea polyphenols induce apoptosis through mitochondrial pathway and by inhibiting nuclear factor-kappaB and Akt activation in human cervical cancer cells, *Oncology Research* 19 (2011) 245–257.
- [23] U. Bhattacharya, B. Halder, S. Mukhopadhyay, et al., Role of oxidation-triggered activation of JNK and p38 MAPK in black tea polyphenols induced apoptotic death of A375 cells, *Cancer Science* 100 (2009) 1971–1978.
- [24] H. Sil, T. Sen, S. Moulik, et al., Black tea polyphenol (theaflavin) downregulates MMP-2 in human melanoma cell line A375 by involving multiple regulatory molecules, *Journal of Environmental Pathology, Toxicology and Oncology* 29 (2010) 55–68.
- [25] L. Lahiry, B. Saha, J. Chakraborty, et al., Theaflavins target Fas/caspase-8 and Akt/pBad pathways to induce apoptosis in p53-mutated human breast cancer cells, *Carcinogenesis* 31 (2010) 259–268.
- [26] C.W. Yeh, W.J. Chen, C.T. Chiang, et al., Suppression of fatty acid synthase in MCF-7 breast cancer cells by tea and tea polyphenols: a possible mechanism for their hypolipidemic effects, *Pharmacogenomics Journal* 3 (2003) 267–276.
- [27] S.S. Banerjee, S.S. Manna, S. Mukherjee, et al., Black tea polyphenols restrict benzopyrene-induced mouse lung cancer progression through inhibition of Cox-2 and induction of caspase-3 expression, *Asian Pacific Journal of Cancer Prevention* 7 (2006) 661–666.
- [28] S. Banerjee, S. Manna, P. Saha, et al., Black tea polyphenols suppress cell proliferation and induce apoptosis during benzo(a)pyrene-induced lung carcinogenesis, *European Journal of Cancer Prevention* 14 (2005) 215–221.
- [29] G.Y. Yang, Z. Liu, D.N. Seril, et al., Black tea constituents, theaflavins, inhibit 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK)-induced lung tumorigenesis in A/J mice, *Carcinogenesis* 18 (1997) 2361–2365.
- [30] T. Mujtaba, Q.P. Dou, Black tea polyphenols inhibit tumor proteasome activity, *In Vivo* 26 (2012) 197–202.
- [31] A. Gossau, D.L. En Jao, M.T. Huang, et al., Effects of the black tea polyphenol theaflavin-2 on apoptotic and inflammatory pathways *in vitro* and *in vivo*, *Molecular Nutrition & Food Research* 55 (2011) 198–208.
- [32] J. Lu, C.T. Ho, G. Ghai, et al., Differential effects of theaflavin monogallates on cell growth, apoptosis, and Cox-2 gene expression in cancerous versus normal cells, *Cancer Research* 60 (2000) 6465–6471.
- [33] M.H. Pan, Y.C. Liang, S.Y. Lin-Shiau, et al., Induction of apoptosis by the oolong tea polyphenol theasinensin A through cytochrome c release and activation of caspase-9 and caspase-3 in human U937 cells, *Journal of Agricultural and Food Chemistry* 48 (2000) 6337–6346.
- [34] A. Rajavelu, Z. Tulyasheva, R. Jaiswal, et al., The inhibition of the mammalian DNA methyltransferase 3a (Dnmt3a) by dietary black tea and coffee polyphenols, *BMC Biochemistry* 12 (2011) 16.
- [35] Y.C. Liang, Y.C. Chen, Y.L. Lin, et al., Suppression of extracellular signals and cell proliferation by the black tea polyphenol, theaflavin-3,3'-digallate, *Carcinogenesis* 20 (1999) 733–736.
- [36] H. Mizuno, Y.Y. Cho, F. Zhu, et al., Theaflavin-3,3'-digallate induces epidermal growth factor receptor downregulation, *Molecular Carcinogenesis* 45 (2006) 204–212.

- [37] A.G. Schuck, M.B. Ausubel, H.L. Zuckerbraun, et al., Theaflavin-3,3'-digallate, a component of black tea: an inducer of oxidative stress and apoptosis, *Toxicology In Vitro* 22 (2008) 598–609.
- [38] W. Li, J.X. Wu, Y.Y. Tu, Synergistic effects of tea polyphenols and ascorbic acid on human lung adenocarcinoma SPC-A-1 cells, *Journal of Zhejiang University Science B* 11 (2010) 458–464.
- [39] J. Mestas, K. Ley, Monocyte-endothelial cell interactions in the development of atherosclerosis, *Trends in Cardiovascular Medicine* 18 (2008) 228–232.
- [40] Y. Higashi, K. Noma, M. Yoshizumi, et al., Endothelial function and oxidative stress in cardiovascular diseases, *Circulation Journal* 73 (2009) 411–418.
- [41] T. Baborun, A. Luximon-Ramma, T.K. Gunness, et al., Black tea reduces uric acid and C-reactive protein levels in humans susceptible to cardiovascular diseases, *Toxicology* 278 (2010) 68–74.
- [42] T. Baborun, A. Luximon-Ramma, V.S. Neergheen-Bhujun, et al., The effect of black tea on risk factors of cardiovascular disease in a normal population, *Preventive Medicine* 54 (Suppl. 1) (2012) 98–102.
- [43] S.J. Duffy, J.F. Keaney, M. Holbrook, et al., Short- and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease, *Circulation* 104 (2001) 151–156.
- [44] J. Sugatani, N. Fukazawa, K. Ujihara, et al., Tea polyphenols inhibit acetyl-CoA:1-alkyl-sn-glycero-3-phosphocholine acetyltransferase (a key enzyme in platelet-activating factor biosynthesis) and platelet-activating factor-induced platelet aggregation, *International Archives of Allergy and Immunology* 134 (2004) 17–28.
- [45] L.K. Leung, Y. Su, R. Chen, et al., Theaflavins in black tea and catechins in green tea are equally effective antioxidants, *Journal of Nutrition* 131 (2001) 2248–2251.
- [46] H. Ma, X. Huang, Q. Li, et al., ATP-dependent potassium channels and mitochondrial permeability transition pores play roles in the cardioprotection of theaflavin in young rat, *Journal of Physiological Sciences* 61 (2011) 337–342.
- [47] W.M. Loke, J.M. Proudfoot, J.M. Hodgson, et al., Specific dietary polyphenols attenuate atherosclerosis in apolipoprotein E-knockout mice by alleviating inflammation and endothelial dysfunction, *Arteriosclerosis, Thrombosis, and Vascular Biology* 30 (2010) 749–757.
- [48] W. Luczaj, E. Zapor, M. Szczepanski, et al., Polyphenols action against oxidative stress formation in endothelial cells, *Acta Poloniae Pharmaceutica* 66 (2009) 617–624.
- [49] M. Lorenz, J. Urban, U. Engelhardt, et al., Green and black tea are equally potent stimuli of NO production and vasodilation: new insights into tea ingredients involved, *Basic Research in Cardiology* 104 (2009) 100–110.
- [50] H.L. Wright, R.J. Moots, R.C. Bucknall, et al., Neutrophil function in inflammation and inflammatory diseases, *Rheumatology* 49 (2010) 1618–1631.
- [51] M. Averbek, C. Gebhardt, F. Emmrich, et al., Immunologic principles of allergic disease, *Journal Der Deutschen Dermatologischen Gesellschaft* 5 (2007) 1015–1028.
- [52] G.R. Mundy, Osteoporosis and inflammation, *Nutrition Reviews* 65 (2007) S147–S151.
- [53] R. Scrivo, F.M. Di, A. Spadaro, et al., The immunology of rheumatoid arthritis, *Annals of the New York Academy of Sciences* 1108 (2007) 312–322.
- [54] M.H. Pan, C.S. Lai, C.T. Ho, Anti-inflammatory activity of natural dietary flavonoids, *Food & Function* 1 (2010) 15–31.
- [55] C. Chattopadhyay, N. Chakrabarti, M. Chatterjee, et al., Black tea (*Camellia sinensis*) decoction shows immunomodulatory properties on an experimental animal model and in human peripheral mononuclear cells, *Pharmacognosy Research* 4 (2012) 15–21.
- [56] Y.A. Song, Y.L. Park, K.Y. Kim, et al., Black tea extract prevents lipopolysaccharide-induced NF-kappaB signaling and attenuates dextran sulfate sodium-induced experimental colitis, *BMC: Complementary and Alternative Medicine* 11 (2011) 91.
- [57] B. Adhikary, S.K. Yadav, S. Chand, et al., Black tea and theaflavins suppress various inflammatory modulators and i-NOS mediated nitric oxide synthesis during gastric ulcer healing, *Free Radical Research* 45 (2011) 767–778.
- [58] P.P. Tak, G.S. Firestein, NF-kappaB: a key role in inflammatory diseases, *Journal of Clinical Investigation* 107 (2001) 7–11.
- [59] A. Sarkar, A. Bhaduri, Black tea is a powerful chemopreventor of reactive oxygen and nitrogen species: comparison with its individual catechin constituents and green tea, *Biochemical and Biophysical Research Communications* 284 (2001) 173–178.
- [60] F. Gutierrez-Orozco, B.R. Stephens, A.P. Neilson, et al., Green and black tea inhibit cytokine-induced IL-8 production and secretion in AGS gastric cancer cells via inhibition of NF-kappaB activity, *Planta Medica* 76 (2010) 1659–1665.
- [61] M. Nomura, W. Ma, N. Chen, et al., Inhibition of 12-O-tetradecanoylphorbol-13-acetate-induced NF-kappaB activation by tea polyphenols, (–)-epigallocatechin gallate and theaflavins, *Carcinogenesis* 21 (2000) 1885–1890.
- [62] R. Aneja, K. Odoms, A.G. Denenberg, et al., Theaflavin, a black tea extract, is a novel anti-inflammatory compound, *Critical Care Medicine* 32 (2004) 2097–2103.
- [63] S. Kim, Y.E. Joo, Theaflavin inhibits LPS-induced IL-6, MCP-1, and ICAM-1 expression in bone marrow-derived macrophages through the blockade of NF-kappaB and MAPK signaling pathways, *Chonnam Medical Journal* 47 (2011) 104–110.
- [64] Y.A. Song, Y.L. Park, S.H. Yoon, Black tea polyphenol theaflavin suppresses LPS-induced ICAM-1 and VCAM-1 expression via blockage of NF-kappaB and JNK activation in intestinal epithelial cells, *Inflammation Research* 60 (2011) 493–500.
- [65] K. Yoshino, K. Yamazaki, M. Sano, Preventive effects of black tea theaflavins against mouse type IV allergy, *Journal of the Science of Food and Agriculture* 90 (2010) 1983–1987.
- [66] M.H. Pan, S.Y. Lin-Shiau, C.T. Ho, et al., Suppression of lipopolysaccharide-induced nuclear factor-kappaB activity by theaflavin-3,3'-digallate from black tea and other polyphenols through down-regulation of IkappaB kinase activity in macrophages, *Biochemical Pharmacology* 59 (2000) 357–367.
- [67] M.T. Huang, Y. Liu, D. Ramji, et al., Inhibitory effects of black tea theaflavin derivatives on 12-O-tetradecanoylphorbol-13-acetate-induced inflammation and arachidonic acid metabolism in mouse ears, *Molecular Nutrition & Food Research* 50 (2006) 115–122.
- [68] A. Ukil, S. Maity, P.K. Das, Protection from experimental colitis by theaflavin-3,3'-digallate correlates with inhibition of IKK and NF-kappaB activation, *British Journal of Pharmacology* 149 (2006) 121–131.
- [69] Y. Hosokawa, I. Hosokawa, K. Ozaki, et al., Tea polyphenols inhibit IL-6 production in tumor necrosis factor superfamily 14-stimulated human gingival fibroblasts, *Molecular Nutrition & Food Research* 54 (Suppl. 2) (2010) 151–158.
- [70] Y. Hosokawa, I. Hosokawa, S. Shindo, et al., Black tea polyphenol inhibits CXCL10 production in oncostatin M-stimulated human gingival fibroblasts, *International Immunopharmacology* 11 (2011) 670–674.
- [71] T. Ara, K. Kurata, K. Hirai, et al., Human gingival fibroblasts are critical in sustaining inflammation in periodontal disease, *Journal of Periodontal Research* 44 (2009) 21–27.
- [72] A.S. Das, M. Mukherjee, D. Das, et al., Protective action of aqueous black tea (*Camellia sinensis*) extract (BTE) against ovariectomy-induced oxidative stress of mononuclear cells and its associated progression of bone loss, *Phytotherapy Research* 23 (2009) 1287–1294.
- [73] O.M. Crosbie, R. Freaney, M.J. McKenna, Bone density, vitamin D status, and disordered bone remodeling in end-stage chronic liver disease, *Calcified Tissue International* 64 (1999) 295–300.
- [74] S. Karmakar, S. Majumdar, A. Maiti, et al., Protective role of black tea extract against nonalcoholic steatohepatitis-induced skeletal dysfunction, *Journal of Osteoporosis* (2011) 426863.
- [75] Y. Oka, S. Iwai, H. Amano, et al., Tea polyphenols inhibit rat osteoclast formation and differentiation, *Journal of Pharmacological Sciences* 118 (2012) 55–64.
- [76] S.M. Arent, M. Senso, D.L. Golem, et al., The effects of theaflavin-enriched black tea extract on muscle soreness, oxidative stress, inflammation, and endocrine responses to acute anaerobic interval training: a randomized, double-blind, crossover study, *Journal of the International Society of Sports Nutrition* 7 (2010) 11.

- [77] W. Li, T.O. Khor, C. Xu, et al., Activation of Nrf2-antioxidant signaling attenuates NF-kappaB-inflammatory response and elicits apoptosis, *Biochemical Pharmacology* 76 (2008) 1485–1489.
- [78] G. Reaven, F. Abbasi, T. McLaughlin, Obesity, insulin resistance, and cardiovascular disease, *Recent Progress in Hormone Research* 59 (2004) 207–223.
- [79] S. Nishiumi, H. Bessyo, M. Kubo, Green and black tea suppress hyperglycemia and insulin resistance by retaining the expression of glucose transporter 4 in muscle of high-fat diet-fed C57BL/6J mice, *Journal of Agricultural and Food Chemistry* 58 (2010) 12916–12923.
- [80] N. Chen, R. Bezzina, E. Hinch, et al., Green tea, black tea, and epigallocatechin modify body composition, improve glucose tolerance, and differentially alter metabolic gene expression in rats fed a high-fat diet, *Nutrition Research* 29 (2009) 784–793.
- [81] S. Uchiyama, Y. Taniguchi, A. Saka, et al., Prevention of diet-induced obesity by dietary black tea polyphenols extract *in vitro* and *in vivo*, *Nutrition* 27 (2011) 287–292.
- [82] H.C. Huang, J.K. Lin, Pu-erh tea, green tea, and black tea suppresses hyperlipidemia, hyperleptinemia and fatty acid synthase through activating AMPK in rats fed a high-fructose diet, *Food & Function* 3 (2012) 170–177.
- [83] D.G. Hardie, Sensing of energy and nutrients by AMP-activated protein kinase, *American Journal of Clinical Nutrition* 93 (2011) 891S–896S.
- [84] A.K. Wong, J. Howie, J.R. Petrie, et al., AMP-activated protein kinase pathway: a potential therapeutic target in cardiometabolic disease, *Clinical Science (London)* 116 (2009) 607–620.
- [85] C.L. Lin, H.C. Huang, J.K. Lin, Theaflavins attenuate hepatic lipid accumulation through activating AMPK in human HepG2 cells, *Journal of Lipid Research* 48 (2007) 2334–2343.
- [86] M.A. Vermeer, T.P. Mulder, H.O. Molhuizen, Theaflavins from black tea, especially theaflavin-3-gallate, reduce the incorporation of cholesterol into mixed micelles, *Journal of Agricultural and Food Chemistry* 56 (2008) 12031–12036.
- [87] I. Ikeda, T. Yamahira, M. Kato, et al., Black-tea polyphenols decrease micellar solubility of cholesterol *in vitro* and intestinal absorption of cholesterol in rats, *Journal of Agricultural and Food Chemistry* 58 (2010) 8591–8595.
- [88] M. Kobayashi, M. Ichitani, Y. Suzuki, et al., Black-tea polyphenols suppress postprandial hypertriacylglycerolemia by suppressing lymphatic transport of dietary fat in rats, *Journal of Agricultural and Food Chemistry* 57 (2009) 7131–7136.
- [89] J.A. Bryans, P.A. Judd, P.R. Ellis, The effect of consuming instant black tea on postprandial plasma glucose and insulin concentrations in healthy humans, *Journal of the American College of Nutrition* 26 (2007) 471–477.
- [90] R.A. Anderson, M.M. Polansky, Tea enhances insulin activity, *Journal of Agricultural and Food Chemistry* 50 (2002) 7182–7186.
- [91] A.R. Cameron, S. Anton, L. Melville, et al., Black tea polyphenols mimic insulin/insulin-like growth factor-1 signalling to the longevity factor FOXO1a, *Aging Cell* 7 (2008) 69–77.
- [92] A. Barthel, D. Schmoll, K.D. Kruger, et al., Regulation of the forkhead transcription factor FKHR (FOXO1a) by glucose starvation and AICAR, an activator of AMP-activated protein kinase, *Endocrinology* 143 (2002) 3183–3186.
- [93] M. Dixit, E. Bess, B. Fisslthaler, et al., Shear stress-induced activation of the AMP-activated protein kinase regulates FoxO1a and angiotensin-2 in endothelial cells, *Cardiovascular Research* 77 (2008) 160–168.
- [94] X.G. Luo, J.Q. Ding, S.D. Chen, Microglia in the aging brain: relevance to neurodegeneration, *Molecular Neurodegeneration* 5 (2010) 12.
- [95] M.G. Tansey, M.S. Goldberg, Neuroinflammation in Parkinson's disease: its role in neuronal death and implications for therapeutic intervention, *Neurobiology of Disease* 37 (2010) 510–518.
- [96] L.C. Tan, W.P. Koh, J.M. Yuan, et al., Differential effects of black versus green tea on risk of Parkinson's disease in the Singapore Chinese Health Study, *American Journal of Epidemiology* 167 (2008) 553–560.
- [97] R.K. Chaturvedi, S. Shukla, K. Seth, et al., Neuroprotective and neurorescue effect of black tea extract in 6-hydroxydopamine-lesioned rat model of Parkinson's disease, *Neurobiology of Disease* 22 (2006) 421–434.
- [98] A. Anandhan, K. Tamilselvam, T. Radhiga, et al., Theaflavin, a black tea polyphenol, protects nigral dopaminergic neurons against chronic MPTP/probenecid induced Parkinson's disease, *Brain Research* 1433 (2012) 104–113.
- [99] A. Anandhan, U. Janakiraman, T. Manivasagam, Theaflavin ameliorates behavioral deficits, biochemical indices and monoamine transporters expression against subacute 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced mouse model of Parkinson's disease, *Neuroscience* 218 (2012) 257–267.
- [100] A. Anandhan, M.M. Essa, T. Manivasagam, Therapeutic attenuation of neuroinflammation and apoptosis by black tea theaflavin in chronic MPTP/probenecid model of Parkinson's disease, *Neurotoxicity Research* 23 (2013) 166–173.
- [101] A.J. Gauci, M. Caruana, A. Giese, et al., Identification of polyphenolic compounds and black tea extract as potent inhibitors of lipid membrane destabilization by Abeta aggregates, *Journal of Alzheimer's Disease* 27 (2011) 767–779.
- [102] S. Bastianetto, Z.X. Yao, V. Papadopoulos, et al., Neuroprotective effects of green and black teas and their catechin gallate esters against beta-amyloid-induced toxicity, *European Journal of Neuroscience* 23 (2006) 55–64.
- [103] G. Grelle, A. Otto, M. Lorenz, et al., Black tea theaflavins inhibit formation of toxic amyloid-beta and alpha-synuclein fibrils, *Biochemistry* 50 (2011) 10624–10636.
- [104] K. Rezai-Zadeh, G.W. Arendash, H. Hou, et al., Green tea epigallocatechin-3-gallate (EGCG) reduces beta-amyloid mediated cognitive impairment and modulates tau pathology in Alzheimer transgenic mice, *Brain Research* 1214 (2008) 177–187.
- [105] J. Jankun, M. Skotnicka, W. Lysiak-Szydlowska, et al., Diverse inhibition of plasminogen activator inhibitor type 1 by theaflavins of black tea, *International Journal of Molecular Medicine* 27 (2011) 525–529.
- [106] E. Skrzypczak-Jankun, J. Jankun, Theaflavin digallate inactivates plasminogen activator inhibitor: could tea help in Alzheimer's disease and obesity? *International Journal of Molecular Medicine* 26 (2010) 45–50.