

Chemistry and health beneficial effects of oolong tea and theasinensins

Monthana Weerawatanakorn^a, Wei-Lun Hung^b, Min-Hsiung Pan^c, Shiming Li^b, Daxiang Li^d,
Xiaochun Wan^d, Chi-Tang Ho^{b,*}

^a Department of Agro-Industry, Faculty of Agriculture, Natural Resources and Environment, Naresuan University, Phitsanulok 65000, Thailand

^b Department of Food Science, Rutgers University, 65 Dudley Road, New Brunswick, NJ 08901, USA

^c Institute of Food Science and Technology, National Taiwan University, Taipei 10617, Taiwan, China

^d State Key Laboratory of Tea Plant Biology and Utilization, Anhui Agricultural University, 130 West Changjiang Rd., Hefei 230036, Anhui, China

Received 28 September 2015; received in revised form 15 October 2015; accepted 25 October 2015

Abstract

Among six major types of tea (white, green, oolong, yellow, black, and dark teas) from *Camellia sinensis*, oolong tea, a semi-fermented tea, with its own unique aroma and taste, has become a popular consumption as indicated by the increasing production. Representing the characteristic flavonoids of oolong tea, theasinensins are dimeric flavan-3-ols. Many recent studies have indicated that oolong tea and theasinensins possess several health benefit properties. We consider it significant and necessary to have a comprehensive review in the recent advances of oolong tea. Therefore, the aim of the present review is to provide a new perspective on oolong tea and its characteristic phytochemicals, theasinensins associated with health benefits, molecular action pathway, and chemical mechanism of theasinensin formation from scientific evidences available on the literature. Furthermore, the chemical characterization of the oxidation products and the model oxidation system to the chemical changes of theasinensins are also discussed.

© 2015 Beijing Academy of Food Sciences. Production and hosting by Elsevier B.V. All rights reserved.

1. Introduction

Generally, functional foods may be classified into three categories: food to which a component has been added, food in which a component has been modified in nature and/or bioavailability, and conventional food containing naturally occurring bioactive substance [1]. For the later class, tea from *Camellia sinensis* is one of the most important functional foods and has held the second most popular beverage in consumption among all beverages except water worldwide. Tea contains over 4,000 chemicals and some of which have health promoting properties [2].

Based on different degree of fermentation during tea process, tea can be simplistically divided into three major types: green (unfermented) tea produced from fresh tea leaves and enzymatic oxidation is inhibited using steaming or pan-frying; oolong (partially-fermented) tea made by wilting fresh leaves by sun, then slightly bruising; and black (fully fermented) tea made by crushing tea leaves to release the polyphenol oxidase and peroxidase for fully catalyzing the enzymatic oxidation and polymerization of original tea catechins [3–6]. Many potential health promotion properties associated with these three types of tea consumption have been reported [7,8]. Comparing with

Abbreviations: AGH, alpha-glucosidase; Akt, protein kinase B (PKB); ALPHA-TOH, alpha-tocopherol; AMPK, 5' adenosine monophosphate-activated protein kinase; CaMKK, Ca(2+)/calmodulin-dependent protein kinase; COX-2, cyclooxygenase-2; EC, (–)-epicatechin; ECG, (–)-epicatechin-3-gallate; EGC, (–)-epigallocatechin; EGCG, (–)-epigallocatechin 3-O-gallate; EGFR, epidermal growth factor receptor; ERK, extracellular signal-regulated protein kinases; HSV, herpes simplex virus; iNOS, inducible nitric oxide synthase; IL-12 (p70), interleukin-12; MAPK, mitogen-activated protein kinases; MCP-1, monocyte chemoattractant protein-1; MIC, minimum inhibitory concentration; MMP-2, matrix metalloproteinase-2; MNIC, maximum non-inhibitory concentration; MRSA, methicillin-resistant *Staphylococcus aureus*; 2-NBDG, 2-[N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)amino]-2-deoxyglucose; NF-κB, nuclear factor kappa-light-chain-enhancer of activated B cells; NO, nitric oxide; P13K, phosphoinositide 3-kinase; PGE2, prostaglandin E2; TR, thearubigin; TF, theaflavin; TAK1, transforming growth factor beta-activated kinase 1; TNF-alpha, tumor necrosis factor alpha; Sp1, transcription factor (specificity protein 1); SGLT1, sodium-dependent glucose cotransporters (sodium-glucose linked transporter); SOD, superoxide dismutase.

* Corresponding author at: Department of Food Science, Rutgers University, 65 Dudley Road, New Brunswick, NJ 08901, USA. Tel.: +1 848 932 5553.

E-mail address: ho@aesop.rutgers.edu (C.-T. Ho).

Peer review under responsibility of Beijing Academy of Food Sciences.

Table 1
Different fermented tea and catechin contents.

| Types of tea | Fermentation | Total catechins (% w/w) |
|-----------------|--|-------------------------|
| Pu-erh tea | Microbial fermentation | 6.07 ± 0.18 |
| Iron Buddha tea | Semi-fermentation | 7.49 ± 0.22 |
| Oolong tea | Semi-fermentation | 8.05 ± 0.18 |
| Jasmine tea | Less fermentation (reprocessed from loose green tea scented with fresh Jasmine flower) | 12.72 ± 0.70 |
| Lung Chen tea | Less fermentation | 14.57 ± 1.08 |

Adapted and modified from Sajilata et al. [5].

green tea having only monomeric catechins, *i.e.* (–)-epicatechin (EC), (–)-epicatechin-3-gallate (ECG), (–)-epigallocatechin (EGC), and (–)-epigallocatechin-3-gallate (EGCG), fully fermented black tea and semi-fermented oolong tea contain a mixture of catechins and their oxidized polymeric substances such as theaflavins and thearubigins [9]. Comparison of catechin contents in different fermented and semi-fermented teas is illustrated in Table 1. Theaflavins (TFs) and thearubigins (TRs), main secondary polyphenols formed during fermentation process by enzymatic oxidation, have been extensively studied on bioactivities and formation mechanism. The orange red or brown color and astringent taste of black tea infusion is attributed to TFs as TRs contribute to rusty color and richness taste [4,10]. There are four major TFs in black tea and oolong tea, that is, theaflavin (TF1), theaflavin-3-gallate (TF2a), theaflavin-3'-gallate (TF2b), and theaflavin-3,3'-digallate (TF3) [6,11,12]. Chemical structures of various types of tea catechins and theaflavins are shown in Fig. 1.

Among three types of tea, black tea is the most popular tea produced and consumed preferentially in the United State, England, and other Western countries with 78% of global market, followed by 20% of green tea consumed primarily in Asian and Northern African countries, and about 2% of oolong tea consumed mainly in Taiwan, southern China, and most Eastern countries [13,14]. It has been noted that production and consumption of oolong tea worldwide have increased over the past decades. For example, the production of oolong tea in China from 2000 to 2014 had been nearly doubled and increased from 67.6×10^3 to 254×10^3 metric tons [15]. Catechin prior to oolong tea is oxidized in the range of 10–80% during processing depending on the demand of customers [9,16]. The taste quality of oolong tea depends on several properties, such as smell of volatile fragrance, taste sensation of sweetness, umami, and intensity of astringency. The differentiation of green tea, black tea and oolong tea, regardless of degree of fermentation, is also depended on their contents of free amino acids, mainly *L*-theanine and several natural amino acids including glutamic acid, asparagine, serine, alanine, leucine, and isoleucine [17].

Major contents of oolong tea infusion are listed in Table 2, which has two categories: monomeric polyphenols and polymeric substances. Oolong tea have been demonstrated to possess various pharmacological activities such as antioxidant activity




Fig. 1. Chemical structure of tea catechins (EC, ECG, EGCG, EGC), theaflavins, and theasinsensins A–E.

Table 2
Components of oolong tea beverage.

| Compounds | Contents (mg/100 mL) |
|--------------------------|----------------------|
| Catechin | 1.65 |
| Gallocatechin | 6.68 |
| Epigallocatechin | 16.14 |
| Epicatechin | 5.08 |
| Catechin gallate | 0.6 |
| Epicatechin gallate | 5.73 |
| Epigallocatechin gallate | 25.73 |
| Allocatechin gallate | 1.85 |
| Gallic acid | 2.19 |
| Caffeine | 23.51 |
| Polymerized | 33.65 |
| Total polyphenols | 99.32 |

Sajilata et al. [5].

by reducing oxidative stress, anti-cancer, anti-obesity, anti-diabetes, preventive effect of atherosclerosis, heart disease, hypertension, anti-allergic effect, and antiseptic effects [18–27]. However, all of the studies were applied to oolong tea extract, not single characteristic oolong tea polyphenols. Until 1984, a new group of polymeric oxidized flavan-3-ols was isolated and identified from oolong tea as theasinensins A, B, C, and later in 1988 for D, E, F and G, have been confirmed from oolong tea by the Japanese scientists Nonaka and Hashimoto [28–33]. As a matter of fact, this group of compounds was formerly discovered by Robert in 1958 [34,35] and was known as bisflavanol A, B, and C, which were formed by coupling of EGCG [36]. Based on the literatures listed above, theasinensins were implied as the bioactive flavonoids in oolong tea.

2. Theasinensins – structure and occurrence

Theasinensins are formed from catechin dimerization at their B-rings, *i.e.* two catechin B-rings are connected through C–C bonds [35,37]. The major tea polyphenol components found in black and oolong teas are shown in Fig. 1. The configuration of theasinensins A, B, and C differ from that of theasinensins D and E in which the biphenyl bonds of theasinensins A, B, and C, carrying a *R*-biphenyl configuration, whereas theasinensins D, and E embed with *S*-biphenyl bonds. Hashimoto et al. [33] also concluded that theasinensins D and E were characterized as the atropisomers/stereoisomers of theasinensin A and C, respectively. Theasinensin A and D are the dimers of EGCG with an *R*- and an *S*-biphenyl bond, respectively [37]. Theasinensin B is the dimer of EGCG and EGC and theasinensin C is the dimer of EGC [38].

Theasinensins, mainly exist in black tea and oolong tea [39], are transformed from the unstable intermediate produced by the oxidation of original catechins in tea leaves [28–33]. Among tea catechins found in fresh tea leaves, EGCG is the dominant catechin (63%) followed by EGC (25%) [5,40,41]. Therefore, oxidation of EGC and EGCG, two pyrogallol-type catechins, is important during fermentation process. Many studies had been focused on the oxidation of these two catechins to understand the formation of theasinensins. Several studies have been designed to clarify the underlying mechanisms of theasinensin formation in the fermentation process including a model system to mimic catechin oxidation [33,35,37,42,43].

3. Proposed formation mechanism of theasinensin via gallocatechin dimerization

Hashimoto et al. [42] indicated that tea catechins are easily transformed to theasinensins by endogeneous polyphenol oxidase than to theaflavins which are the characteristic flavonoids in black tea. This finding has unequivocally confirmed that theasinensin production involved in the enzymatic oxidative coupling of two pyrogallol rings of EGCG. The pyrogallol rings are also susceptible to be oxidized by chemical agents [44]. There are different pathways toward the formation of theasinensins and theaflavins. The benzotropolone core of TFs is formed from an oxidative condensation reaction between a catechol moiety of

EC or ECG and a pyrogallol moiety of EGC or EGCG [42], whereas theasinensin carrying a *R*-biphenyl bond is generated by a pyrogallol-type B rings of EGCG and EGC.

Once the fresh tea leaves are crushed and kneaded, TFs are formed in the leaves, but theasinensins are not detected in this step [28,33]. The formation of theasinensins is observed in heating the leaves to 80 °C [28,33,35]. Several researches who tried to investigate theasinensin formation pathway initially focused on characterizing the intermediates and later known that they were heat and chemical susceptible intermediates. The structure elucidation from derivatization of the unstable intermediates has concluded that they are quinone dimers of EGC and EGCG produced by these two catechin quinone monomers. To date, one of the intermediates has been successfully synthesized by an enzymatic oxidation of EGCG and named dehydrotheasinensin A with a pale yellow color [37,43]. In the heating and drying process of the tea leaves at the final stage of black and oolong tea manufacturing, dehydrotheasinensin A undergoes redox dismutation to generate theasinensin A and its atropisomer [37,43]. The production of unstable dehydrotheasinensins from the oxidation of pyrogallol-type B-ring catechins is the most important reaction in theasinensin formation. Takana et al. [38] have utilized an *in vitro* experiment to investigate the chemical mechanism of theasinensin formation and the proposed pathway is also given in Fig. 2.

More than a decade ago, the oxidized EGCG and EGC as theasinensin quinones were trapped for the first time with *o*-phenylenediamine, yielding five phenazine derivatives of catechin dimer quinones [35]. The result was also the first evidence supporting the formation and accumulation of catechin dimer quinones in tea fermentation process and proving that the formation mechanism of theasinensin A is not simply produced by enzymatic oxidation of EGCG, the main catechin in fresh tea leaves. These theasinensin quinones are unstable and decomposed readily in particular the process of extract solution concentration by rotary evaporator [35]. Later, Tanaka's team also suggested that the addition of 0.1% trifluoroacetic acid (TFA) in the mobile phase increased the stability of theasinensin quinones [37].

Tanaka and his team were also focused on the chemical examination of theasinensin production using *in vitro* experiments to mimic black tea production process in particular on the fermentation step. EGCG and enzymes from homogeneous pear were applied at room temperature for 2 h. The study elucidated the structure of dehydrotheasinensin A and mechanism of theasinensin formation, and successfully obtained dehydrotheasinensin A, a theasinensin precursor from enzymatic oxidation of EGCG. The results also revealed that dehydrotheasinensin A produced from coupling of EGCG quinone monomers is equivalent to a hydrated *ortho*-quinone of the theasinensin A, which is readily converted to theasinensin A and D by oxidation reduction dismutation. Previous studies from Takana et al. [35,37] suggested that several chemical reactions involved in EGCG oxidation subsequently produced theasinensins (A and D).

EGCG was initially oxidized to the *ortho*-quinone, and subsequently an unstable dimer, theasinensin quinone was formed after stereoselective dimerization (Fig. 2). This unique quinone




Fig. 2. Possible formation mechanism of theasinensin.

was equivalent to dehydrotheasinensin A, generated by both enzymatic and non-enzymatic reaction [28,33,35,37,43,45,46]. To further understand the chemical mechanism of theasinensin formation, many studies had concentrated on the enzymatic oxidation of catechins such as EGCG oxidation. However, several studies also focused on the non-enzymatic oxidation of catechins.

The non-enzymatic oxidation of EGCG providing theasinensin quinones can be developed from not only chemical oxidation with either potassium ferricyanide or copper salts, but also autoxidation in a phosphate buffer (pH 7.4 or 6.8) in which the oxidized product is unstable even at a neutral pH [37]. Therefore, it can be implied that the production of theasinensin A and D derived from dehydrotheasinensin A during black tea and oolong tea manufacturing may possibly be a non-enzymatic process which occurs spontaneously when tea leaves are heated and dried. Shii further showed that non-enzymatic oxidation of EGCG by copper salts with the optimum pH of 4–5 at room temperature yielded dehydrotheasinensin A and the product increased with the elevated temperature [43]. Initial oxidation reaction was carried out at room temperature and elevated temperature increased byproduct generation. The difference between enzymatic oxidation in tea leaves and *in vitro* chemical oxidation by copper salt is that catechol-type catechins prefer to enzyme catalyzed oxidation, such as EC and ECG, whereas pyrogallol-type catechins, such as EGC and EGCG, are less reactive [43,45]. By enzymatic oxidation (banana homogenate) of catechin, EGC was oxidized to the corresponding catechin quinone but the reaction was accelerated in presence of ECG. The oxidation of EGC was much faster than EGC due to its higher redox potential than pyrogallol-type catechin (EGCG, EGC) [35,43]. Technically, the catechol-type catechins were not chemically oxidized by copper salt. Therefore, the chemical oxidation of catechins does not yield theaflavins which are generated from the oxidative

couplings between the quinones produced from catechol-type and pyrogallol-type catechins [47–49].

Enzymatic oxidation of EGC generates a new quinone dimer with a hydrated cyclohexenetrione structure, which may be equivalent to dehydrotheasinensin C whereas theasinensins C and E were produced through oxidation–reduction dismutation [39]. In addition, the enzymes isolated from pear homogenates catalyzed the oxidation of EGC at the pyrogallol B-ring, yielding predominantly unstable quinone products known as dehydrotheasinensin C trapped with *o*-phenylenediamine [45]. Non-enzymatic coupling reaction of dehydrotheasinensin C was subsequently occurred to give theasinensins A and C [37,38,45,46].

From commercial back tea, an interesting derivative of theasinensin, *N*-ethylpyrrolidinonyl theasinensin has been isolated and identified [44,50]. Tea leaves contain an amino acid named *L*-theanine (*N*²-ethyl-*L*-glutamine) accounting for over 50% of the total amino acids. Consequently, the catechin quinones generated from the beginning of tea process maybe react with *L*-theanine to generate theanine Strecker aldehyde [51,52] as demonstrated in Fig. 3. The Strecker aldehyde of theanine subsequently reacts with theasinensin to yield *N*-ethylpyrrolidinonyl theasinensin.

In addition, the reduction of dehydrotheasinensin A to yield theasinensin derived from several paths. The formation of theasinensins through several chemical reactions was involved in EGCG oxidation [35,37,43]. Dehydrotheasinensin A could be reduced by the reductants such as ascorbic acid or sulfur containing compounds at room temperature and subsequently forming theasinensin A [37]. It is also known that reduction of dehydrotheasinensin A with 2-mercaptoethanol afforded theasinensin A only. Stereoselective formation of dehydrotheasinensin A from EGCG created a chiral center at the benzylic position of the pyrogallol ring [46,53]. Unstable dehydrotheasinensin A was rapidly degraded by heating alone or heating with ascorbic acid, higher




Fig. 3. Proposed route for the formation of *N*-ethylpyrrolidinonyl theasinensin A [44,50].

pH, and theasinensin A subsequently was formed [37,43]. Even so, the reducing agent such as 2-mercaptoethanol had lower yield of theasinensin A compared to ascorbic acid [43]. On the other hand, the reduction of dehydrotheasinensin A occurred at 80 °C and thermodynamically underwent the oxidation–reduction dismutation to yield theasinensin A. Theasinensin D also was produced as it is probably isomerized at an elevated temperature during heating and drying in black tea manufacturing. Furthermore, theasinensin quinones were gradually decomposed to a mixture of theasinensin A and D in a pH 6.8 phosphate buffer at 20 °C [37]. Taken together, it is clear that the products from enzymatic oxidation of tea catechins both in oolong and black tea process can be classified into two major oxidation routes [4,35,38,42,46]: the condensation with coexisting epicatechin forming theaflavins [47,48] and oxidative dimerization of two gallo moiety and reduction yielding epigallocatechin dimers, such as theasinensins and oolongtheanins [28,33].

4. Potential bioactive compounds

Evidence from experimental and clinical studies has indicated that tea exerts antioxidative, anti-inflammatory, cancer prevention and vasodilating effects among others [54]. Epidemiological studies, both *in vitro* and *in vivo*, have indicated that tea consumption is positively associated with reduced risk of chronic diseases, causing 60% global death such as coronary heart disease, stroke and cancer [54–58]. One of the health benefits of tea polyphenols is generally attributed to

its potent antioxidant property. Act as free radical scavengers, tea polyphenols may remove endogenously superoxide, peroxy, and hydroxyl radicals [22,59,60]. Tea polyphenols and their metabolites also possess antibacterial properties against pathogenic bacteria such as *Clostridium perfringens*, *Clostridium difficile*, *Escherichia Coli*, *Salmonella*, and *Pseudomonas* and enhance probiotics such as *Bifidobacterium* and *Lactobacillus* species, which improve the intestinal microbial balance [61,62].

The study on bioactivities of oolong tea and its characteristic compounds, theasinesins on health promotion property in the literature is still limited compared to that of black and green teas. Several studies have revealed that oolong tea and theasinesins have biological activities such as antioxidative effects against lipid peroxidation, anti-inflammatory activity, antibacterial properties and anti-obesity. Herein, we summarize the health benefits of oolong tea and theasinensins in Tables 3 and 4.

4.1. Antioxidant activities of oolong tea and theasinensins

The FRAP (ferric reducing/antioxidant power) assay has tested that antioxidant value of oolong tea ranged between 233 and 532 $\mu\text{mol/g}$ [63]. In an *in vivo* study, modest transient increase in human plasma antioxidant capacity was noticed upon oolong tea consumption [22]. It is observed that oolong tea reduced oxidative stress, especially oxidative DNA damage [22]. Furthermore, human studies on athletes showed that oolong tea ingestion significantly reduced plasma malondialdehyde levels in rest and post-exhaustive exercise athletes, as well as

Table 3
Health benefits and proposed molecular mechanisms of theasinensins.

| Bioactivity | Experimental model | | Compound tested/control | Mechanism/biomarker | Ref. |
|-----------------------|---|---------------------------------------|--------------------------------------|--|-------|
| | <i>In vitro</i> | <i>In vivo</i> | | | |
| Antioxidant activity | Ferric thiocyanate assay | | Theasinensins A–E/alpha-tocopherol | Decreasing lipid peroxidation | [64] |
| Anti-inflammation | LPS-activated murine macrophage RAW264.7 cells | | Theasinensins A–E | Reducing gene expression of cyclooxygenase-2 (COX-2) and PGE ₂ | [68] |
| | LPS-activated murine macrophage RAW264.7 cells (a genome-wide microarray) | | Theasinensin A | 22,050 genes of inflammatory and immune response | [69] |
| Anti-cancer | LPS-activated murine macrophage RAW264.7 | Mouse paw edema model | Theasinensin A | Reducing the production of NO/iNOS, IL-12 (p70), TNF- α , and MCP-1 | [70] |
| | Human fibrosarcoma HT1080 cells | | Theasinensin D | Suppressing invasion by reducing Gelatinase/Type IV Collagenases (MMP-2 and -9) activities | [84] |
| Hypoglycemic effect | Human histolytic lymphoma (U937) cell line and acute T cell leukemia (Jurkat) cell line | | Theasinensin A | Inducing DNA fragmentation, and caspase activation | [85] |
| | Rat skeletal muscle cells | KKAy mice and Sprague-Dawley rats | Theasinensin A | Regulation of serum glucose, lipid serum, hepatic fatty acid synthase activity | [9] |
| Anti-microbial effect | Alpha-glucosidase from rat intestinal acetone powder | | Theasinensin A and B | Increasing glucose uptake | [90] |
| | | | Catechin, theaflavin, theasinensin A | Increasing alpha-glucosidase (AGH) inhibitory activity | [89] |
| Anti-microbial effect | | MRSA (strains OM48, 505,584, and 623) | Theasinensin A and EGCG | Increasing antibiotic resistance | [103] |
| | | HSV-1 and HSV-2 | Theasinensin A, theaflavin, and EGCG | Enhancing protein aggregation | [108] |

NBDG; 2-[N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)amino]-2-deoxyglucose; herpes simplex virus (HSV)

MRSA; methicillin-resistant *Staphylococcus aureus*.

resting levels of superoxide dismutase activity, suggesting that the decrease of oxidative stress is resulted from reduction of the lipid peroxidation level and its free radical scavenging activity [25]. Hashimoto reported that the antioxidant activity evaluated by the 5-day lipid peroxidation of theasinensins A–E were ranged from 9 to 13% compared to 3 and 17% for BHA and alpha-tocopherol, respectively. The result suggests that the inhibitory activity on lipid oxidation of theasinensins A–E were lower than BHA (the synthetic antioxidant), but higher than alpha-tocopherol [64]. Among theasinensin isomers, theasinensin C has the highest ability against lipid oxidation inhibition. Until now, there is no data reported concerning on bio-antioxidative effect of oolong tea or theasinensin on the antioxidant defense systems.

4.2. Anti-inflammatory effect

Previous studies have indicated that ethanol extract of oolong tea was profoundly increased adiponectin gene expression in epididymal fat, consistent with an anti-inflammatory effect, and angiogenesis during adipose tissue expansion [65]. The increase of blood vessel formation in adipose tissue contributes to their anti-inflammatory effects by maintaining adipocyte perfusion [65]. This finding is consistent with the pro-angiogenic activity of oolong tea ethanol extract during adipose tissue expansion

mediating protective effects on metabolism and inflammation, although this finding is in contrast to the traditional concept that inhibition of angiogenesis results in weight loss [65,66]. In addition, ethanol extract of oolong tea decreased the concentration of monocyte chemoattractant protein-1 (MCP-1) in serum. The reduction of plasma MCP-1 also explains the anti-inflammatory effect of oolong tea polyphenol since MCP-1 is a small cytokine recruiting monocytes, memory T cells, and dendritic cells to the site of injury [67]. The anti-inflammatory activity of oolong tea might be due to flavonoids such as catechins, theaflavins and theasinensins [68]. There are some reports on the anti-inflammatory properties of theasinensins. Results from the first molecular basis of the anti-inflammatory properties of oolong tea theasinensins showed that theasinensin A and D exhibited better activities in reducing LSP-stimulated COX-2 and PGE₂ production than theasinensins B, C and E. Structure analysis of theasinensins A and D with two galloyl moieties and theasinensins C and E without galloyl moieties, showed that the galloyl moiety played an important role on anti-inflammatory effects of oolong tea theasinensins. Molecular mechanisms of anti-inflammatory property of oolong tea theasinensins demonstrated that the down-regulation of TAK1-mediated NF- κ B and MAPK signaling pathway might be involved in the inhibition of COX-2 expression by theasinensin A [68]. Recently, in clarifying the molecular mechanism of anti-inflammatory effects of

Table 4
Health benefits of oolong tea.

| Bioactivity | Experiment model | | Compound tested/control | Mechanism/biomarker | Ref. |
|----------------------|---|---|---|--|-------|
| | <i>In vitro</i> | <i>In vivo</i> | | | |
| Antioxidant activity | Lipoxygenase inhibition activity | | Fraction compound from oolong tea | | [114] |
| | | FRAP assay | Tea infusion (green, oolong and black tea) | Antioxidant power | [63] |
| | | Human plasma | Consumption oolong tea infusion | Reducing oxidative DNA damage | [22] |
| | | Athletes before and after exhaustive exercise | Daily consumption of oolong infusion for 30 days | Normalizing the cholesterol profiles, reducing lipid peroxidation level, and superoxide dismutase activity | [25] |
| Anti-inflammation | | Mice | Crude ethanol extract of oolong tea | Reducing MCP-1 plasma concentration | [65] |
| Anti-obesity | | Mice | Crude ethanol extract of oolong tea | Reducing MCP-1 gene expression | [65] |
| | | High-fat diet-induced obese mice | Oolong tea | Fat cells and a cell-free system consisting of lipid droplets and hormone-sensitive lipase | [72] |
| | | Rats | Oolong, black, pu-erh, and green teas leaves | Reducing plasma lipid Increasing plasma enzyme SOD Weight ratios of liver to epididymal adipose tissue | [74] |
| | | | Fifty-four polyphenols isolated from tea leaves Oolonghomobisflavans A and B | Elevating pancreatic lipase activity | [73] |
| Anti-cancer | Salmonella/microsome reverse mutation assay (<i>Salmonella typhimurium</i> TA98 and TA100) | | Tea water extract (green, oolong, pouchong, and black teas) | Anti-mutagenicity | [19] |
| | Salmonella/microsome reverse mutation assay (<i>Salmonella typhimurium</i> TA98 and TA100) | | Tea water extracts (green, pouchong, oolong and black tea) | Anti-mutagenicity | [79] |
| | Male F344 rats induced by diethylnitrosamine (rat model is <i>in vivo</i> study) | | Tea extract (black and oolong tea/tea catechins) | Suppressing hepatocarcinogenesis | [19] |
| | Salmonella/microsome reverse mutation assay (<i>Salmonella typhimurium</i> TA100, TA98 and TA97) | | Tea water extracts (green, pouchong, oolong and black tea/EGCG, gallic acid and caffeine) | Anti-mutagenic activities | [78] |
| | AH109A rat ascites hepatoma cell line | | Tea water extracts (green, oolong and black tea/catechin, theaflavins) | Preventing proliferation and invasion of AH109A cells | [21] |
| | | Male Donryu rats | Tea water extracts (green, oolong and black tea/catechin, theaflavins) | Inducing apoptosis, and cell cycle arrest | [81] |
| | Human stomach cancer KATO III cells | | Oolong tea extract | Apoptosis by fragmentation of DNA to oligonucleosomal sized fragments | [82] |
| Hypoglycemic effect | Salmonella/microsome reverse mutation assay (<i>Salmonella typhimurium</i> TA1535/pSK 1002) | | Tea methanol extract (green, pouchong, oolong and black tea) | Anti-genotoxic abilities (suppressive effects against <i>umu</i> gene expression) | [80] |
| | | Adult female frogs (<i>Xenopus laeVis</i>) | Tea extract by hot normal frog Ringer Solution (green, oolong, and black tea) | Inhibition of SGLT1 Response | [86] |
| | Rat epididymal adipocytes | | Tea infusion (green, oolong, and black tea) | Enhancing insulin-activity | [87] |

Table 4 (Continued)

| Bioactivity | Experiment model | | Compound tested/control | Mechanism/biomarker | Ref. |
|-----------------------------|---|--|---|---|-------|
| | <i>In vitro</i> | <i>In vivo</i> | | | |
| Prevention of Heart Disease | | 10 men and 10 women (average age 61.2 years, duration of diabetes 4.8 years) Male Sprague-Dawley rats | 1500 ml of oolong tea per day (tea bags) for 10 weeks | Decreasing plasma glucose and fructosamine | [88] |
| | | | Tea solutions for 9 weeks (green, Jasmine, Iron Buddha, oolong and Pu erh tea) | Decreasing serum and liver lipids | [101] |
| | | 12 healthy men | Water plus caffeine and oolong tea | Increasing EE and fat oxidation | [100] |
| | | 12 patients with previous myocardial infarction and 10 patients with stable angina pectoris | Medication during their additional oolong tea for 1 month | Decreasing plasma adiponectin, glucose and hemoglobin A1c levels | [97] |
| | | 12 healthy university students, 3 males and 9 females | The polymerized-polyphenol extract from oolong tea as beverage for 10 days | Increasing fecal lipid excretion | [98] |
| | | 12 healthy Japanese females | Oolong tea and green tea | Increasing energy expenditure and resting energy expenditure Anti-hypertension | [99] |
| Anti-microbial effect | | 711 men and 796 women without hypertensive history | Epidemiology study (tea consumption) | | [94] |
| | <i>Streptococcus sobrinus</i> 6715 | | Oolong tea extract and its polymeric polyphenols | Inhibition of GTase | [105] |
| | Pathogen | | Aqueous tea extract (green and oolong tea) catechin, theasinensin A, theaflavin | Increasing MNIC and MIC | [20] |
| | <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Proteus vulgaris</i> , <i>Pseudomonas fluorescens</i> , <i>Salmonella</i> sp. and <i>Staphylococcus aureus</i> | | Dry tea extract (green, oolong, and black tea) | Decrease of growth and survival | [104] |
| | <i>S. mutans</i> MT8148R and <i>Streptococcus sobrinus</i> 6715 | | Methanol extract of oolong tea | Antibacterial activity | [106] |

Superoxide dismutase (SOD); minimum inhibitory concentration (MIC); maximum non-inhibitory concentration (MNIC); energy expenditure (EE); glucosyl transferase (GTase).

theasinensin A, gene expression profiling in macrophage-like cells treated with theasinensin A through a genome-wide DNA microarray were used to detect the changes of 22,050 genes involving inflammatory and immune response. The changes of 1382 genes suggested that theasinensin A has exerted anti-inflammatory effects by regulating the relevant expression networks of chemokines, interleukins, and interferons [69]. Recently, an *in vitro* study on anti-inflammatory activity of theasinensin A by LPS-activated macrophages indicated that the levels of pro-inflammatory mediators including inducible nitric oxide synthase (iNOS), nitric oxide (NO), interleukin-12 (IL-12) (p70), tumor necrosis factor alpha (TNF- α), and MCP-1 were significantly reduced by theasinensin A. Cellular signaling pathway of this study uncovers that theasinensin A downregulated MAPK/ERK kinase (MEK)-extracellular signal-regulated kinase (ERK) signaling through directly binding to MEK-ERK for the inhibitory action. The *in vivo* study also demonstrates that the theasinensin A suppressed the production of IL-12 (p70), TNF- α , and MCP-1 and attenuated mouse paw edema induced by LPS [70].

4.3. Anti-obesity

Nowadays obesity has emerged as a major health concerning problem and a risk factor of metabolic disorders. Functional foods affecting energy metabolism and fat partitioning may be helpful adjuncts to a dietary approach to weight control. The mechanism of anti-obesity effects of tea catechins, especially EGCG and theaflavins, appear to be related to modulation of energy balance, endocrine systems, food intake, lipid and carbohydrate metabolism, and activities of different cell types including fat, liver, muscle and β -pancreatic cells [7,27]. Tea catechins inhibit lipogenesis through down-regulation of gene expression of fatty acid synthases in the nucleus, resulting in the down-regulation of EGFR/P13K/Akt/Sp1 signal transduction pathways, and stimulation of cell energy expenditure in the mitochondria [27,69]. One of the mechanisms on fat reduction by oolong tea is the inhibition of pancreatic lipase activity. A mouse study suggested that inhibition of digestive lipase activity can significantly affect dietary lipid absorption and increase lipid excretion into the feces [71]. Another study was

also demonstrated that the oolong tea water extract enhanced noradrenaline-induced lipolysis and inhibited pancreatic lipase activity, resulting in anti-obesity effects. This might be partially due to the effects of caffeine or some other bioactive compounds in oolong tea [72]. The homobisflavans A and B, typical compounds in oolong tea, exhibit more potent inhibitory activities against pancreatic lipase with IC₅₀ values of 0.048, and 0.108 $\mu\text{mol/L}$, respectively, than EGCG with IC₅₀ values of 0.349 $\mu\text{mol/L}$. The data hinted that galloyl moieties in the molecule were crucial for pancreatic lipase inhibition [73]. Supplementation with oolong tea leaves reduced body weights and plasma triacylglycerol, cholesterol and LDL-cholesterol of rats. Pu-erh tea and oolong tea can significantly decrease the levels of triacylglycerol more significantly than green tea and black tea. In addition, superoxide dismutase activity (SOD) was increased and the relative weight ratios of liver to epididymal adipose tissue reduced by oolong tea. This study also suggested that oolong tea was more effective on their growth suppressive and hypolipidemic effects as compared to green tea [74]. Scientific evidence has revealed the role of MCP-1 gene expression in the etiologies of obesity- and diabetes-related diseases [75]. Plasma concentration of MCP-1 is positively associated with obesity [67]. A recent study shows that the ethanol extract of oolong tea polyphenol caused weight loss in mice fed with a high-fat diet, and decreased plasma MCP-1 protein as well as its gene expression in mesenteric fat and epididymal fat [65].

4.4. Anti-cancer

Among the biological activities of tea polyphenols, the cancer-chemopreventive effects in various animal models have been intensively investigated [76]. Many studies have reported that the anti-cancer effect of oolong tea mostly focused on catechins. However, some studies paid much attention to evaluate the anti-cancer effects of theasinensins. It is well known that inflammation plays a key role in the initiation and/or progression of multiple types of cancers, including liver, bladder and gastric cancers by inducing oxidative stress and promoting cell growth [77]. As mentioned above, oolong tea extract exhibits a higher anti-inflammatory activity than green tea and black tea extracts [65]. The anti-cancer activity of oolong tea may be originated from its antioxidant and anti-inflammatory activities.

The mutagenic effects of carcinogens such as heterocyclic amines are reduced by oolong tea extracts. In *Salmonella* reverse mutation assay (Ames test), Yen and Chen [18] indicated that the greater anti-mutagenic effect was found in oolong tea than in green tea and black tea and some anti-mutagenic substances might be formed during manufacturing processes of tea. Ames test with bacterium *Salmonella typhimurium* has confirmed anti-mutagenic activities of EGCG, GC, and caffeine from green tea [78]. Furthermore, oolong tea extract remarkably inhibited the mutagenicity of 2-amino-3-methylimidazo(4,5-*f*)quinoline (IQ), 3-amino-1,4-dimethyl-5-*H*-pyrido-(4,3-*b*)indole (Trp-P-1), 2-amino-6-methyl-dipyrido(1,2-*a*:3',2'-*d*)imidazole (Glu-P-1), benzo[*a*]pyrene (B[*a*]P) and aflatoxin B, (AFB₁) and the

inhibitory effect was associated with the contents of catechins and ascorbic acid [79]. Oolong tea extract possessed the highest anti-mutagenic activity against several mutagens, including four nitroarenes, two nitro compounds and one alkylating agent (1-nitropyrene (1-NP), 2-nitropyrene (2-NF), 3-nitropyrene (3-NF) and 2,4-dinitrophenol (DNP) among the different teas [79]. In addition, oolong tea extracts have a chemopreventive action against hepatocarcinogenesis. Different concentrations of oolong tea extract (0.05 or 0.1%) significantly decreased the number and area of diethylnitrosamine- and phenobarbital-induced preneoplastic glutathione S-transferase placental form-positive foci in the liver of rats [19]. Oolong tea extracts also inhibited the formation of reactive oxygen species (ROS) and induced cytochromes P450 1A1, 1A2, and 2B1, and glucuronosyl transferase, leading to glucuronide, which is an important mechanism in biological detoxification system [76]. Therefore, one of the anti-carcinogenic mechanisms of oolong tea may be involved in regulation of catalytic activities of the P450 enzymes and glucuronosyl transferase [7,76].

Besides the anti-inflammatory effects, anti-genotoxic and anti-mutagenic properties of oolong tea also contribute to its chemopreventive effect. Zhang et al. [21,81] suggested that the chemopreventive mechanism of oolong tea might be attributed to its bioactivities against the invasion and proliferation of cancer cells (AH109A) through the loss of cell viability, apoptosis, and cell cycle arrest at the G1 phase in a rat hepatoma cell line (AH109A) and murine B16 melanoma cells. Furthermore, the oolong tea extract inhibited the growth of human stomach cancer KATO III cells by the induction of apoptosis [82]. Saeki et al. [83] has confirmed that a pyrogallol type structure in the B-ring of catechin induced apoptosis of cancer cells compared with the catechins without a pyrogallol-type.

In 1999, an *in vitro* study showed that comparing with ECG and EGCG, theasinensin D and theaflavin-3,3'-digallate exhibited a weak invasion inhibitory effect determined by the suppression of the gelatin degradation mediated through matrix metalloproteinase (MMP) [84]. However, using human histolytic lymphoma (U937) cell line and acute T cell leukemia (Jurkat) cell line, the study demonstrates that theasinensin A from oolong tea induced the process of cell death through the release of cytochrome *c* and activation of caspase-9 and caspase-3 [85]. Their results also suggested that a linear and specific activation cascade between caspase-9 and caspase-3 in response to cytochrome *c* released from mitochondrial and apoptosis by activation of the caspases, leading to the process of cell death [85].

4.5. Hypoglycemic effect

Hyperglycemia is the major cause of diabetic angiopathy. The Na⁺-dependent glucose cotransporter (SGLT1) in jejunum transports glucose into epithelial cells. The inhibition of glucose uptake *via* SGLT1 in the small intestine may prevent hyperglycemia [86]. Aqueous extracts of green, oolong, and black tea failed to exhibit effects on SGLT1 response compared with tea catechins. Black, green, and oolong teas as tea beverages had

shown to increase insulin activity by a minimum of 15-fold in an epididymal fat cell assay [87]. Oolong tea extract can reduce plasma glucose and have a complicated impact on antioxidant systems in diabetic rats [7,87]. Oolong tea is an effective adjunct to oral hypoglycemic agents in the treatment of type 2 diabetes. An *in vivo* study with respect to type 2 diabetes subjects indicated that oolong tea remarkably reduced concentrations of plasma glucose from 229 to 162.2 mg/dL and fructosamine (from 409.9 to 323.3 $\mu\text{mol/L}$) [88]. This study also strongly supports the concept of combination therapy because the ingestion of oral anti-hyperglycemic agents and oolong tea simultaneously was more effective in lowering plasma glucose than taking the drugs alone [88].

Theasinensins might play a key role in anti-hyperglycemic activity of oolong tea. Theasinensin A induces anti-hyperglycemic responses in diabetic mice and shows hypotriacylglycerolemic effect in rats by suppressing intestinal fat absorption [89]. Miyata indicated that feeding male KK-Ay mice with diets containing 0.1% theasinensin A for 6 weeks reduced serum glucose levels by greater than 30% and feeding rats with diets containing 0.2% theasinensin A for 4 weeks had higher fecal fat excretion and 33% lower hepatic triacylglycerol without the effect on hepatic fatty acid synthase activity [10]. High fecal excretion of fatty acid is associated with the inhibition of pancreatic lipase by theasinensin A [89]. The result suggested that it might be due to suppression of postprandial hypertriacylglycerolemia, theasinensin A inhibited glucose production in the intestine through suppression of α -glucosidase activity. Theasinensin A inhibited α -glucosidase (AGH) activity (IC₅₀ of 142 $\mu\text{mol/L}$ for maltase and 286 $\mu\text{mol/L}$ for sucrase) evaluated by an immobilized AGH assay system and thus provides a substantially useful prediction of the *in vivo* suppression of glucose absorption [89]. Qiu et al. [90] proposed the mechanism of anti-hyperglycemic activity of theasinensins A and B using rat skeletal muscle cells (L6 myotubes). Theasinensins A and B were found to promote GLUT4 translocation to the plasma membrane in L6 myotubes through the CaMKK/AMPK signaling pathway, but not through the PI3K/Akt pathway. Moreover, theasinensin A is more effective in stimulating 2-NBDG (2-(*N*-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)amino)-2-deoxyglucose; a fluorescent glucose analog) uptake than theasinensin B, which is as potent as EGCG, suggesting that the number of galloyl moieties may be associated with the promotion of 2-NBDG uptake in cells [90]. GLUT4 plays a pivotal role in regulating insulin-stimulated glucose transport predominantly in muscle of skeleton, cardiac and adipose tissue [91]. Glucose uptake by muscle and fat cells is regulated by modulating the number of GLUT4 on cell surface. Promoting glucose uptake or improving insulin resistance by increasing GLUT4 translocation to the plasma membrane *via* the AMP-activated protein kinase (AMPK) pathway raises the possibility to prevent hyperglycemia. The effects of theasinensins A and B on regulation of GLUT4 are differs from that of EGCG that involved in stimulating the phosphatidylinositol 3-kinase (PI3K)/Akt pathway in skeletal muscle to improve insulin sensitivity [90].

4.6. Prevention of atherosclerosis, heart disease, and hypertension

High blood pressure affects millions of people globally [92] and it is associated with atherosclerosis and plaque build-up in the arteries, both of which are considered as main cardiovascular risk factors [93]. The prevention of atherosclerosis can also reduce the risk of developing hypertension and heart disease. An epidemiology study has shown that people with habitual and moderate oolong tea consumption, such as 120 ml/day or more for a year period, significantly reduced the risk of developing hypertension in Taiwan [94].

Not only the level but also the particle size of low-density lipoprotein (LDL) is the major risk factor in the early development of coronary heart disease (CAD) [95,96]. Small particle size of LDL is considered as a great risk factor for CAD [97]. Oolong tea increased plasma adiponectin levels and LDL particle size in CAD patients and decreased hemoglobin A1c (HbA1c). Therefore, oolong tea may have beneficial effects on the progression of atherosclerosis in patients with CAD. As mentioned earlier, oolong tea decreased levels of triglyceride in Sprague-Dawley rats and lowered the relative weight ratios of liver to epididymal adipose tissues [74]. The result also showed that oolong tea was more effective in suppressing the growth of adipose tissues as compared to green tea [74]. In addition, oolong tea may inhibit the oxidized-LDL cholesterol, which is a risk factor for atherosclerosis and heart disease, and reduced the formation of 8-hydroxydeoxyguanosine, a marker of oxidative DNA damage [76].

There is an affiliation between the ingestion of dietary lipid and some diseases such as obesity and cardiovascular diseases. Hence reduction of lipid intake is a logical nutritional intervention strategy. Moreover, the more energy expenditure means less accumulation of lipid, which can decrease the incidence of atherosclerosis and heart disease. To absorb dietary lipids, the lipid digestion by pancreatic lipase is a key step. Inhibition of digestive lipase activity significantly affects dietary lipid absorption, leading to increase lipid excretion into the feces [71]. Thus, it is a possible strategy responsible for the prevention of atherosclerosis and heart disease. Polyphenol-enriched oolong tea extract increased lipid excretion into feces in subjects with high fat diet, and polymerized-polyphenol extract from oolong tea appears to increase cholesterol excretion into feces [98]. Furthermore, the consumption of oolong tea in healthy women leads to the increase of energy expenditure (EE) and this effect is higher than that of green tea [99]. The same trend of the effect of oolong tea on energy expenditure was reported in Japanese subjects [99] and the result showed that oolong tea stimulates fat oxidation. Therefore, the consumption of oolong tea increased metabolic rate and fat oxidation, potential beneficial effect on an individual's ability to maintain a lower body fat content, associated with lower risk of atherosclerosis and hypertension [100]. Oolong tea decreased atherogenic index and increased HDL-total cholesterol ratio in hypercholesterolemia rats [101]. However, there is still no report of the bioactivity of theasinensins on prevention of atherosclerosis and heart disease in the literature.

4.7. Antiseptic effects

Oxacillin (methicillin)-resistant *Staphylococcus aureus* (MRSA) caused infection is a problem in health care institutions in the United States and worldwide, especially for intensive care unit patients [102]. This type of bacteria is resistant to a number of widely used antibiotics, so MRSA infections can be more difficult to cure than other bacterial infections. Aqueous extract of oolong tea inhibits various pathogens including *Staphylococcus aureus* (a methicillin-resistant strain). A previous study indicated that theasinensin A has the maximum non-inhibitory concentration at 130–180 mg/mL. Hatano et al. [103] found that theasinensin A suppressed the oxacillin resistance of MRSA and the minimum inhibition concentration (MICs) of oxacillin decreased from 64 to 4 µg/mL. It decreased the MICs of other β-lactam including penicillin G, ampicillin, and streptomycin, antibiotics for MRSA strain [104].

Pathogenesis is closely associated with the ability to synthesize water-insoluble glucans from sucrose by glucosyltransferases (GTases) and to release acids from various fermentable sugars [105]. Oolong tea extract can decrease the cellular surface hydrophobicity of almost all the oral streptococcus and inhibit bacterial adherence to the tooth surfaces, as well as reduce the growth rate and the rate of acid production of mutant streptococci [104]. Oolong tea extract and its purified polymeric polyphenols identified as dehydro-dicatechin A exhibit the inhibition of glucosyltransferase (GTase) of mutants streptococci, *Streptococcus sobrinus* 6715. As the degree of polymerization of catechin increased, GTase was inhibited more effectively [105]. Furthermore, antibacterial activity of oolong tea extract on oral streptococci, including *Streptococcus mutans* and *S. sobrinus* has been confirmed by Sasaki's study who reported that the oolong tea extract have antibacterial activity against all of the oral streptococci examined, with the highest activity against *S. mutans* MT8148R in which the activity was attributed to a monomeric polyphenol-rich fraction. The results also suggested that the antibacterial activity of oolong tea extract was resulted from a synergistic effect of monomeric polyphenols, which can easily bind to proteins [106]. Besides, theasinensin A weakening virus has been reported. For example, herpes simplex virus (HSV), both HSV-1 and HSV-2, is the leading cause of genital ulcers in the developed world [107]. Theasinensin A caused aggregation of HSV-1 glycoprotein B (gB) and the effect is faster than EGCG suggesting that dimers may inhibit the function of viral proteins. This effect is similar to EGCG, on herpes simplex virus (HSV) infectivity. As microbicide agents against HSV, theasinensin A appears to have excellent potential at acidic and neutral pHs [108].

5. Bioavailability

Theasinensins have been reported to have many biological properties closely associated with the antihyperglycemic, anti-obesity, anticancer, anti-inflammatory, antibacterial activity. Usually, biological properties of polyphenols are associated with their bioavailability [109]. Tea polyphenols are biologically regarded as xenobiotics that will be extensively metabolized for

elimination from the body [110]. Metabolism of tea polyphenols can be affected by interaction with other dietary ingredients, solubility, molecular transformations, different cellular transporters and the action of gut microbiota, interindividual variations [110]. The Caco-2 cell monolayers, derived from human colon carcinoma, have been widely used as an *in vitro* intestinal absorption model for studying permeability and transport of drugs [111,112]. It represents morphologically the enterocytes of the small intestine and exhibits brush-border characteristics at the apical side [112].

Qiu et al. [112] illustrated the *in vivo* and *in vitro* absorption of theasinensins A and B and evaluated their transport pathway across intestinal membrane. The rat study showed that a single oral administration of theasinensins demonstrated the intact absorption of theasinensins into the blood system, which was estimated to be a greater than 10-fold lower absorption amount than EGCG. The *in vitro* absorption study indicated that theasinensins can be transported across Caco-2 cell monolayers, while their permeability coefficients were also >10-fold lower than those of EGCG and EGC. In addition, theasinensins were transported across Caco-2 cells in a tight junction (TJ) paracellular diffusion pathway which is the same route as EGCG [112].

6. Future studies of theasinensin

We have known that the concentration of theasinensin A is higher than theasinensin D in black tea [44]. Hashimoto et al. [42] also found that the concentration of theasinensin A is higher than theasinensin B in fermented tea leaves. The average contents of theasinensins in green tea and oolong tea are 0.05% and 0.65%, respectively [36,113]. Data on the content of theasinensins in black or oolong tea are still scarce. Comparison of each isoform of theasinensins between black tea and oolong tea is necessary. Many work are needed to clarify pharmacokinetics and bioactivity of theasinensins both *in vivo* and *in vitro* model. Despite there are studies reporting the potential health benefits of theasinensins, the potential biological activities of their metabolites (conjugates or microbial) is poorly understood. Finally, it is also important to highlight that no studies regarding the mechanism of theasinensins (A–E) on obesity and other bioactivity such as hypertension and atherosclerosis associated with heart disease prevention.

7. Conclusion

In summary, oolong tea and its characteristic compounds theasinensins have been reviewed for chemical formation mechanism of theasinensins and their corresponding biological property and potential action pathway, specifically, the bioactivity of oolong tea and theasinensins and their structural information. These studies clearly indicated the various bioactivities on health benefit of oolong tea coming from theasinensins. Regardless of the polymolecular nature of oolong tea extract, the results shown in the published literatures implicated a relationship between oolong tea and theasinensins and their biological properties. A better knowledge of bioavailability of theasinensins is required. It is foreseeable that theasinensins

may become a promising class of health products against the development of many diseases.

Conflict of interest

The authors have no conflicts of interest to declare.

Acknowledgments

This study was made possible by Naresuan University, Phisanulok, Thailand under the International Research University (IRU) program and Anhui Major Demonstration Project for Leading Talent Team on Tea Chemistry and Health, Anhui Department of Education, Hefei, China.

References

- [1] I. Baldi, D. Gregori, Design considerations on the proof of efficacy of functional foods, *Int. J. Food. Sci. Nutr.* 63 (S1) (2012) 43–50.
- [2] L.M. Yung, F.P. Leung, W.T. Wong, et al., Tea polyphenols benefit vascular function, *Inflammopharmacology* 16 (5) (2008) 230–234.
- [3] N. Harold, P.D. Graham, Green tea composition, consumption and polyphenol chemistry, *Prev. Med.* 21 (1992) 334–350.
- [4] E. Haslam, Thoughts on thearubigins, *Phytochemistry* 64 (2003) 61–73.
- [5] M.G. Sajilata, P.R. Bajaj, R.S. Singhal, Tea polyphenols as nutraceuticals, *Compr. Rev. Food Sci. Food Saf.* 7 (3) (2008) 229–254.
- [6] S. Li, M.H. Pan, C.Y. Lo, C.S. Lai, C.T. Ho, Black tea: chemical analysis and stability, *Food Funct.* 4 (2013) 10–18.
- [7] Y. Chen, J. Duan, Y.M. Jiang, et al., Production, quality, and biological effects of oolong tea (*Camellia sinensis*), *Food Rev. Int.* 27 (2011) 1–15.
- [8] M.S. Pinto, Tea: a new perspective on health benefits, *Food Res. Int.* 53 (2013) 558–567.
- [9] Y. Miyata, S. Tamaru, T. Tanaka, et al., Theflavins and theasinensin A derived from fermented tea have antihyperglycemic and hypotriacylglycerolemic effects in KKA^y mice and Sprague-Dawley rats, *J. Agric. Food Chem.* 61 (2013) 9366–9372.
- [10] L.H. Yao, Y.M. Jiang, N. Caffin, et al., Phenolic compounds in tea from Australian supermarkets, *Food Chem.* 96 (2006) 614–620.
- [11] L. Leung, Y. Su, R. Chen, et al., Theaflavins in black tea and catechins in green tea are equally effective antioxidants, *J. Nutr.* 131 (2001) 2248–2251.
- [12] C. Cabrera, R. Artacho, R. Gimenez, Beneficial effects of green tea – a review, *J. Am. College Nutr.* 25 (2006) 79–99.
- [13] C.S. Yang, J.Y. Chung, G. Yang, et al., Tea and tea polyphenols in cancer prevention, *J. Nutr.* 130 (2000) 472S–478S.
- [14] A.M. Bode, Z. Dong, Signal transduction pathways: targets for green and black tea polyphenols, *J. Biochem. Mol. Biol.* 36 (1) (2002) 66–77.
- [15] China China Stat. (2014).
- [16] G.H. Chen, C.Y. Yang, S.J. Lee, et al., Catechin content and the degree of its galloylation in oolong tea are inversely correlated with cultivation altitude, *J. Food Drug Anal.* 22 (2014) 303–309.
- [17] A. Alcazar, O. Ballesteros, J.M. Jurado, et al., Differentiation of green, white, black, oolong, and Pu-erh teas according to their free amino acids content, *J. Agric. Food Chem.* 55 (2007) 5960–5965.
- [18] G.C. Yen, H.Y. Chen, Comparison of antimutagenic effect of various tea extracts (green, oolong, pouchong, and black tea), *J. Food Prot.* 57 (1) (1994) 54–58.
- [19] N. Matsumoto, T. Kohri, K. Okushio, et al., Inhibitory effects of tea catechins, black tea extract and oolong tea extract on hepatocarcinogenesis in rat, *Jpn. J. Cancer Res.* 87 (1996) 1034–1038.
- [20] T.S. Yam, S. Shah, J.M.T. Hamilton Miller, Microbiological activity of whole and fractionated crude extracts of tea (*Camellia sinensis*), and of tea components, *FEMS Microbiol. Lett.* 152 (1997) 169–174.
- [21] G.Y. Zhang, Y. Miura, K. Yagasaki, Effects of green, oolong and black teas and related components on proliferation and invasion of hepatoma cells, in: Y. Kitagawa, T. Matsuda, S. Iijima (Eds.), *Animal Cell Technology: Basic & Applied Aspects*, vol. 10, Kluwer Academic Publishers, Dordrecht, Netherlands, 1999, pp. 135–139.
- [22] J.V. Higdon, B. Frei, Tea catechins and polyphenols: health effects, metabolism, and antioxidant functions, *Crit. Rev. Food Sci. Nutr.* 43 (2003) 89–143.
- [23] M. Matsumoto, T. Tanaka, M. Maeda, et al., Antibacterial activity of polyphenol components in oolong tea extract against *Streptococcus mutans*, *Caries Res.* 38 (2004) 2–8.
- [24] J.S. Seo, Q. Zhejiu, Y.M. Jeong, Effect of dietary oolong tea extracts on the concentrations of plasma glucose, antioxidant vitamins and minerals in diabetic rats, *FASEB J.* 18 (2004) 519.
- [25] P.H. Tsai, N.B. Kan, S.C. Ho, et al., Effects of oolong tea supplementation on lipid peroxidation of athletes at rest and post-exhaustive exercise, *J. Food Sci.* 70 (2005) S581–S585.
- [26] T.F. Hsu, A. Kusumoto, K. Abe, et al., Polyphenol-enriched oolong tea increases fecal lipid excretion, *Eur. J. Clin. Nutr.* 60 (2006) 1330–1336.
- [27] J.K. Lin, S.Y. Lin-Shiau, Mechanisms of hypolipidemic and anti-obesity effects of tea and tea polyphenols, *Mol. Nutr. Food Res.* 50 (2006) 211–217.
- [28] G. Nonaka, O. Kawahara, I. Nishioka, Tannins and related compounds. XV. A new class of dimeric flavan-3-ol gallates, theasinensins A and B, and proanthocyanidin gallates from green tea leaf, *Chem. Pharm. Bull.* 31 (1983) 3906–3914.
- [29] G. Nonaka, R. Sakai, I. Nishioka, Hydrolysable tannins and proanthocyanidins from green tea, *Phytochemistry* 23 (1984) 1753–1755.
- [30] G. Nonaka, F. Hashimoto, I. Nishioka, Tannins and related compounds. XXXVI. Isolation and structures of theaflagallins, new red pigments from black tea, *Chem. Pharm. Bull.* 34 (1986) 61–65.
- [31] F. Hashimoto, G. Nonaka, I. Nishioka, Tannins and related compounds. LVI. Isolation of four new acylated flavan-3-ols from oolong tea, *Chem. Pharm. Bull.* 35 (1987) 611–616.
- [32] Y. Ikeya, H. Taguchi, I. Yosioka, et al., The constituents of *Schizandra chinensis* Baill. I. Isolation and structure determination of five new lignans, gomisin A, B, C, F and G, and the absolute structure of schizandrin, *Chem. Pharm. Bull.* 27 (1979) 1383.
- [33] F. Hashimoto, G. Nonaka, I. Nishioka, Tannins and related compounds LXIX. Isolation and structure elucidation of B,B-linked bisflavanoids, theasinensins D–G and oolongtheanine from oolong tea, *Chem. Pharm. Bull.* 36 (1988) 1676–1684.
- [34] E.A.H. Robert, The phenolic substances of manufactured tea IV. Enzymic oxidations of individual substances, *J. Sci. Food Agric.* 10 (1959) 167–172.
- [35] T. Tanaka, C. Mine, S. Watarumi, et al., Accumulation of epigallocatechin quinone dimers during tea fermentation and formation of theasinensins, *J. Nat. Prod.* 65 (2002) 1582–1587.
- [36] U.H. Engellhardt, Chemistry of tea. Comprehensive natural products II, *Chem. Biol.* 3 (2010) 999–1032.
- [37] T. Tanaka, S. Watarumi, Y. Matsuo, et al., Production of theasinensins A and D, epigallocatechin gallate dimers of black tea, by oxidation–reduction dismutation of dehydrotheasinensin A, *Tetrahedron* 59 (2003) 7939–7947.
- [38] Y. Matsuo, T. Tanaka, I. Kouno, A new mechanism for oxidation of epigallocatechin and production of benzotropolone pigments, *Tetrahedron* 62 (2006) 4774–4783.
- [39] A. Ferretti, V.P. Flanagan, H.A. Bondarovich, et al., The chemistry of tea. Structures of compounds A and B of Roberts and reactions of some model compounds, *J. Agric. Food Chem.* 16 (5) (1968) 756–761.
- [40] F. Shahidi, M. Nacz, Phenolics in Food and Nutraceuticals, CRC Press, Boca Raton, FL, USA, 2004, pp. 241–250.
- [41] C.T. Ho, J.K. Lin, F. Shahidi, Tea and Tea Products, Chemistry and Health-Promoting Properties, CRC Press, Boca Raton, 2009.
- [42] F. Hashimoto, G. Nonaka, I. Nishioka, Tannins related compounds CXIV. Structure of novel fermentation products, theogallin, theaflavinonin and desgalloyl theaflavinonin from black tea, and changes of tea leaf polyphenols during fermentation, *Chem. Pharm. Bull.* 40 (1992) 1383–1389.
- [43] T. Shii, M. Miyamoto, Y. Matsuo, et al., Biomimetic one-pot preparation of a black tea polyphenol theasinensin A from epigallocatechin gallate

- by treatment with copper(II) chloride and ascorbic acid, *Chem. Pharm. Bull.* 59 (9) (2011) 1183–1185.
- [44] T. Tanaka, Y. Matsuo, I. Kouno, Chemistry of secondary polyphenols produced during processing of tea and selected foods, *Int. J. Mol. Sci.* 11 (2010) 14–40.
- [45] T. Tanaka, C. Mine, K. Inoue, et al., Synthesis of theaflavin from epicatechin and epigallocatechin by plant homogenates and role of epicatechin quinone in the synthesis and degradation of theaflavin, *J. Agric. Food Chem.* 50 (2002) 2142–2148.
- [46] Y. Matsuo, Y. Yamada, T. Tanaka, et al., Enzymatic oxidation of gallic acid and epigallocatechin: effects of C-ring configuration on the reaction products, *Phytochemistry* 69 (2008) 3054–3061.
- [47] Y. Takino, H. Imagawa, H. Horikawa, et al., Study on the mechanism of the oxidation of tea leaf catechins, Part III. Formation a reddish orange pigment and its spectral relationship to some benzotropolone derivatives, *Agric. Biol. Chem.* 28 (1964) 64–71.
- [48] Y. Takino, A. Ferretti, V. Flanagan, et al., The structure of theaflavin, A polyphenol of black tea, *Tetrahedron Lett.* 6 (1965) 4019–4025.
- [49] Y. Matsuo, T. Tanaka, I. Kouno, Production mechanism of proepitheafllagallin, a precursor of benzotropolone-type black tea pigment, derived from epigallocatechin via a bicycle [3.2.1] octane-type intermediate, *Tetrahedron Lett.* 50 (2009) 1348–1351.
- [50] T. Tanaka, S. Watarumi, M. Fujieda, et al., New black tea polyphenol having N-ethyl-2-pyrrolidinone moiety derived from tea amino acid theanine: isolation, characterization and partial synthesis, *Food Chem.* 93 (2005) 81–87.
- [51] R. Saijo, T. Takeo, Formation of aldehydes from amino acids by tea leaves extracts, *Agric. Biol. Chem.* 34 (1970) 227–233.
- [52] G.P. Rizzi, Formation of Strecker aldehydes from polyphenol-derived quinones and α -amino acids in a nonenzymic model system, *J. Agric. Food Chem.* 54 (2006) 1893–1897.
- [53] Y. Li, T. Tanaka, I. Kouno, Oxidative coupling of the pyrogallol B-ring with a galloyl group during enzymatic oxidation of epigallocatechin 3-O-gallate, *Phytochemistry* 68 (2007) 1081–1088.
- [54] M.H. Pan, C.S. Lai, H. Wang, C.Y. Lo, C.T. Ho, S. Li, Black tea in chemoprevention of cancer and other human diseases, *Food Sci. Hum. Wellness* 2 (2013) 13–22.
- [55] M.G. Hertog, E.J.M. Feskens, P.C.H. Hollman, et al., Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly study, *Lancet* 342 (1993) 1007–1011.
- [56] C.S. Yang, Z.Y. Wang, Tea and cancer, *J. Natl. Cancer Inst.* 85 (1993) 1038–1049.
- [57] J.M. Geleijnse, L.J. Launer, A. Hofman, et al., Tea flavonoids may protect against atherosclerosis. The Rotterdam Study, *Arch. Intern. Med.* 159 (18) (1999) 2170–2174.
- [58] I.C.W. Arts, P.C.H. Hollman, E.J.M. Feskens, et al., Catechin intake might explain the inverse relation between tea consumption and ischemic heart disease: the Zutphen Elderly Study, *Am. J. Clin Nutr.* 74 (2001) 227–232.
- [59] G.C. Yen, H.Y. Chen, Antioxidant activity of various tea extracts in relation to their antimutagenicity, *J. Agric. Food Chem.* 43 (1995) 27–32.
- [60] C. Rice-Evans, Implications of the mechanisms of action of tea polyphenols as antioxidants in vitro for chemoprevention in humans, *Proc. Soc. Exp. Biol. Med.* 220 (1999) 262–266.
- [61] H.C. Lee, A.M. Jenner, C.S. Low, et al., Effect of tea phenolics and their aromatic fecal bacterial metabolites on intestinal microbiota, *Res. Microbiol.* 157 (2006) 876–884.
- [62] Y.S. Cho, N.L. Schiller, H.Y. Kahng, et al., Cellular responses and proteomic analysis of *Escherichia coli* exposed to green tea polyphenols, *Curr. Microbiol.* 55 (2007) 501–506.
- [63] I.F. Benzie, Y.T. Szeto, Total antioxidant capacity of teas by the ferric reducing/antioxidant power assay, *J. Agric. Food Chem.* 47 (1999) 633–636.
- [64] F. Hashimoto, M. Ono, C. Masuoka, et al., Evaluation of the antioxidative effect (in vitro) of tea polyphenols, *Biosci. Biotechnol. Biochem.* 67 (2003) 396–401.
- [65] D. Heber, Y. Zhang, J. Yang, et al., Green tea, black tea, and oolong tea polyphenols reduce visceral fat and inflammation in mice fed high-fat, high-sucrose obesogenic diets, *J. Nutr.* 144 (9) (2014) 1385–1393.
- [66] K. Sun, I. Wernstedt Asterholm, C.M. Kusminski, et al., Dichotomous effects of VEGF-A on adipose tissue dysfunction, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) 5874–5879.
- [67] A. Tchernof, J.P. Despres, Pathophysiology of human visceral obesity: an update, *Physiol. Rev.* 93 (2013) 359–404.
- [68] D.X. Hou, S. Masuzaki, S. Tanigawa, et al., Oolong tea theasinensins attenuate cyclooxygenase-2 expression in lipopolysaccharide (LPS)-activated mouse macrophages: structure–activity relationship and molecular mechanisms, *J. Agric. Food Chem.* 58 (2010) 12735–12743.
- [69] J. Chen, S. Qin, J. Xiao, et al., A genome-wide microarray highlights the antiinflammatory genes targeted by oolong tea theasinensin A in macrophages, *Nutr. Cancer* 63 (7) (2011) 1064–1073.
- [70] A. Hisanaga, H. Ishida, K. Sakao, et al., Anti-inflammatory activity and molecular mechanism of oolong tea theasinensin, *Food Funct.* 5 (2014) 1891–1897.
- [71] F. Carriere, C. Renou, S. Ransac, et al., Inhibition of gastrointestinal lipolysis by Orlistat during digestion of test meals in healthy volunteers, *Am. J. Physiol. Gastrointest. Liver Physiol.* 281 (2001) G16–G28.
- [72] L.K. Han, T. Takaku, J. Li, et al., Anti-obesity action of oolong tea, *Int. J. Obes.* 23 (1999) 98–105.
- [73] M. Nakai, Y. Fukui, S. Asami, et al., Inhibitory effects of oolong tea polyphenols on pancreatic lipase in vitro, *J. Agric. Food Chem.* 53 (2005) 4593–4598.
- [74] K.L. Kuo, M.S. Weng, C.T. Chiang, et al., Comparative studies on the hypolipidemic and growth suppressive effects of oolong, black, pu-erh, and green tea leaves in rats, *Agric. Food Chem.* 53 (2005) 480–489.
- [75] J. Panee, Monocyte chemoattractant protein 1 (MCP-1) in obesity and diabetes, *Cytokine* 60 (2012) 1–12.
- [76] J.H. Weisburger, F.L. Chung, Mechanisms of chronic disease causation by nutritional factors and tobacco products and their prevention by tea polyphenols, *Food Chem. Toxicol.* 40 (2002) 1145–1154.
- [77] L.M. Coussens, Z. Werb, Inflammation and cancer, *Nature* 420 (2002) 860–867.
- [78] T.C. Hour, Y.C. Liang, I.S. Chu, et al., Inhibition of eleven mutagens by various tea extracts, (–)-epigallocatechin-3-gallate, gallic acid and caffeine, *Food Chem. Toxicol.* 37 (1999) 569–579.
- [79] G.C. Yen, H.Y. Chen, Relationship between antimutagenic activity and major components of various teas, *Mutagenesis* 11 (1996) 37–41.
- [80] T. Ohe, K. Marutani, S. Nakase, Catechins are not major components responsible for antigenotoxic effects of tea extracts against nitroarenes, *Mutat. Res. Genet. Toxicol. Environ. Mutagen.* 496 (2001) 75–81.
- [81] G.Y. Zhang, Y. Miura, K. Yagasaki, Induction of apoptosis and cell cycle arrest in cancer cells by in vivo metabolites of teas, *Nutr. Cancer* 38 (2000) 265–273.
- [82] H. Hibasami, Z.X. Jin, M. Hasegawa, et al., Oolong tea polyphenol extract induces apoptosis in human stomach cancer cells, *Anticancer Res.* 20 (2000) 4403–4406.
- [83] K. Saeki, S. Hayakawa, M. Isemura, et al., Importance of a pyrogallol-type structure in catechin compounds for apoptosis-inducing activity, *Phytochemistry* 53 (3) (2000) 391–394.
- [84] M. Maeda-Yamamoto, H. Kawahara, N. Tahara, et al., Effects of tea polyphenols on the invasion and matrix metalloproteinases activities of human fibrosarcoma HT1080 cells, *J. Agric. Food Chem.* 47 (1999) 2350–2354.
- [85] M.H. Pan, Y.C. Liang, S.Y. Lin-Shiau, et al., Induction of apoptosis by the oolong tea polyphenol theasinensin A through cytochrome *c* release and activation of caspase-9 and caspase-3 in human U937 cells, *J. Agric. Food Chem.* 48 (2000) 6337–6346.
- [86] S.J. Hossain, H. Kato, H. Aoshima, et al., Polyphenol induced inhibition of the response of Na⁺/glucose cotransporter expressed in xenopus oocytes, *J. Agric. Food Chem.* 50 (2002) 5215–5219.
- [87] R.A. Anderson, M.M. Polansky, Tea enhances insulin activity, *J. Agric. Food Chem.* 50 (2002) 7182–7186.
- [88] K. Hosoda, M.F. Wanf, M.L. Liao, et al., Antihyperglycemic effect of oolong tea in type 2 diabetes, *Diabetes Care* 26 (2003) 1714–1718.
- [89] T. Matsui, T. Tanaka, S. Tamura, et al., Alpha-glucosidase inhibitory profile of catechins and theaflavins, *J. Agric. Food Chem.* 55 (2007) 99–105.

- [90] J. Qiu, K. Maekawa, Y. Kitamura, et al., Stimulation of glucose uptake by theasinensins through the AMP-activated protein kinase pathway in rat skeletal muscle cells, *Biochem. Pharmacol.* 87 (2014) 344–351.
- [91] H.G. Joost, G.I. Bell, J.D. Best, et al., Nomenclature of the GLUT/SLC2A family of sugar/polyol transport facilitators, *Am. J. Physiol. Endocrinol. Metab.* 282 (2002) 974–976.
- [92] J. Moline, I.F. Bukharovich, M.S. Wolff, et al., Dietary flavonoids and hypertension: is there a link? *Med. Hypotheses* 55 (2000) 306–309.
- [93] M. Safar, P. Chamiot-Clerc, G. Dagher, et al., Pulse pressure, endothelium function, and arterial stiffness in spontaneously hypertensive rats, *Hypertension* 38 (2001) 1416–1421.
- [94] Y.C. Yang, F.H. Lu, J.S. Wu, et al., The protective effect of habitual tea consumption on hypertension, *Arch. Int. Med.* 164 (2004) 1534–1540.
- [95] C.D. Gardner, S.P. Fortmann, R.M. Krauss, Association of small low-density lipoprotein particles with the incidence of coronary artery disease in men and women, *JAMA* 276 (1996) 875–881.
- [96] M.J. Stampfer, R.M. Krauss, J. Ma, et al., A prospective study of triglyceride level, low-density lipoprotein particle diameter, and risk of myocardial infarction, *JAMA* 276 (1996) 882–888.
- [97] K. Shimada, T. Kawarabayashi, A. Tanaka, et al., Oolong tea increases plasma adiponectin levels and low-density lipoprotein particle size in patients with coronary artery disease, *Diabetes Res. Clin. Pract.* 65 (2004) 227–234.
- [98] T.F. Hsu, A. Kusumoto, K. Abe, et al., Polyphenol enriched oolong tea increases fecal lipid excretion, *Eur. J. Clin. Nutr.* 60 (2006) 1330–1336.
- [99] T. Komatsu, M. Nakamori, K. Komatsu, et al., Oolong tea increases energy metabolism in Japanese females, *J. Med. Investig.* 50 (2003) 170–175.
- [100] W. Rumpfer, J.J. Seale, B. Clevidence, et al., Oolong tea increases metabolic rate and fat oxidation in men, *J. Nutr.* 131 (2001) 2848–2852.
- [101] T.T. Yang, M.W. Koo, Hypercholesterolemic effects of Chinese tea, *Pharmacol. Res.* 35 (1997) 505–512.
- [102] R. Monina Kleven, J.R. Edwards, F.C. Tenover, et al., National nosocomial infections surveillance system changes in the epidemiology of methicillin-resistant *Staphylococcus aureus* in intensive care units in US hospitals, 1992–2003, *Clin. Infect. Dis.* 42 (2006) 389–391.
- [103] T. Hatano, M. Kusuda, M. Hori, et al., Thesinesin A, a tea polyphenol formed from (–)-epigallocatechin gallate, suppresses antibiotic resistance of methicillin-resistant *Staphylococcus aureus*, *Planta Med.* 69 (2003) 984–989.
- [104] C.C. Chou, L.L. Lin, K.T. Chung, Antimicrobial activity of tea as affected by the degree of fermentation and manufacturing season, *Int. J. Food Microbiol.* 48 (1999) 125–130.
- [105] S. Hamada, M. Kontani, H. Hosono, et al., Peroxidase-catalyzed generation of catechin oligomers that inhibit glucosyltransferase from *Streptococcus sobrinus*, *FEMS Microbiol. Lett.* 143 (1996) 35–40.
- [106] H. Sasaki, M. Matsumoto, T. Tanaka, et al., Antibacterial activity of polyphenol components in oolong tea extract against *Streptococcus mutans*, *Caries Res.* 38 (2004) 2–8.
- [107] A. Mindel, Genital herpes—how much of a public-health problem? *Lancet* 351 (1998) 16–18.
- [108] C.E. Isaacs, W. Xu, G. Merz, et al., Digallate dimers of (–)-epigallocatechin gallate inactivate herpes simplex virus, *Antimicrob. Agents Chemother.* 55 (12) (2011) 5646–5653.
- [109] A. Scalbert, G. Williamson, Dietary intake and bioavailability of polyphenols, *J. Nutr.* 130 (2000) 2073S–2085S.
- [110] A.P. Neilson, M.G. Ferruzzi, Influence of formulation and processing on absorption and metabolism of flavan-3-ols from tea and cocoa, *Annu. Rev. Food Sci. Technol.* 2 (2011) 125–151.
- [111] P. Artursson, K. Palma, K. Luthman, Caco-2 monolayers in experimental and theoretical predictions of drug transport, *Adv. Drug Deliv. Rev.* 46 (1996) 27–43.
- [112] J. Qiu, Y. Kitamura, Y. Miyata, et al., Transepithelial transport of theasinensins through caco-2 cell monolayers and their absorption in Sprague-Dawley rats after oral administration, *J. Agric. Food Chem.* 60 (2012) 8036–8043.
- [113] U. Stodt, U.H. Engelhardt, Progress in the analysis of selected tea constituents over the past 20 years, *Food Res. Int.* 53 (2013) 636–648.
- [114] B.J. Xie, H. Shi, Q.Y. Chen, et al., Antioxidant properties of fractions and polyphenol constituents from green, oolong, and black teas, *Proc. Natl. Sci. Council Rep. China [B]* 17 (1993) 77–84.